

# Achim A Jungbluth

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6677931/publications.pdf>

Version: 2024-02-01

15

papers

3,237

citations

623734

14

h-index

996975

15

g-index

15

all docs

15

docs citations

15

times ranked

4147

citing authors

#	ARTICLE	IF	CITATIONS
1	Cancer/testis antigens, gametogenesis and cancer. <i>Nature Reviews Cancer</i> , 2005, 5, 615-625.	28.4	1,415
2	Immunohistochemical analysis of NY-ESO-1 antigen expression in normal and malignant human tissues. <i>International Journal of Cancer</i> , 2001, 92, 856-860.	5.1	310
3	PRAME Expression in Melanocytic Tumors. <i>American Journal of Surgical Pathology</i> , 2018, 42, 1456-1465.	3.7	248
4	A103. <i>American Journal of Surgical Pathology</i> , 1998, 22, 595-602.	3.7	233
5	Monophasic and biphasic synovial sarcomas abundantly express cancer/testis antigen ny-eso-1 but not mage-a1 or ct7. <i>International Journal of Cancer</i> , 2001, 94, 252-256.	5.1	182
6	Expression of MAGE-antigens in normal tissues and cancer. <i>International Journal of Cancer</i> , 2000, 85, 460-465.	5.1	179
7	The cancer-testis antigens CT7 (MAGE-C1) and MAGE-A3/6 are commonly expressed in multiple myeloma and correlate with plasma-cell proliferation. <i>Blood</i> , 2005, 106, 167-174.	1.4	172
8	A monoclonal antibody recognizing human cancers with amplification/overexpression of the human epidermal growth factor receptor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 639-644.	7.1	158
9	NY-ESO-1 is a ubiquitous immunotherapeutic target antigen for patients with myxoid/round cell liposarcoma. <i>Cancer</i> , 2012, 118, 4564-4570.	4.1	107
10	T311â€”An Anti-Tyrosinase Monoclonal Antibody for the Detection of Melanocytic Lesions in Paraffin Embedded Tissues. <i>Pathology Research and Practice</i> , 2000, 196, 235-242.	2.3	73
11	CT7 (MAGE-C1) antigen expression in normal and neoplastic tissues. <i>International Journal of Cancer</i> , 2002, 99, 839-845.	5.1	60
12	Cancer/testis (CT) antigens are expressed in fetal ovary. <i>Cancer Immunity</i> , 2007, 7, 1.	3.2	35
13	POU2F3 in SCLC: Clinicopathologic and Genomic Analysis With a Focus on Its Diagnostic Utility in Neuroendocrine-Low SCLC. <i>Journal of Thoracic Oncology</i> , 2022, 17, 1109-1121.	1.1	29
14	MAGE antigen expression in monophasic and biphasic synovial sarcoma. <i>Human Pathology</i> , 2002, 33, 225-229.	2.0	22
15	Immunohistochemical Detection of Cancer-Testis Antigen PRAME. <i>International Journal of Surgical Pathology</i> , 2021, 29, 826-835.	0.8	14