

P Monica Lind

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6677973/p-monica-lind-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

115
papers

3,583
citations

36
h-index

54
g-index

134
ext. papers

4,104
ext. citations

7.1
avg. IF

5.47
L-index

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 115 | Serum levels of perfluoroalkyl substances (PFAS) and body composition - A cross-sectional study in a middle-aged population.. <i>Environmental Research</i> , 2022 , 209, 112677 | 7.9 | 0 |
| 114 | Obesity II: Establishing Causal Links Between Chemical Exposures and Obesity.. <i>Biochemical Pharmacology</i> , 2022 , 115015 | 6 | 6 |
| 113 | Changes in plasma levels of per- and polyfluoroalkyl substances (PFAS) are associated with changes in plasma lipids - A longitudinal study over 10 years.. <i>Environmental Research</i> , 2022 , 112903 | 7.9 | 1 |
| 112 | High Serum-Induced AhRL Is Associated with Prevalent Metabolic Syndrome and Future Impairment of Glucose Tolerance in the Elderly. <i>Endocrinology and Metabolism</i> , 2021 , 36, 436-446 | 3.5 | 1 |
| 111 | Concentrations of nine endogenous steroid hormones in 70-year-old men and women. <i>Endocrine Connections</i> , 2021 , 10, 511-520 | 3.5 | 2 |
| 110 | Lipophilic Environmental Chemical Mixtures Released During Weight-Loss: The Need to Consider Dynamics. <i>BioEssays</i> , 2020 , 42, e1900237 | 4.1 | 4 |
| 109 | Are Persistent Organic Pollutants Linked to Lipid Abnormalities, Atherosclerosis and Cardiovascular Disease? A Review. <i>Journal of Lipid and Atherosclerosis</i> , 2020 , 9, 334-348 | 3 | 5 |
| 108 | The associations between p,pSDDE levels and plasma levels of lipoproteins and their subclasses in an elderly population determined by analysis of lipoprotein content. <i>Lipids in Health and Disease</i> , 2020 , 19, 249 | 4.4 | 3 |
| 107 | Association of Exposure to Persistent Organic Pollutants With Mortality Risk: An Analysis of Data From the Prospective Investigation of Vasculature in Uppsala Seniors (PIVUS) Study. <i>JAMA Network Open</i> , 2019 , 2, e193070 | 10.4 | 16 |
| 106 | Developmental low-dose exposure to bisphenol A induces chronic inflammation, bone marrow fibrosis and reduces bone stiffness in female rat offspring only. <i>Environmental Research</i> , 2019 , 177, 108584 | 7.9 | 16 |
| 105 | Developmental exposure to a very low dose of bisphenol A induces persistent islet insulin hypersecretion in Fischer 344 rat offspring. <i>Environmental Research</i> , 2019 , 172, 127-136 | 7.9 | 18 |
| 104 | Urinary bisphenol A and serum lipids: a meta-analysis of six NHANES examination cycles (2003-2014). <i>Journal of Epidemiology and Community Health</i> , 2019 , 73, 1012-1019 | 5.1 | 11 |
| 103 | Identification of metabolic profiles associated with human exposure to perfluoroalkyl substances. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2019 , 29, 196-205 | 6.7 | 36 |
| 102 | DDT and its metabolites could contribute to the aetiology of chronic kidney disease of unknown aetiology (CKDu) and more studies are a priority. <i>Science of the Total Environment</i> , 2019 , 649, 1638-1639 | 10.2 | 2 |
| 101 | A method for analysis of marker persistent organic pollutants in low-volume plasma and serum samples using 96-well plate solid phase extraction. <i>Journal of Chromatography A</i> , 2018 , 1546, 18-27 | 4.5 | 20 |
| 100 | Circulating levels of perfluoroalkyl substances and left ventricular geometry of the heart in the elderly. <i>Environment International</i> , 2018 , 115, 295-300 | 12.9 | 4 |
| 99 | High plasma organochlorine pesticide levels are related to increased biological age as calculated by DNA methylation analysis. <i>Environment International</i> , 2018 , 113, 109-113 | 12.9 | 18 |

| | | | |
|----|---|------|----|
| 98 | Neurotoxic chemicals in adipose tissue: A role in puzzling findings on obesity and dementia. <i>Neurology</i> , 2018 , 90, 176-182 | 6.5 | 12 |
| 97 | The association between p,pSDDE levels and left ventricular mass is mainly mediated by obesity. <i>Environmental Research</i> , 2018 , 160, 541-546 | 7.9 | 7 |
| 96 | Studies of indirect and direct effects of hypervitaminosis A on rat bone by comparing free access to food and pair-feeding. <i>Uppsala Journal of Medical Sciences</i> , 2018 , 123, 82-85 | 2.8 | 5 |
| 95 | Longitudinal changes in persistent organic pollutants (POPs) from 2001 to 2009 in a sample of elderly Swedish men and women. <i>Environmental Research</i> , 2018 , 165, 193-200 | 7.9 | 12 |
| 94 | Prenatal Bisphenol A Exposure is Linked to Epigenetic Changes in Glutamate Receptor Subunit Gene Grin2b in Female Rats and Humans. <i>Scientific Reports</i> , 2018 , 8, 11315 | 4.9 | 24 |
| 93 | Changes in plasma levels of perfluoroalkyl substances (PFASs) are related to increase in carotid intima-media thickness over 10 years - a longitudinal study. <i>Environmental Health</i> , 2018 , 17, 59 | 6 | 11 |
| 92 | High serum levels of p,pSDDE are associated with an accelerated decline in GFR during 10 years follow-up. <i>Science of the Total Environment</i> , 2018 , 644, 371-374 | 10.2 | 9 |
| 91 | Changes in markers of liver function in relation to changes in perfluoroalkyl substances - A longitudinal study. <i>Environment International</i> , 2018 , 117, 196-203 | 12.9 | 36 |
| 90 | Endocrine-disrupting chemicals and risk of diabetes: an evidence-based review. <i>Diabetologia</i> , 2018 , 61, 1495-1502 | 10.3 | 69 |
| 89 | Low-dose exposure to Bisphenol A during development has limited effects on male reproduction in midpubertal and aging Fischer 344 rats. <i>Reproductive Toxicology</i> , 2018 , 81, 196-206 | 3.4 | 8 |
| 88 | Low-dose developmental bisphenol A exposure alters fatty acid metabolism in Fischer 344 rat offspring. <i>Environmental Research</i> , 2018 , 166, 117-129 | 7.9 | 18 |
| 87 | Population attributable risks and costs of diabetogenic chemical exposures in the elderly. <i>Journal of Epidemiology and Community Health</i> , 2017 , 71, 111-114 | 5.1 | 13 |
| 86 | Dismissing manufactured uncertainties, limitations and competing interpretations about chemical exposures and diabetes. <i>Journal of Epidemiology and Community Health</i> , 2017 , 71, 942 | 5.1 | 1 |
| 85 | Effects of Low-Dose Developmental Bisphenol A Exposure on Metabolic Parameters and Gene Expression in Male and Female Fischer 344 Rat Offspring. <i>Environmental Health Perspectives</i> , 2017 , 125, 067018 | 8.4 | 40 |
| 84 | Relationships between serum-induced AhR bioactivity or mitochondrial inhibition and circulating polychlorinated biphenyls (PCBs). <i>Scientific Reports</i> , 2017 , 7, 9383 | 4.9 | 12 |
| 83 | Low-dose developmental exposure to bisphenol A induces sex-specific effects in bone of Fischer 344 rat offspring. <i>Environmental Research</i> , 2017 , 159, 61-68 | 7.9 | 23 |
| 82 | The effect of drinking water contaminated with perfluoroalkyl substances on a 10-year longitudinal trend of plasma levels in an elderly Uppsala cohort. <i>Environmental Research</i> , 2017 , 159, 95-102 | 7.9 | 19 |
| 81 | Mixture effects of 30 environmental contaminants on incident metabolic syndrome-A prospective study. <i>Environment International</i> , 2017 , 107, 8-15 | 12.9 | 15 |

| | | | |
|----|---|------|----|
| 80 | Genetic and methylation variation in the CYP2B6 gene is related to circulating p,pSdde levels in a population-based sample. <i>Environment International</i> , 2017 , 98, 212-218 | 12.9 | 4 |
| 79 | Serum levels of brominated flame retardants (BFRs: PBDE, HBCD) and influence of dietary factors in a population-based study on Swedish adults. <i>Chemosphere</i> , 2017 , 167, 485-491 | 8.4 | 41 |
| 78 | Circulating levels of perfluoroalkyl substances (PFASs) and carotid artery atherosclerosis. <i>Environmental Research</i> , 2017 , 152, 157-164 | 7.9 | 25 |
| 77 | Low-dose exposure to bisphenol A in combination with fructose increases expression of genes regulating angiogenesis and vascular tone in juvenile Fischer 344 rat cardiac tissue. <i>Uppsala Journal of Medical Sciences</i> , 2017 , 122, 20-27 | 2.8 | 14 |
| 76 | Association between background exposure to organochlorine pesticides and the risk of cognitive impairment: A prospective study that accounts for weight change. <i>Environment International</i> , 2016 , 89-90, 179-84 | 12.9 | 37 |
| 75 | Altered heart proteome in fructose-fed Fisher 344 rats exposed to bisphenol A. <i>Toxicology</i> , 2016 , 347-349, 6-16 | 4.4 | 17 |
| 74 | The metabolic fingerprint of p,pSDDE and HCB exposure in humans. <i>Environment International</i> , 2016 , 88, 60-66 | 12.9 | 47 |
| 73 | Uppsala Consensus Statement on Environmental Contaminants and the Global Obesity Epidemic. <i>Environmental Health Perspectives</i> , 2016 , 124, A81-3 | 8.4 | 27 |
| 72 | Circulating levels of perfluoroalkyl substances are associated with dietary patterns - A cross sectional study in elderly Swedish men and women. <i>Environmental Research</i> , 2016 , 150, 59-65 | 7.9 | 22 |
| 71 | Changes in serum levels of perfluoroalkyl substances during a 10-year follow-up period in a large population-based cohort. <i>Environment International</i> , 2016 , 95, 86-92 | 12.9 | 37 |
| 70 | Perfluoroalkyl substances (PFAS) including structural PFOS isomers in plasma from elderly men and women from Sweden: Results from the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS). <i>Environment International</i> , 2015 , 82, 21-7 | 12.9 | 22 |
| 69 | Expression of the aryl hydrocarbon receptor in growth plate cartilage and the impact of its local modulation on longitudinal bone growth. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 8059-69 | 6.3 | 5 |
| 68 | Genome-wide association study of plasma levels of polychlorinated biphenyls disclose an association with the CYP2B6 gene in a population-based sample. <i>Environmental Research</i> , 2015 , 140, 95-101 | 7.9 | 10 |
| 67 | Genome-wide association study of toxic metals and trace elements reveals novel associations. <i>Human Molecular Genetics</i> , 2015 , 24, 4739-45 | 5.6 | 75 |
| 66 | Manufacturing doubt about endocrine disrupter science--A rebuttal of industry-sponsored critical comments on the UNEP/WHO report "State of the Science of Endocrine Disrupting Chemicals 2012". <i>Regulatory Toxicology and Pharmacology</i> , 2015 , 73, 1007-17 | 3.4 | 46 |
| 65 | Circulating levels of environmental contaminants are associated with dietary patterns in older adults. <i>Environment International</i> , 2015 , 75, 93-102 | 12.9 | 13 |
| 64 | Circulating levels of perfluoroalkyl substances and prevalent diabetes in the elderly. <i>Diabetologia</i> , 2014 , 57, 473-9 | 10.3 | 83 |
| 63 | Exposure to bisphenol A affects lipid metabolism in <i>Drosophila melanogaster</i> . <i>Basic and Clinical Pharmacology and Toxicology</i> , 2014 , 114, 414-20 | 3.1 | 19 |

| | | | |
|----|---|------|----|
| 62 | Persistent organic pollutants are related to the change in circulating lipid levels during a 5 year follow-up. <i>Environmental Research</i> , 2014 , 134, 190-7 | 7.9 | 31 |
| 61 | Persistent organic pollutants and liver dysfunction biomarkers in a population-based human sample of men and women. <i>Environmental Research</i> , 2014 , 134, 251-6 | 7.9 | 31 |
| 60 | The identification of complex interactions in epidemiology and toxicology: a simulation study of boosted regression trees. <i>Environmental Health</i> , 2014 , 13, 57 | 6 | 32 |
| 59 | Serum levels of monobenzylphthalate (MBzP) is related to carotid atherosclerosis in the elderly. <i>Environmental Research</i> , 2014 , 133, 348-52 | 7.9 | 15 |
| 58 | Influence of persistent organic pollutants on the complement system in a population-based human sample. <i>Environment International</i> , 2014 , 71, 94-100 | 12.9 | 20 |
| 57 | Genetic variation in the CYP2B6 gene is related to circulating 2,2,5,4,4-tetrabromodiphenyl ether (BDE-47) concentrations: an observational population-based study. <i>Environmental Health</i> , 2014 , 13, 34 | 6 | 8 |
| 56 | Bisphenol A is related to circulating levels of adiponectin, leptin and ghrelin, but not to fat mass or fat distribution in humans. <i>Chemosphere</i> , 2014 , 112, 42-8 | 8.4 | 47 |
| 55 | Circulating levels of p,p-DDDE are related to prevalent hypertension in the elderly. <i>Environmental Research</i> , 2014 , 129, 27-31 | 7.9 | 42 |
| 54 | Genetic variation in the CYP1A1 gene is related to circulating PCB118 levels in a population-based sample. <i>Environmental Research</i> , 2014 , 133, 135-40 | 7.9 | 10 |
| 53 | Influence of persistent organic pollutants on oxidative stress in population-based samples. <i>Chemosphere</i> , 2014 , 114, 303-9 | 8.4 | 27 |
| 52 | Does mortality risk of cigarette smoking depend on serum concentrations of persistent organic pollutants? Prospective investigation of the vasculature in Uppsala seniors (PIVUS) study. <i>PLoS ONE</i> , 2014 , 9, e95937 | 3.7 | 8 |
| 51 | Persistent organic pollutants and inflammatory markers in a cross-sectional study of elderly Swedish people: the PIVUS cohort. <i>Environmental Health Perspectives</i> , 2014 , 122, 977-83 | 8.4 | 37 |
| 50 | Whole blood and serum concentrations of metals in a Swedish population-based sample. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 143-8 | 2 | 53 |
| 49 | An environmental wide association study (EWAS) approach to the metabolic syndrome. <i>Environment International</i> , 2013 , 55, 1-8 | 12.9 | 49 |
| 48 | A rapid method for the determination of perfluoroalkyl substances including structural isomers of perfluorooctane sulfonic acid in human serum using 96-well plates and column-switching ultra-high performance liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2013 , 1305, 164-70 | 4.5 | 46 |
| 47 | Circulating levels of persistent organic pollutants are related to retrospective assessment of life-time weight change. <i>Chemosphere</i> , 2013 , 90, 998-1004 | 8.4 | 17 |
| 46 | Bisphenol A exposure increases liver fat in juvenile fructose-fed Fischer 344 rats. <i>Toxicology</i> , 2013 , 303, 125-32 | 4.4 | 43 |
| 45 | Global DNA hypermethylation is associated with high serum levels of persistent organic pollutants in an elderly population. <i>Environment International</i> , 2013 , 59, 456-61 | 12.9 | 63 |

| | | | |
|----|--|------|-----|
| 44 | Circulating levels of persistent organic pollutants (POPs) are associated with left ventricular systolic and diastolic dysfunction in the elderly. <i>Environmental Research</i> , 2013 , 123, 39-45 | 7.9 | 25 |
| 43 | Circulating levels of persistent organic pollutants in relation to visceral and subcutaneous adipose tissue by abdominal MRI. <i>Obesity</i> , 2013 , 21, 413-8 | 8 | 39 |
| 42 | Persistent organic pollutants and abnormal geometry of the left ventricle in the elderly. <i>Journal of Hypertension</i> , 2013 , 31, 1547-53 | 1.9 | 14 |
| 41 | Quantification of total and visceral adipose tissue in fructose-fed rats using water-fat separated single echo MRI. <i>Obesity</i> , 2013 , 21, E388-95 | 8 | 4 |
| 40 | Circulating levels of metals are related to carotid atherosclerosis in elderly. <i>Science of the Total Environment</i> , 2012 , 416, 80-8 | 10.2 | 38 |
| 39 | Microarray profiling of diaphyseal bone of rats suffering from hypervitaminosis A. <i>Calcified Tissue International</i> , 2012 , 90, 219-29 | 3.9 | 9 |
| 38 | Elevated circulating levels of copper and nickel are found in elderly subjects with left ventricular hypertrophy. <i>Ecotoxicology and Environmental Safety</i> , 2012 , 86, 66-72 | 7 | 10 |
| 37 | Circulating levels of bisphenol A (BPA) and phthalates in an elderly population in Sweden, based on the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS). <i>Ecotoxicology and Environmental Safety</i> , 2012 , 75, 242-8 | 7 | 39 |
| 36 | Associations between circulating levels of bisphenol A and phthalate metabolites and coronary risk in the elderly. <i>Ecotoxicology and Environmental Safety</i> , 2012 , 80, 179-83 | 7 | 89 |
| 35 | Associations of persistent organic pollutants with abdominal obesity in the elderly: The Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study. <i>Environment International</i> , 2012 , 40, 170-178 | 12.9 | 99 |
| 34 | Circulating levels of persistent organic pollutants (POPs) among elderly men and women from Sweden: results from the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS). <i>Environment International</i> , 2012 , 44, 59-67 | 12.9 | 36 |
| 33 | Background exposure to persistent organic pollutants predicts stroke in the elderly. <i>Environment International</i> , 2012 , 47, 115-20 | 12.9 | 56 |
| 32 | Serum concentrations of phthalate metabolites are related to abdominal fat distribution two years later in elderly women. <i>Environmental Health</i> , 2012 , 11, 21 | 6 | 83 |
| 31 | Circulating levels of phthalate metabolites are associated with prevalent diabetes in the elderly. <i>Diabetes Care</i> , 2012 , 35, 1519-24 | 14.6 | 130 |
| 30 | A rapid method for screening of the Stockholm Convention POPs in small amounts of human plasma using SPE and HRGC/HRMS. <i>Chemosphere</i> , 2012 , 86, 747-53 | 8.4 | 55 |
| 29 | Gender differences for associations between circulating levels of metals and coronary risk in the elderly. <i>International Journal of Hygiene and Environmental Health</i> , 2012 , 215, 411-7 | 6.9 | 28 |
| 28 | Circulating levels of persistent organic pollutants (POPs) and carotid atherosclerosis in the elderly. <i>Environmental Health Perspectives</i> , 2012 , 120, 38-43 | 8.4 | 83 |
| 27 | An investigation of the co-variation in circulating levels of a large number of environmental contaminants. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2012 , 22, 476-82 | 6.7 | 19 |

| | | | |
|----|---|------|-----|
| 26 | High dietary intake of retinol leads to bone marrow hypoxia and diaphyseal endosteal mineralization in rats. <i>Bone</i> , 2011 , 48, 496-506 | 4.7 | 35 |
| 25 | Circulating levels of bisphenol A and phthalates are related to carotid atherosclerosis in the elderly. <i>Atherosclerosis</i> , 2011 , 218, 207-13 | 3.1 | 131 |
| 24 | Circulating levels of persistent organic pollutants associate in divergent ways to fat mass measured by DXA in humans. <i>Chemosphere</i> , 2011 , 85, 335-43 | 8.4 | 58 |
| 23 | Polychlorinated biphenyls and organochlorine pesticides in plasma predict development of type 2 diabetes in the elderly: the prospective investigation of the vasculature in Uppsala Seniors (PIVUS) study. <i>Diabetes Care</i> , 2011 , 34, 1778-84 | 14.6 | 189 |
| 22 | Transgenic mice with a constitutively active aryl hydrocarbon receptor display a gender-specific bone phenotype. <i>Toxicological Sciences</i> , 2010 , 114, 48-58 | 4.4 | 30 |
| 21 | Effects on bone tissue in ewes (<i>Ovis aries</i>) and their foetuses exposed to PCB 118 and PCB 153. <i>Toxicology Letters</i> , 2010 , 192, 126-33 | 4.4 | 20 |
| 20 | Pregnant ewes exposed to multiple endocrine disrupting pollutants through sewage sludge-fertilized pasture show an anti-estrogenic effect in their trabecular bone. <i>Science of the Total Environment</i> , 2010 , 408, 2340-6 | 10.2 | 22 |
| 19 | Quantitative characterization of changes in bone geometry, mineral density and biomechanical properties in two rat strains with different Ah-receptor structures after long-term exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. <i>Toxicology</i> , 2010 , 273, 1-11 | 4.4 | 24 |
| 18 | Exposure to pastures fertilised with sewage sludge disrupts bone tissue homeostasis in sheep. <i>Science of the Total Environment</i> , 2009 , 407, 2200-8 | 10.2 | 31 |
| 17 | Short-term exposure to dioxin impairs bone tissue in male rats. <i>Chemosphere</i> , 2009 , 75, 680-684 | 8.4 | 24 |
| 16 | Effects of 3,3,4,4,5-pentachlorobiphenyl (PCB126) on vertebral bone mineralization and on thyroxin and vitamin D levels in Sprague-Dawley rats. <i>Toxicology Letters</i> , 2009 , 187, 63-8 | 4.4 | 34 |
| 15 | Health of herring gulls (<i>Larus argentatus</i>) in relation to breeding location in the early 1990s. III. Effects on the bone tissue. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2008 , 71, 1448-56 | 3.2 | 17 |
| 14 | Bone mineral density changes in relation to environmental PCB exposure. <i>Environmental Health Perspectives</i> , 2008 , 116, 1162-6 | 8.4 | 52 |
| 13 | In utero and lactational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) affects bone tissue in rhesus monkeys. <i>Toxicology</i> , 2008 , 253, 147-52 | 4.4 | 35 |
| 12 | Effects of short-term exposure to the DDT metabolite p,pSDDE on bone tissue in male common frog (<i>Rana temporaria</i>). <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2007 , 70, 614-9 | 3.2 | 13 |
| 11 | Osteopontin: a rapid and sensitive response to dioxin exposure in the osteoblastic cell line UMR-106. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 341, 116-20 | 3.4 | 12 |
| 10 | Perinatal exposure to PCB 153, but not PCB 126, alters bone tissue composition in female goat offspring. <i>Toxicology</i> , 2006 , 228, 33-40 | 4.4 | 37 |
| 9 | Vitamin D fails to prevent serum starvation- or staurosporine-induced apoptosis in human and rat osteosarcoma-derived cell lines. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 330, 891-7 | 3.4 | 9 |

| | | | |
|---|--|-----|----|
| 8 | Subchronic Toxicity of Baltic Herring Oil and its Fractions in the Rat (III) Bone Tissue Composition and Dimension, and Ratio of n-6/n-3 Fatty Acids in Serum Phospholipids. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2005 , 96, 453-464 | 3.1 | 8 |
| 7 | Altered retinoid metabolism in female Long-Evans and Han/Wistar rats following long-term 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-treatment. <i>Toxicological Sciences</i> , 2005 , 86, 264-72 | 4.4 | 24 |
| 6 | Abnormal bone composition in female juvenile American alligators from a pesticide-polluted lake (Lake Apopka, Florida). <i>Environmental Health Perspectives</i> , 2004 , 112, 359-62 | 8.4 | 72 |
| 5 | Estrogen supplementation modulates effects of the endocrine disrupting pollutant PCB126 in rat bone and uterus: diverging effects in ovariectomized and intact animals. <i>Toxicology</i> , 2004 , 199, 129-36 | 4.4 | 28 |
| 4 | The dioxin-like pollutant PCB 126 (3,3,4,4,5-pentachlorobiphenyl) affects risk factors for cardiovascular disease in female rats. <i>Toxicology Letters</i> , 2004 , 150, 293-9 | 4.4 | 97 |
| 3 | Bone mineral density in male Baltic grey seal (<i>Halichoerus grypus</i>). <i>Ambio</i> , 2003 , 32, 385-8 | 6.5 | 48 |
| 2 | Subchronic toxicity of Baltic herring oil and its fractions in the rat II: Clinical observations and toxicological parameters. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2002 , 91, 232-44 | | 5 |
| 1 | Circulating Levels of Persistent Organic Pollutants in Relation to Visceral and Subcutaneous Adipose Tissue by Abdominal MRI. <i>Obesity</i> , | 8 | 3 |