

# Sylvain Dubucquoi

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/6684109/sylvain-dubucquoi-publications-by-citations.pdf>

**Version:** 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62

papers

1,506

citations

23

h-index

37

g-index

66

ext. papers

1,767

ext. citations

6.1

avg, IF

3.72

L-index

| #  | Paper  | IF   | Citations |
|----|--|------|-----------|
| 62 | Distinct immune response in two MERS-CoV-infected patients: can we go from bench to bedside?. <i>PLoS ONE</i> , <b>2014</b> , 9, e88716  | 3.7  | 180       |
| 61 | Hierarchical cluster and survival analyses of antisynthetase syndrome: phenotype and outcome are correlated with anti-tRNA synthetase antibody specificity. <i>Autoimmunity Reviews</i> , <b>2012</b> , 12, 210-7  | 13.6 | 177       |
| 60 | The Spectrum of FIP1L1-PDGFR $\alpha$ -Associated Chronic Eosinophilic Leukemia: New Insights Based on a Survey of 44 Cases. <i>Medicine (United States)</i> , <b>2013</b> , 92, e1-e9   | 1.8  | 62        |
| 59 | Clinical manifestations of anti-synthetase syndrome positive for anti-alanyl-tRNA synthetase (anti-PL12) antibodies: a retrospective study of 17 cases. <i>Rheumatology</i> , <b>2010</b> , 49, 972-6  | 3.9  | 60        |
| 58 | New antigenic candidates in multiple sclerosis: identification by serological proteome analysis. <i>Proteomics</i> , <b>2004</b> , 4, 2184-94  | 4.8  | 60        |
| 57 | Prevalence of anti-RNA polymerase III antibodies in systemic sclerosis: New data from a French cohort and a systematic review and meta-analysis. <i>Arthritis and Rheumatology</i> , <b>2014</b> , 66, 407-17  | 9.5  | 57        |
| 56 | New insights into cell responses involved in experimental autoimmune encephalomyelitis and multiple sclerosis. <i>Immunology Letters</i> , <b>2005</b> , 96, 11-26   | 4.1  | 56        |
| 55 | Total soluble HLA class I and soluble HLA-G in multiple myeloma and monoclonal gammopathy of undetermined significance. <i>Clinical Cancer Research</i> , <b>2005</b> , 11, 7297-303   | 12.9 | 52        |
| 54 | Endothelial dysfunction in systemic lupus patients with low disease activity: evaluation by quantification and characterization of circulating endothelial microparticles, role of anti-endothelial cell antibodies. <i>Rheumatology</i> , <b>2010</b> , 49, 1049-55 | 3.9  | 44        |
| 53 | Characterization of discriminant human brain antigenic targets in neuropsychiatric systemic lupus erythematosus using an immunoproteomic approach. <i>Arthritis and Rheumatism</i> , <b>2007</b> , 56, 3420-32   |      | 44        |
| 52 | Antineutrophil Cytoplasmic Antibody-Positive Small-Vessel Vasculitis Associated with Antithyroid Drug Therapy: How Significant Is the Clinical Problem?. <i>Thyroid</i> , <b>2015</b> , 25, 1273-81  | 6.2  | 41        |
| 51 | Distortion of the self-reactive IgG antibody repertoire in multiple sclerosis as a new diagnostic tool. <i>Journal of Immunology</i> , <b>2004</b> , 172, 669-78   | 5.3  | 41        |
| 50 | Intravenous immunoglobulins in systemic sclerosis: Data from a French nationwide cohort of 46 patients and review of the literature. <i>Autoimmunity Reviews</i> , <b>2017</b> , 16, 377-384   | 13.6 | 35        |
| 49 | Seronegative polyarthritis revealing antisynthetase syndrome: a multicentre study of 40 patients. <i>Rheumatology</i> , <b>2015</b> , 54, 927-32   | 3.9  | 35        |
| 48 | Interleukin-5 messenger RNA and immunoreactive protein expression by activated eosinophils in lesional atopic dermatitis skin. <i>Journal of Investigative Dermatology</i> , <b>1994</b> , 103, 589-92   | 4.3  | 34        |
| 47 | Altered B lymphocyte homeostasis and functions in systemic sclerosis. <i>Autoimmunity Reviews</i> , <b>2018</b> , 17, 244-255  | 13.6 | 31        |
| 46 | Serum free light chains of immunoglobulins as biomarkers for systemic sclerosis characteristics, activity and severity. <i>Autoimmunity Reviews</i> , <b>2014</b> , 13, 974-80   | 13.6 | 29        |

|    |  |      |    |
|----|--|------|----|
| 45 | Effect of in vitro and in vivo anakinra on cytokines production in Schnitzler syndrome. <i>PLoS ONE</i> , <b>2013</b> , 8, e59327  | 3.7  | 28 |
| 44 | Heterogeneous clinical spectrum of anti-SRP myositis and importance of the methods of detection of anti-SRP autoantibodies: a multicentric study. <i>Immunologic Research</i> , <b>2016</b> , 64, 677-86   | 4.3  | 26 |
| 43 | Comparison of the Farr radioimmunoassay, 3 commercial enzyme immunoassays and Crithidia luciliae immunofluorescence test for diagnosis and activity assessment of systemic lupus erythematosus. <i>Clinica Chimica Acta</i> , <b>2010</b> , 411, 959-64  | 6.2  | 26 |
| 42 | Evidence for eosinophil activation in eosinophilic cystitis. <i>European Urology</i> , <b>1994</b> , 25, 254-8   | 10.2 | 25 |
| 41 | CD3-CD4+ lymphoid variant of hypereosinophilic syndrome: nodal and extranodal histopathological and immunophenotypic features of a peripheral indolent clonal T-cell lymphoproliferative disorder. <i>Haematologica</i> , <b>2015</b> , 100, 1086-95   | 6.6  | 24 |
| 40 | Humoral immunity links <i>Candida albicans</i> infection and celiac disease. <i>PLoS ONE</i> , <b>2015</b> , 10, e0121776  | 3.7  | 23 |
| 39 | B-cell subsets up-regulate $\alpha$ 4 integrin and accumulate in the cerebrospinal fluid in clinically isolated syndrome suggestive of multiple sclerosis onset. <i>Neuroscience Letters</i> , <b>2011</b> , 487, 273-7  | 3.3  | 23 |
| 38 | A functional haplotype of PADI4 gene in rheumatoid arthritis: positive correlation in a French population. <i>Journal of Rheumatology</i> , <b>2009</b> , 36, 881-6  | 4.1  | 23 |
| 37 | Diversified serum IgG response involving non-myelin CNS proteins during experimental autoimmune encephalomyelitis. <i>Journal of Neuroimmunology</i> , <b>2006</b> , 179, 53-64  | 3.5  | 23 |
| 36 | Proinflammatory B-cell profile in the early phases of MS predicts an active disease. <i>Neurology: Neuroimmunology and NeuroInflammation</i> , <b>2018</b> , 5, e431   | 9.1  | 18 |
| 35 | Primed status of transitional B cells associated with their presence in the cerebrospinal fluid in early phases of multiple sclerosis. <i>Clinical Immunology</i> , <b>2011</b> , 139, 12-20   | 9    | 17 |
| 34 | Low endocan levels are predictive of Acute Respiratory Distress Syndrome in severe sepsis and septic shock. <i>Journal of Critical Care</i> , <b>2018</b> , 47, 121-126  | 4    | 16 |
| 33 | Autoantibodies against alpha-fodrin in Sjögren's syndrome with neurological manifestations. <i>Journal of Rheumatology</i> , <b>2004</b> , 31, 500-3   | 4.1  | 16 |
| 32 | C-Reactive protein as a diagnostic tool in differential diagnosis of hypereosinophilic syndrome and antineutrophil cytoplasmic antibody-negative eosinophilic granulomatosis with polyangiitis. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , <b>2019</b> , 7, 1347-1351.e3 | 5.4  | 15 |
| 31 | Value of the Overall Pneumococcal Polysaccharide Response in the Diagnosis of Primary Humoral Immunodeficiencies. <i>Frontiers in Immunology</i> , <b>2017</b> , 8, 1862   | 8.4  | 14 |
| 30 | Interleukin-10 promoter polymorphism in multiple sclerosis: association with disease progression. <i>European Cytokine Network</i> , <b>2002</b> , 13, 200-6   | 3.3  | 13 |
| 29 | Endocan is a stable circulating molecule in ICU patients. <i>Clinical Biochemistry</i> , <b>2017</b> , 50, 870-877   | 3.5  | 12 |
| 28 | Prevalence and Clinical Associations of Antiphospholipid Antibodies in Systemic Sclerosis: New Data From a French Cross-Sectional Study, Systematic Review, and Meta-Analysis. <i>Frontiers in Immunology</i> , <b>2018</b> , 9, 2457  | 8.4  | 11 |

|    |   |      |    |
|----|---|------|----|
| 27 | Specific Polysaccharide Antibody Deficiency Revealed by Severe Bacterial Infections in Adulthood: A Report on 11 Cases. <i>Clinical Infectious Diseases</i> , <b>2017</b> , 65, 328-331   | 11.6 | 10 |
| 26 | IgM-specific measles-virus antibody in families with a high frequency of Crohn's disease. <i>Lancet, The</i> , <b>1995</b> , 346, 967   | 4.0  | 9  |
| 25 | An Optimized Fluorescence-Based Bidimensional Immunoproteomic Approach for Accurate Screening of Autoantibodies. <i>PLoS ONE</i> , <b>2015</b> , 10, e0132142   | 3.7  | 8  |
| 24 | Diluted Russell viper-venom time improves identification of antiphospholipid syndrome in a lupus anticoagulant-positive patient population. <i>Thrombosis and Haemostasis</i> , <b>2009</b> , 101, 577-581  | 7    | 8  |
| 23 | CD28+ intraepithelial lymphocytes with long telomeres are recruited within the inflamed ileal mucosa in Crohn disease. <i>Human Immunology</i> , <b>2001</b> , 62, 694-700  | 2.3  | 8  |
| 22 | Impact of acute renal failure on plasmatic levels of cleaved endocan. <i>Critical Care</i> , <b>2019</b> , 23, 55   | 10.8 | 7  |
| 21 | Vaginal Mucosal Homeostatic Response May Determine Pregnancy Outcome in Women With Bacterial Vaginosis: A Pilot Study. <i>Medicine (United States)</i> , <b>2016</b> , 95, e2668  | 1.8  | 7  |
| 20 | Antiphospholipid Syndrome With Isolated Isotype M Anticardiolipin and/or Anti-B2GPI Antibody Is Associated With Stroke. <i>Stroke</i> , <b>2018</b> , 49, 2770-2772   | 6.7  | 7  |
| 19 | Significance of persistent antiphospholipid antibodies in the elderly. <i>Journal of Rheumatology</i> , <b>2006</b> , 33, 1559-62   | 4.1  | 7  |
| 18 | Immunoglobulin G (IgG) and IgG subclass reference intervals in children, using Optilite <sup>®</sup> reagents. <i>Clinical Chemistry and Laboratory Medicine</i> , <b>2018</b> , 56, 1319-1327  | 5.9  | 5  |
| 17 | B Cell Homeostasis and Functional Properties Are Altered in an Hypochlorous Acid-Induced Murine Model of Systemic Sclerosis. <i>Frontiers in Immunology</i> , <b>2017</b> , 8, 53   | 8.4  | 5  |
| 16 | Are immunoglobulin A anti-gliadin antibodies helpful in diagnosing coeliac disease in children younger than 2 years?. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , <b>2012</b> , 54, 110-2  | 2.8  | 5  |
| 15 | Serological proteome analysis reveals new specific biases in the IgM and IgG autoantibody repertoires in autoimmune polyendocrine syndrome type 1. <i>Autoimmunity</i> , <b>2015</b> , 48, 532-41   | 3    | 4  |
| 14 | Decrease of the plasmatic endocan cleavage ratio is associated with the hyperinflammatory phenotype of acute respiratory distress syndrome. <i>Critical Care</i> , <b>2019</b> , 23, 252  | 10.8 | 4  |
| 13 | Soluble form of receptor for advanced glycation end-products (sRAGE): do sRAGE ligands or anti-sRAGE auto-antibodies interfere with sRAGE quantification?. <i>Annals of Clinical Biochemistry</i> , <b>2014</b> , 51, 248-57  | 2.2  | 4  |
| 12 | Assessment of T-cell polarization on the basis of surface marker expression: Diagnosis and potential therapeutic implications in lymphocytic variant hypereosinophilic syndrome. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , <b>2020</b> , 8, 1110-1114.e2 | 5.4  | 4  |
| 11 | Extended myositis-specific and -associated antibodies profile in systemic sclerosis: A cross-sectional study. <i>Joint Bone Spine</i> , <b>2021</b> , 88, 105048  | 2.9  | 4  |
| 10 | Clinical significance of anti-Ro52 (TRIM21) antibodies in adult patients with connective tissue diseases. <i>European Journal of Internal Medicine</i> , <b>2021</b> , 91, 45-52  | 3.9  | 4  |

|   |   |      |   |
|---|---|------|---|
| 9 | Whole-Body Distribution and Clinical Association of Telangiectases in Systemic Sclerosis. <i>JAMA Dermatology</i> , <b>2018</b> , 154, 796-805  | 5.1  | 3 |
| 8 | The complex kinetics of blood endocan during the time course of sepsis and acute respiratory distress syndrome. <i>Critical Care</i> , <b>2019</b> , 23, 86                               | 10.8 | 1 |
| 7 | Changes Related to Age in Natural and Acquired Systemic Self-IgG Responses in Malaria. <i>Interdisciplinary Perspectives on Infectious Diseases</i> , <b>2011</b> , 2011, 462767          | 1.7  | 0 |
| 6 | Classical pathway activity C3c, C4 and C1-inhibitor protein reference intervals determination in EDTA plasma. <i>Biochemia Medica</i> , <b>2019</b> , 29, 030707                          | 2.5  | 0 |
| 5 | Anti-Ephrin Type-B Receptor 2 (EphB2) and Anti-Three Prime Histone mRNA EXonuclease 1 (THEX1) Autoantibodies in Scleroderma and Lupus. <i>PLoS ONE</i> , <b>2016</b> , 11, e0160283       | 3.7  | 0 |
| 4 | Diagnostic performance of a new vimentin-derived ACPA (CCP high sensitive) in patients with rheumatoid arthritis. <i>Immunologic Research</i> , <b>2016</b> , 64, 455-60                  | 4.3  |   |
| 3 | Reply to Gilchrist et al. and to Musher. <i>Clinical Infectious Diseases</i> , <b>2018</b> , 66, 637-638  | 11.6 |   |
| 2 | Passages in culture and stimulation conditions influence protein expression of primary fibroblasts. <i>Proteomics</i> , <b>2021</b> , e2100116  | 4.8  |   |
| 1 | Étude transversale des profils anticorps spécifiques et associés aux myosites dans la sclérodermie systémique. <i>Revue Du Rhumatisme (Edition Française)</i> , <b>2021</b> , 88, 284-290 | 0.1  |   |