Frederik Vandenberghe

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6684985/publications.pdf

Version: 2024-02-01

23 papers 314 citations

8 h-index 17 g-index

23 all docs 23 docs citations

times ranked

23

544 citing authors

#	Article	IF	CITATIONS
1	Importance of Early Weight Changes to Predict Long-Term Weight Gain During Psychotropic Drug Treatment. Journal of Clinical Psychiatry, 2015, 76, e1417-e1423.	1.1	67
2	Genetics-Based Population Pharmacokinetics and Pharmacodynamics of Risperidone in a Psychiatric Cohort. Clinical Pharmacokinetics, 2015, 54, 1259-1272.	1.6	44
3	Influence of <i>CRTC1</i> Polymorphisms on Body Mass Index and Fat Mass in Psychiatric Patients and the General Adult Population. JAMA Psychiatry, 2013, 70, 1011.	6.0	42
4	Association of CRTC1 polymorphisms with obesity markers in subjects from the general population with lifetime depression. Journal of Affective Disorders, 2016, 198, 43-49.	2.0	18
5	Tools for optimising pharmacotherapy in psychiatry (therapeutic drug monitoring, molecular brain) Tj ETQq1 1 0. Psychiatry, 2021, 22, 561-628.	.784314 1.3	l rgBT /Overlo <mark>ck</mark> 17
6	Influence of MCHR2 and MCHR2-AS1 Genetic Polymorphisms on Body Mass Index in Psychiatric Patients and In Population-Based Subjects with Present or Past Atypical Depression. PLoS ONE, 2015, 10, e0139155.	1.1	16
7	Early changes of blood lipid levels during psychotropic drug treatment as predictors of long-term lipid changes and of new onset dyslipidemia. Journal of Clinical Lipidology, 2018, 12, 219-229.	0.6	15
8	Second-Generation Antipsychotics in Adolescent Psychiatric Patients: Metabolic Effects and Impact of an Early Weight Change to Predict Longer Term Weight Gain. Journal of Child and Adolescent Psychopharmacology, 2018, 28, 258-265.	0.7	13
9	Psychological trauma occurring during adolescence is associated with an increased risk of greater waist circumference in Early Psychosis patients treated with psychotropic medication. PLoS ONE, 2020, 15, e0242569.	1.1	10
10	Prediction of early weight gain during psychotropic treatment using a combinatorial model with clinical and genetic markers. Pharmacogenetics and Genomics, 2016, 26, 547-557.	0.7	9
11	Olanzapineâ€associated doseâ€dependent alterations for weight and metabolic parameters in a prospective cohort. Basic and Clinical Pharmacology and Toxicology, 2022, 130, 531-541.	1.2	9
12	Influence of polygenic risk scores on lipid levels and dyslipidemia in a psychiatric population receiving weight gain-inducing psychotropic drugs. Pharmacogenetics and Genomics, 2017, 27, 464-472.	0.7	8
13	Association of variants in SH2B1 and RABEP1 with worsening of low-density lipoprotein and glucose parameters in patients treated with psychotropic drugs. Gene, 2017, 628, 8-15.	1.0	7
14	Clinical Pharmacy in Psychiatry: Towards Promoting Clinical Expertise in Psychopharmacology. Pharmacy (Basel, Switzerland), 2021, 9, 146.	0.6	7
15	Insomnia disorders are associated with increased cardiometabolic disturbances and death risks from cardiovascular diseases in psychiatric patients treated with weight-gain-inducing psychotropic drugs: results from a Swiss cohort. BMC Psychiatry, 2022, 22, 342.	1.1	7
16	Bioavailability of Vortioxetine After a Roux-en-Y Gastric Bypass. Obesity Surgery, 2021, 31, 1353-1356.	1.1	6
17	Genetic immune and inflammatory markers associated with diabetes in solid organ transplant recipients. American Journal of Transplantation, 2019, 19, 238-246.	2.6	5
18	Association of genetic risk scores with body mass index in Swiss psychiatric cohorts. Pharmacogenetics and Genomics, 2016, 26, 208-217.	0.7	3

#	Article	IF	CITATIONS
19	Association Between Plasma Caffeine and Other Methylxanthines and Metabolic Parameters in a Psychiatric Population Treated With Psychotropic Drugs Inducing Metabolic Disturbances. Frontiers in Psychiatry, 2018, 9, 573.	1.3	3
20	Impact of drugs on hypoglycaemia in hospitalised patients. European Journal of Hospital Pharmacy, 2019, 26, 199-204.	0.5	3
21	Lipid Disturbances in Adolescents Treated With Second-Generation Antipsychotics. Journal of Clinical Psychiatry, 2019, 80, .	1.1	3
22	Associations Between High Plasma Methylxanthine Levels, Sleep Disorders and Polygenic Risk Scores of Caffeine Consumption or Sleep Duration in a Swiss Psychiatric Cohort. Frontiers in Psychiatry, 2021, 12, 756403.	1.3	2
23	F227. PSYCHOLOGICAL TRAUMA OCCURRING DURING ADOLESCENCE IS ASSOCIATED WITH AN INCREASED RISK OF GREATER WAIST CIRCUMFERENCE IN EARLY PSYCHOSIS PATIENTS INDEPENDENTLY OF MEDICATION. Schizophrenia Bulletin, 2018, 44, S310-S310.	2.3	0