

Madeleine Duvic

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6688498/madeleine-duvic-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

352
papers

19,940
citations

73
h-index

133
g-index

363
ext. papers

22,508
ext. citations

4
avg, IF

6.45
L-index

| # | Paper | IF | Citations |
|-----|---|-----|-----------|
| 352 | New Practical Aspects of Sweet Syndrome.. <i>American Journal of Clinical Dermatology</i> , 2022 , 1 | 7.1 | 5 |
| 351 | Vulvar Primary Cutaneous CD8+ Aggressive Epidermotropic Cytotoxic T-Cell Lymphoma. <i>International Journal of Gynecological Pathology</i> , 2021 , 40, 229-233 | 3.2 | 1 |
| 350 | Teledermatology During COVID-19: An Updated Review. <i>American Journal of Clinical Dermatology</i> , 2021 , 22, 467-475 | 7.1 | 7 |
| 349 | Response to brentuximab vedotin versus physician choice by CD30 expression and large cell transformation status in patients with mycosis fungoides: An ALCANZA sub-analysis. <i>European Journal of Cancer</i> , 2021 , 148, 411-421 | 7.5 | 6 |
| 348 | Post-transplantation donor-derived Sezary syndrome in a patient with A91V PRF1 variant hemophagocytic lymphohistiocytosis. <i>American Journal of Hematology</i> , 2021 , 96, E350-E353 | 7.1 | 1 |
| 347 | Determination of immunophenotypic aberrancies provides better assessment of peripheral blood involvement by mycosis fungoides/Sezary syndrome than quantification of CD26- or CD7- CD4+ T-cells. <i>Cytometry Part B - Clinical Cytometry</i> , 2021 , 100, 183-191 | 3.4 | 8 |
| 346 | Quality of Life Effect of the Anti-CCR4 Monoclonal Antibody Mogamulizumab Versus Vorinostat in Patients With Cutaneous T-cell Lymphoma. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2021 , 21, 97-105 ² | | 7 |
| 345 | Is immunohistochemical expression of GATA3 helpful in the differential diagnosis of transformed mycosis fungoides and primary cutaneous CD30-positive T cell lymphoproliferative disorders?. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2021 , 479, 377-383 | 5.1 | 1 |
| 344 | Randomized phase 3 ALCANZA study of brentuximab vedotin vs physician choice in cutaneous T-cell lymphoma: final data. <i>Blood Advances</i> , 2021 , 5, 5098-5106 | 7.8 | 5 |
| 343 | Monitoring malignant T-cell clones by direct TCR expression assay in patients with leukemic cutaneous T-cell lymphoma during extracorporeal photopheresis. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2021 , | 2.4 | 0 |
| 342 | Granuloma Annulare: An Updated Review of Epidemiology, Pathogenesis, and Treatment Options. <i>American Journal of Clinical Dermatology</i> , 2021 , 1 | 7.1 | 4 |
| 341 | Unmasking a T cell lymphoma: Folliculotropic mycosis fungoides with a gamma-delta phenotype. <i>JAAD Case Reports</i> , 2020 , 6, 1316-1319 | 1.4 | 2 |
| 340 | Patient-reported quality of life in patients with relapsed/refractory cutaneous T-cell lymphoma: Results from the randomised phase III ALCANZA study. <i>European Journal of Cancer</i> , 2020 , 133, 120-130 | 7.5 | 11 |
| 339 | CD209 monocyte-derived myeloid dendritic cells were increased in patients with leukemic cutaneous T-cell lymphoma undergoing extracorporeal photopheresis via the CELLEX system. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2020 , 36, 290-298 | 2.4 | 2 |
| 338 | Non-Classic Signs of Sezary Syndrome: A Review. <i>American Journal of Clinical Dermatology</i> , 2020 , 21, 383-391 | 7.1 | 4 |
| 337 | Cutaneous T-Cell Lymphoma and Cutaneous B-Cell Lymphoma 2020 , 1948-1964.e5 | | |
| 336 | Methicillin-resistant Staphylococcus aureus (MRSA) is an important pathogen in erythrodermic cutaneous T-cell lymphoma (CTCL) patients. <i>Archives of Dermatological Research</i> , 2020 , 312, 283-288 | 3.3 | 7 |

| | | | |
|-----|---|------|-----|
| 335 | Lymphomatoid Papulosis With a Unique T Follicular Helper-Like Phenotype. <i>American Journal of Dermatopathology</i> , 2020 , 42, 776-779 | 0.9 | |
| 334 | Renal Cell Carcinoma Associated with Mycosis Fungoides: A Paraneoplastic Syndrome. <i>Case Reports in Nephrology</i> , 2020 , 2020, 8897183 | 0.8 | 0 |
| 333 | Second primary malignancies in blastic plasmacytoid dendritic cell neoplasm: A national database study. <i>Journal of the American Academy of Dermatology</i> , 2020 , 83, 1786-1789 | 4.5 | 1 |
| 332 | United States Cutaneous Lymphoma Consortium recommendations for treatment of cutaneous lymphomas during the COVID-19 pandemic. <i>Journal of the American Academy of Dermatology</i> , 2020 , 83, 703-704 | 4.5 | 15 |
| 331 | Development of Sjögren syndrome following the administration of dupilumab. <i>Dermatology Online Journal</i> , 2020 , 26, | 1 | 2 |
| 330 | Anaphylaxis following administration of extracorporeal photopheresis for cutaneous T cell lymphoma. <i>Dermatology Online Journal</i> , 2020 , 26, | 1 | |
| 329 | Multi-institutional Investigation: Circulating CD4:CD8 ratio is a prognosticator of response to total skin electron beam radiation in mycosis fungoides. <i>Radiotherapy and Oncology</i> , 2019 , 131, 88-92 | 5.3 | 3 |
| 328 | Long-term control of mycosis fungoides of the hands with topical bexarotene: an update 15 years later. <i>International Journal of Dermatology</i> , 2019 , 58, e221-e222 | 1.7 | 2 |
| 327 | Tagraxofusp in Blastic Plasmacytoid Dendritic-Cell Neoplasm. <i>New England Journal of Medicine</i> , 2019 , 380, 1628-1637 | 59.2 | 173 |
| 326 | FINAL DATA FROM THE PHASE 3 ALCANZA STUDY: BRENTUXIMAB VEDOTIN (BV) VS PHYSICIAN'S CHOICE (PC) IN PATIENTS (PTS) WITH CD30-POSITIVE (CD30+) CUTANEOUS T-CELL LYMPHOMA (CTCL). <i>Hematological Oncology</i> , 2019 , 37, 286-288 | 1.3 | 1 |
| 325 | Lichenoid dermatitis from immune checkpoint inhibitor therapy: An immune-related adverse event with mycosis-fungoides-like morphologic and molecular features. <i>Journal of Cutaneous Pathology</i> , 2019 , 46, 872-877 | 1.7 | 3 |
| 324 | Blood transcriptional profiling reveals IL-1 and integrin signaling pathways associated with clinical response to extracorporeal photopheresis in patients with leukemic cutaneous T-cell lymphoma. <i>Oncotarget</i> , 2019 , 10, 3183-3197 | 3.3 | 3 |
| 323 | RESPONSE TO BRENTUXIMAB VEDOTIN BY CD30 EXPRESSION: RESULTS FROM FIVE TRIALS IN PTCL, CTCL, AND B-CELL LYMPHOMAS. <i>Hematological Oncology</i> , 2019 , 37, 470-471 | 1.3 | 2 |
| 322 | Antibody-Based Therapies for Cutaneous T-Cell Lymphoma. <i>American Journal of Clinical Dermatology</i> , 2019 , 20, 115-122 | 7.1 | 15 |
| 321 | Proteomic analysis of stratum corneum in Cutaneous T-Cell Lymphomas and psoriasis. <i>Experimental Dermatology</i> , 2019 , 28, 317-321 | 4 | 2 |
| 320 | Response to pembrolizumab and lenalidomide in advanced refractory mycosis fungoides. <i>Leukemia and Lymphoma</i> , 2019 , 60, 1079-1082 | 1.9 | 2 |
| 319 | Primary cutaneous CD4+ small- to medium-sized pleomorphic T-cell lymphoproliferative disorder in a pediatric patient successfully treated with low-dose radiation. <i>Pediatric Dermatology</i> , 2019 , 36, e23-e26 | 1.9 | 1 |
| 318 | Nail irregularities associated with Sjögren syndrome. <i>Cutis</i> , 2019 , 103, E11-E16 | 0.4 | 2 |

| | | | |
|-----|---|------|-----|
| 317 | Waistband Mycosis Fungoides: A New Clinical Variant of Early-Stage Disease. <i>Skinmed</i> , 2019 , 17, 329-332. | 0.2 | 1 |
| 316 | Necrotizing Granulomatous Dermatitis and Panniculitis Masquerading as T Cell Lymphoma. <i>Skinmed</i> , 2019 , 17, 406-408 | 0.2 | 1 |
| 315 | Image Gallery: Symmetrical whirled eschars on the face in mycosis fungoides. <i>British Journal of Dermatology</i> , 2018 , 178, e224 | 4 | 1 |
| 314 | Childhood alopecia areata-Data from the National Alopecia Areata Registry. <i>Pediatric Dermatology</i> , 2018 , 35, 164-169 | 1.9 | 12 |
| 313 | How to Discern Folliculotropic Mycosis Fungoides From Follicular Mucinosis Using a Pediatric Case. <i>Journal of Cutaneous Medicine and Surgery</i> , 2018 , 22, 336-340 | 1.6 | 11 |
| 312 | Pruritic arthropod bite-like papules in T-cell large granular lymphocytic leukaemia and chronic myelomonocytic leukaemia. <i>Clinical and Experimental Dermatology</i> , 2018 , 43, 449-453 | 1.8 | 2 |
| 311 | The Utility and Validity of the Alopecia Areata Symptom Impact Scale in Measuring Disease-Related Symptoms and their Effect on Functioning. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 2018 , 19, S41-S46 | 1.1 | 10 |
| 310 | Mycosis fungoides occurring at the site of previous herpes zoster eruption. <i>Australasian Journal of Dermatology</i> , 2018 , 59, 217-219 | 1.3 | 2 |
| 309 | Generalized morphea/eosinophilic fasciitis overlap after epoxy exposure. <i>JAAD Case Reports</i> , 2018 , 4, 175-178 | 1.4 | 2 |
| 308 | The Use of Central Pathology Review With Digital Slide Scanning in Advanced-stage Mycosis Fungoides and Sézary Syndrome: A Multi-institutional and International Pathology Study. <i>American Journal of Surgical Pathology</i> , 2018 , 42, 726-734 | 6.7 | 14 |
| 307 | Important considerations for legislation banning commercial tanning beds among minors. <i>Clinics in Dermatology</i> , 2018 , 36, 104-105 | 3 | |
| 306 | Differential expression of CCR4 in primary cutaneous gamma/delta (T) cell lymphomas and mycosis fungoides: Significance for diagnosis and therapy. <i>Journal of Dermatological Science</i> , 2018 , 89, 88-91 | 4.3 | 9 |
| 305 | Responses to romidepsin in patients with cutaneous T-cell lymphoma and prior treatment with systemic chemotherapy. <i>Leukemia and Lymphoma</i> , 2018 , 59, 880-887 | 1.9 | 16 |
| 304 | Alternate dosing regimens of brentuximab vedotin for CD30+ cutaneous T-cell lymphoma. <i>British Journal of Dermatology</i> , 2018 , 178, 302-303 | 4 | 4 |
| 303 | Gene expression profiling and immune cell-type deconvolution highlight robust disease progression and survival markers in multiple cohorts of CTCL patients. <i>Onc Immunology</i> , 2018 , 7, e1467856 | 7.3 | 16 |
| 302 | Mogamulizumab versus vorinostat in previously treated cutaneous T-cell lymphoma (MAVORIC): an international, open-label, randomised, controlled phase 3 trial. <i>Lancet Oncology</i> , 2018 , 19, 1192-1204 | 21.7 | 239 |
| 301 | Cobomarsen, an oligonucleotide inhibitor of miR-155, co-ordinately regulates multiple survival pathways to reduce cellular proliferation and survival in cutaneous T-cell lymphoma. <i>British Journal of Haematology</i> , 2018 , 183, 428-444 | 4.5 | 129 |
| 300 | High-throughput T cell receptor sequencing identifies clonally expanded CD8+ T cell populations in alopecia areata. <i>JCI Insight</i> , 2018 , 3, | 9.9 | 25 |

| | | | |
|-----|--|------|-----|
| 299 | Juvenile mycosis fungoides with large-cell transformation: Successful treatment with psoralen with ultraviolet A light, interferon-alfa, and localized radiation. <i>Pediatric Dermatology</i> , 2018 , 35, e13-e16 | 1.9 | 2 |
| 298 | The "Duvic regimen" for erythrodermic flares secondary to Staphylococcus aureus in mycosis fungoides and Sjögren syndrome. <i>International Journal of Dermatology</i> , 2018 , 57, 123-124 | 1.7 | 8 |
| 297 | Radiotherapy in Patients with Mycosis Fungoides and Central Nervous System Involvement. <i>Case Reports in Oncology</i> , 2018 , 11, 721-728 | 1 | |
| 296 | Inflammatory cytokines and peripheral mediators in the pathophysiology of pruritus in cutaneous T-cell lymphoma. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2018 , 32, 1652-1656 | 4.6 | 12 |
| 295 | Complete resolution of mycosis fungoides tumors with imiquimod 5% cream: a case series. <i>Journal of Dermatological Treatment</i> , 2017 , 28, 567-569 | 2.8 | 13 |
| 294 | Results from a Phase I/II Open-Label, Dose-Finding Study of Pralatrexate and Oral Bexarotene in Patients with Relapsed/Refractory Cutaneous T-cell Lymphoma. <i>Clinical Cancer Research</i> , 2017 , 23, 3552-3556 | 12.9 | 19 |
| 293 | Immunophenotypic Shifts in Primary Cutaneous T-Cell Lymphoma Suggest Antigenic Modulation: A Study of Sequential Biopsy Specimens. <i>American Journal of Surgical Pathology</i> , 2017 , 41, 431-445 | 6.7 | 10 |
| 292 | Forodesine in the treatment of cutaneous T-cell lymphoma. <i>Expert Opinion on Investigational Drugs</i> , 2017 , 26, 771-775 | 5.9 | 5 |
| 291 | Gene expression analysis in Cutaneous T-Cell Lymphomas (CTCL) highlights disease heterogeneity and potential diagnostic and prognostic indicators. <i>Oncolmmunology</i> , 2017 , 6, e1306618 | 7.2 | 52 |
| 290 | Clinical characteristics, risk factors and long-term outcome of 114 patients with folliculotropic mycosis fungoides. <i>Archives of Dermatological Research</i> , 2017 , 309, 453-459 | 3.3 | 22 |
| 289 | Primary Cutaneous T-Cell Lymphomas Showing Gamma-Delta (γδ) Phenotype and Predominantly Epidermotropic Pattern are Clinicopathologically Distinct From Classic Primary Cutaneous T-Cell Lymphomas. <i>American Journal of Surgical Pathology</i> , 2017 , 41, 204-215 | 6.7 | 41 |
| 288 | Effectiveness of low-dose radiation for primary cutaneous anaplastic large cell lymphoma. <i>Advances in Radiation Oncology</i> , 2017 , 2, 363-369 | 3.3 | 5 |
| 287 | Brentuximab vedotin or physician's choice in CD30-positive cutaneous T-cell lymphoma (ALCANZA): an international, open-label, randomised, phase 3, multicentre trial. <i>Lancet, The</i> , 2017 , 390, 555-566 | 4.0 | 303 |
| 286 | Extracorporeal photopheresis for the treatment of early-stage mycosis fungoides. <i>Dermatologic Therapy</i> , 2017 , 30, e12485 | 2.2 | 6 |
| 285 | Brentuximab Vedotin for Patients With Refractory Lymphomatoid Papulosis: An Analysis of Phase 2 Results. <i>JAMA Dermatology</i> , 2017 , 153, 1302-1306 | 5.1 | 20 |
| 284 | Generalised Eruptive Keratoacanthomas of Grzybowski. <i>Journal of Cutaneous Medicine and Surgery</i> , 2017 , 21, 439 | 1.6 | |
| 283 | Primary Cutaneous Peripheral T-Cell Lymphoma in a Sporotrichoid Pattern: A Case Report. <i>Journal of Cutaneous Medicine and Surgery</i> , 2017 , 21, 568-571 | 1.6 | 1 |
| 282 | BRENTUXIMAB VEDOTIN VS PHYSICIAN'S CHOICE IN CTCL PATIENTS FROM THE PHASE 3 ALCANZA STUDY: ANALYSIS OF OUTCOMES BY CD30 EXPRESSION. <i>Hematological Oncology</i> , 2017 , 35, 77-78 | 1.3 | 1 |

| | | | |
|-----|---|------|-----|
| 281 | Oral bexarotene for post-transplant cutaneous T-cell lymphoma. <i>Dermatologic Therapy</i> , 2017 , 30, e12524.2 | | 5 |
| 280 | Mogamulizumab for the treatment of relapsed or refractory adult T-cell leukemia-lymphoma. <i>Expert Review of Hematology</i> , 2017 , 10, 757-760 | 2.8 | 4 |
| 279 | A possible association between mycosis fungoides and Muir-Torre syndrome: Two disorders with microsatellite instability. <i>JAAD Case Reports</i> , 2017 , 3, 358-361 | 1.4 | 1 |
| 278 | RESPONSE BY STAGE IN CD30-POSITIVE (CD30+) CUTANEOUS T CELL LYMPHOMA (CTCL) PATIENTS RECEIVING BRENTUXIMAB VEDOTIN (BV) VS PHYSICIAN'S CHOICE (PC) IN THE PHASE 3 ALCANZA STUDY. <i>Hematological Oncology</i> , 2017 , 35, 245-247 | 1.3 | 1 |
| 277 | PATIENT-REPORTED OUTCOMES AND QUALITY OF LIFE IN PATIENTS WITH CUTANEOUS T CELL LYMPHOMA: RESULTS FROM THE PHASE 3 ALCANZA STUDY. <i>Hematological Oncology</i> , 2017 , 35, 247-248 | 1.3 | 2 |
| 276 | An adolescent with granulomatous mycosis fungoides infiltrating skeletal muscle successfully treated with oral prednisone. <i>JAAD Case Reports</i> , 2017 , 3, 276-279 | 1.4 | 1 |
| 275 | Diverse types of dermatologic toxicities from immune checkpoint blockade therapy. <i>Journal of Cutaneous Pathology</i> , 2017 , 44, 158-176 | 1.7 | 135 |
| 274 | Essential Role of DNA Methyltransferase 1-mediated Transcription of Insulin-like Growth Factor 2 in Resistance to Histone Deacetylase Inhibitors. <i>Clinical Cancer Research</i> , 2017 , 23, 1299-1311 | 12.9 | 20 |
| 273 | Primary cutaneous anaplastic large-cell lymphoma: Complete remission for 13 years after denileukin diftitox. <i>JAAD Case Reports</i> , 2017 , 3, 501-504 | 1.4 | 11 |
| 272 | Recent advances in systemic targeted therapy for cutaneous T-cell lymphoma. <i>Expert Opinion on Pharmacotherapy</i> , 2017 , 18, 1535-1536 | 4 | 4 |
| 271 | Novel Mutations Involving NF- κ B and B-Cell Signaling Pathways in Primary Cutaneous Large B-Cell Lymphoma, Leg-Type and Comparison with Sjögren Syndrome. <i>Journal of Investigative Dermatology</i> , 2017 , 137, 1831-1833 | 4.3 | 5 |
| 270 | TruSeq-Based Gene Expression Analysis of Formalin-Fixed Paraffin-Embedded (FFPE) Cutaneous T-Cell Lymphoma Samples: Subgroup Analysis Results and Elucidation of Biases from FFPE Sample Processing on the TruSeq Platform. <i>Frontiers in Medicine</i> , 2017 , 4, 153 | 4.9 | 12 |
| 269 | Curcumin for the treatment of tumor-stage mycosis fungoides. <i>Dermatologic Therapy</i> , 2017 , 30, e12511 | 2.2 | 3 |
| 268 | ONC201 selectively induces apoptosis in cutaneous T-cell lymphoma cells via activating pro-apoptotic integrated stress response and inactivating JAK/STAT and NF- κ B pathways. <i>Oncotarget</i> , 2017 , 8, 61761-61776 | 3.3 | 16 |
| 267 | The safety profile of vorinostat (suberoylanilide hydroxamic acid) in hematologic malignancies: A review of clinical studies. <i>Cancer Treatment Reviews</i> , 2016 , 43, 58-66 | 14.4 | 40 |
| 266 | Clinicopathological and molecular study of primary cutaneous CD4+ small/medium-sized pleomorphic T-cell lymphoma. <i>Journal of Cutaneous Pathology</i> , 2016 , 43, 1121-1130 | 1.7 | 26 |
| 265 | Loss of CD30 expression after treatment with brentuximab vedotin in a patient with anaplastic large cell lymphoma: a novel finding. <i>Journal of Cutaneous Pathology</i> , 2016 , 43, 1161-1166 | 1.7 | 29 |
| 264 | Primär kutane CD30(+) lymphoproliferative Erkrankungen. <i>JDDG - Journal of the German Society of Dermatology</i> , 2016 , 14, 767-84 | 1.2 | 7 |

| | | | |
|-----|--|------|----|
| 263 | Mogamulizumab in the treatment of cutaneous T cell lymphoma. <i>Expert Opinion on Orphan Drugs</i> , 2016 , 4, 1277-1280 | 1.1 | |
| 262 | Granulomatous Mycosis Fungoides in an Adolescent-A Rare Encounter and Review of the Literature. <i>Pediatric Dermatology</i> , 2016 , 33, e296-8 | 1.9 | 6 |
| 261 | Mogamulizumab for the treatment of cutaneous T-cell lymphoma: recent advances and clinical potential. <i>Therapeutic Advances in Hematology</i> , 2016 , 7, 171-4 | 5.7 | 26 |
| 260 | miR-63/DGCR8-Dependent MicroRNAs Mediate Therapeutic Efficacy of HDAC Inhibitors in Cancer. <i>Cancer Cell</i> , 2016 , 29, 874-888 | 24.3 | 29 |
| 259 | Investigating potential exogenous tumor initiating and promoting factors for Cutaneous T-Cell Lymphomas (CTCL), a rare skin malignancy. <i>Oncolmmunology</i> , 2016 , 5, e1175799 | 7.2 | 22 |
| 258 | BRAF inhibitor therapy-associated melanocytic lesions lack the BRAF V600E mutation and show increased levels of cyclin D1 expression. <i>Human Pathology</i> , 2016 , 50, 79-89 | 3.7 | 16 |
| 257 | Lymphomatoid papulosis: Treatment response and associated lymphomas in a study of 180 patients. <i>Journal of the American Academy of Dermatology</i> , 2016 , 74, 59-67 | 4.5 | 81 |
| 256 | Retrospective Analysis of Prognostic Factors in 187 Cases of Transformed Mycosis Fungoides. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016 , 16, 49-56 | 2 | 38 |
| 255 | A Single-Center Experience With Brentuximab Vedotin in Gamma Delta T-Cell Lymphoma. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016 , 16, e15-9 | 2 | 12 |
| 254 | Brentuximab Vedotin Demonstrates Significantly Superior Clinical Outcomes in Patients with CD30-Expressing Cutaneous T Cell Lymphoma Versus Physician Choice (Methotrexate or Bexarotene): The Phase 3 Alcanza Study. <i>Blood</i> , 2016 , 128, 182-182 | 2.2 | 10 |
| 253 | First-in-Human, Multicenter Phase I Study of IPH4102, First-in-Class Humanized Anti-KIR3DL2 Monoclonal Antibody, in Relapsed/Refractory Cutaneous T-Cell Lymphomas: Preliminary Safety, Exploratory and Clinical Activity Results. <i>Blood</i> , 2016 , 128, 1826-1826 | 2.2 | 3 |
| 252 | Mycosis Fungoides of the Oral Cavity: Fungating Tumor Successfully Treated with Electron Beam Radiation and Maintenance Bexarotene. <i>Case Reports in Dermatological Medicine</i> , 2016 , 2016, 5857935 | 0.8 | 7 |
| 251 | A case of indeterminate dendritic cell tumor presenting with leonine facies. <i>Journal of Cutaneous Pathology</i> , 2016 , 43, 158-63 | 1.7 | 9 |
| 250 | Scleromyxedema: long-term follow-up after high-dose melphalan with autologous stem cell transplantation. <i>International Journal of Dermatology</i> , 2016 , 55, e539-43 | 1.7 | 11 |
| 249 | Molecular signatures define alopecia areata subtypes and transcriptional biomarkers. <i>EBioMedicine</i> , 2016 , 7, 240-7 | 8.8 | 51 |
| 248 | Lymphomatoid Papulosis in Children and Adolescents: A Systematic Review. <i>American Journal of Clinical Dermatology</i> , 2016 , 17, 319-27 | 7.1 | 30 |
| 247 | Clinical Efficacy of Romidepsin in Tumor Stage and Folliculotropic Mycosis Fungoides. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016 , 16, 637-643 | 2 | 15 |
| 246 | Primary cutaneous CD30(+) lymphoproliferative disorders. <i>JDDG - Journal of the German Society of Dermatology</i> , 2016 , 14, 767-82 | 1.2 | 4 |

| | | | |
|-----|--|------|-----|
| 245 | Cutaneous T-cell lymphoma in a patient with celiac disease. <i>Cutis</i> , 2016 , 98, E1-2 | 0.4 | |
| 244 | Characteristics of Sweet Syndrome in patients with acute myeloid leukemia. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2015 , 15, 358-363 | 2 | 37 |
| 243 | An unusual case of cytotoxic peripheral T-cell lymphoma. <i>JAAD Case Reports</i> , 2015 , 1, 257-60 | 1.4 | 4 |
| 242 | Identification of geographic clustering and regions spared by cutaneous T-cell lymphoma in Texas using 2 distinct cancer registries. <i>Cancer</i> , 2015 , 121, 1993-2003 | 6.4 | 34 |
| 241 | Advanced-stage mycosis fungoides and S \ddot{u} ary syndrome: survival and response to treatment. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2015 , 15, e105-12 | 2 | 27 |
| 240 | Clinically significant responses achieved with romidepsin across disease compartments in patients with cutaneous T-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2015 , 56, 2847-54 | 1.9 | 13 |
| 239 | Shared clonality in distinctive lesions of lymphomatoid papulosis and mycosis fungoides occurring in the same patients suggests a common origin. <i>Human Pathology</i> , 2015 , 46, 558-69 | 3.7 | 29 |
| 238 | Histone Deacetylase Inhibitors for Cutaneous T-Cell Lymphoma. <i>Dermatologic Clinics</i> , 2015 , 33, 757-64 | 4.2 | 25 |
| 237 | Results of a Phase II Trial of Brentuximab Vedotin for CD30+ Cutaneous T-Cell Lymphoma and Lymphomatoid Papulosis. <i>Journal of Clinical Oncology</i> , 2015 , 33, 3759-65 | 2.2 | 203 |
| 236 | Depletion of regulatory T cells by targeting CC chemokine receptor type 4 with mogamulizumab. <i>Onc Immunology</i> , 2015 , 4, e1011524 | 7.2 | 22 |
| 235 | Cutaneous Lymphoma International Consortium Study of Outcome in Advanced Stages of Mycosis Fungoides and S \ddot{u} ary Syndrome: Effect of Specific Prognostic Markers on Survival and Development of a Prognostic Model. <i>Journal of Clinical Oncology</i> , 2015 , 33, 3766-73 | 2.2 | 237 |
| 234 | Resimmune, an anti-CD3 \ddot{u} recombinant immunotoxin, induces durable remissions in patients with cutaneous T-cell lymphoma. <i>Haematologica</i> , 2015 , 100, 794-800 | 6.6 | 33 |
| 233 | Leonine facies (LF) and mycosis fungoides (MF): A single-center study and systematic review of the literature. <i>Journal of the American Academy of Dermatology</i> , 2015 , 73, 976-86 | 4.5 | 20 |
| 232 | Allogeneic stem-cell transplantation in patients with cutaneous lymphoma: updated results from a single institution. <i>Annals of Oncology</i> , 2015 , 26, 2490-5 | 10.3 | 63 |
| 231 | Genomic profiling of S \ddot{u} ary syndrome identifies alterations of key T cell signaling and differentiation genes. <i>Nature Genetics</i> , 2015 , 47, 1426-34 | 36.3 | 199 |
| 230 | Long-Term Complete Responses to Combination Therapies and Allogeneic Stem Cell Transplants in Patients With S \ddot{u} ary Syndrome. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2015 , 15, e83-93 | 2 | 30 |
| 229 | Reed syndrome presenting with leiomyosarcoma. <i>JAAD Case Reports</i> , 2015 , 1, 150-2 | 1.4 | 13 |
| 228 | Choosing a systemic treatment for advanced stage cutaneous T-cell lymphoma: mycosis fungoides and S \ddot{u} ary syndrome. <i>Hematology American Society of Hematology Education Program</i> , 2015 , 2015, 529-44 | 2.1 | 11 |

| | | | |
|-----|---|------|-----|
| 227 | Demographic patterns of cutaneous T-cell lymphoma incidence in Texas based on two different cancer registries. <i>Cancer Medicine</i> , 2015 , 4, 1440-7 | 4.8 | 33 |
| 226 | Phase 1/2 study of mogamulizumab, a defucosylated anti-CCR4 antibody, in previously treated patients with cutaneous T-cell lymphoma. <i>Blood</i> , 2015 , 125, 1883-9 | 2.2 | 165 |
| 225 | The effect of extracorporeal photopheresis alone or in combination therapy on circulating CD4(+) Foxp3(+) CD25(-) T cells in patients with leukemic cutaneous T-cell lymphoma. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2015 , 31, 184-94 | 2.4 | 14 |
| 224 | A Phase II trial of Belinostat (PXD101) in patients with relapsed or refractory peripheral or cutaneous T-cell lymphoma. <i>British Journal of Haematology</i> , 2015 , 168, 811-9 | 4.5 | 131 |
| 223 | Low-dose total skin electron beam therapy as an effective modality to reduce disease burden in patients with mycosis fungoides: results of a pooled analysis from 3 phase-II clinical trials. <i>Journal of the American Academy of Dermatology</i> , 2015 , 72, 286-92 | 4.5 | 117 |
| 222 | Lymphomatoid Papulosis: Assessing Treatment Response and Associated Lymphomas in a Study of 180 Patients. <i>Blood</i> , 2015 , 126, 1487-1487 | 2.2 | |
| 221 | Blood Transcriptional Profiling in Patients with Leukemic Cutaneous T-Cell Lymphoma on Extracorporeal Photopheresis Reveals the Integrin Signaling As the Top Pathway Associated with Clinical Response. <i>Blood</i> , 2015 , 126, 3981-3981 | 2.2 | |
| 220 | Vitamin D deficiency in mycosis fungoides and S \bar{z} ary syndrome patients is similar to other cancer patients. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2014 , 14, 518-24 | 2 | 13 |
| 219 | T-cell receptor- $\bar{\gamma}$ in gamma-delta phenotype cutaneous T-cell lymphoma can be accompanied by atypical expression of CD30, CD4, or TCR β 1 and an indolent clinical course. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2014 , 14, e195-200 | 2 | 13 |
| 218 | Clinical presentation, immunopathology, and treatment of juvenile-onset mycosis fungoides: a case series of 34 patients. <i>Journal of the American Academy of Dermatology</i> , 2014 , 71, 1117-26 | 4.5 | 55 |
| 217 | Alopecia areata after HLA-identical BMT from an affected, sibling donor. <i>Bone Marrow Transplantation</i> , 2014 , 49, 592-4 | 4.4 | 4 |
| 216 | Interferon-based treatment for patients with mycosis fungoides and hepatitis C virus infection: a case series. <i>American Journal of Clinical Dermatology</i> , 2014 , 15, 451-6 | 7.1 | 2 |
| 215 | Pralatrexate alone or in combination with bexarotene: long-term tolerability in relapsed/refractory mycosis fungoides. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2014 , 14, 297-304 | 2 | 15 |
| 214 | Results of an open-label multicenter phase 2 trial of lenalidomide monotherapy in refractory mycosis fungoides and S \bar{z} ary syndrome. <i>Blood</i> , 2014 , 123, 1159-66 | 2.2 | 61 |
| 213 | Final results of a multicenter phase II study of the purine nucleoside phosphorylase (PNP) inhibitor forodesine in patients with advanced cutaneous T-cell lymphomas (CTCL) (Mycosis fungoides and S \bar{z} ary syndrome). <i>Annals of Oncology</i> , 2014 , 25, 1807-1812 | 10.3 | 32 |
| 212 | S \bar{z} ary syndrome: an overview of current and future treatment options. <i>Expert Opinion on Orphan Drugs</i> , 2014 , 2, 889-901 | 1.1 | |
| 211 | Concurrent chronic lymphocytic leukemia and cutaneous T cell lymphoma: a case series. <i>Leukemia and Lymphoma</i> , 2014 , 55, 2192-5 | 1.9 | 1 |
| 210 | Mechlorethamine gel for the topical treatment of stage IA and IB mycosis fungoides-type cutaneous T-cell lymphoma. <i>Expert Review of Clinical Pharmacology</i> , 2014 , 7, 591-7 | 3.8 | 14 |

| | | | |
|-----|--|-----|-----|
| 209 | Deregulation in STAT signaling is important for cutaneous T-cell lymphoma (CTCL) pathogenesis and cancer progression. <i>Cell Cycle</i> , 2014 , 13, 3331-5 | 4.7 | 73 |
| 208 | A case of invisible leukemic cutaneous T cell lymphoma with a regulatory T cell clone. <i>International Journal of Dermatology</i> , 2013 , 52, 1111-4 | 1.7 | 10 |
| 207 | A cutaneous lymphoma international prognostic index (CLIPi) for mycosis fungoides and Sezary syndrome. <i>European Journal of Cancer</i> , 2013 , 49, 2859-68 | 7.5 | 87 |
| 206 | Duration of response in cutaneous T-cell lymphoma patients treated with denileukin diftitox: results from 3 phase III studies. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2013 , 13, 377-84 | 2 | 29 |
| 205 | Denileukin diftitox for the treatment of CD25 low-expression mycosis fungoides and Sezary syndrome. <i>Leukemia and Lymphoma</i> , 2013 , 54, 69-75 | 1.9 | 19 |
| 204 | Clinically meaningful reduction in pruritus in patients with cutaneous T-cell lymphoma treated with romidepsin. <i>Leukemia and Lymphoma</i> , 2013 , 54, 284-9 | 1.9 | 34 |
| 203 | Panobinostat activity in both bexarotene-exposed and naïve patients with refractory cutaneous T-cell lymphoma: results of a phase II trial. <i>European Journal of Cancer</i> , 2013 , 49, 386-94 | 7.5 | 104 |
| 202 | Papuloerythroderma of Ofuji associated with an indolent CD3(-), CD4(+), TCR(-) peripheral T-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2013 , 54, 2059-61 | 1.9 | 2 |
| 201 | Hydrochlorothiazide and cutaneous T cell lymphoma: prospective analysis and case series. <i>Cancer</i> , 2013 , 119, 825-31 | 6.4 | 34 |
| 200 | Efficacy and safety of denileukin diftitox retreatment in patients with relapsed cutaneous T-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2013 , 54, 514-9 | 1.9 | 12 |
| 199 | Augmentation of blood dendritic cells by extracorporeal photopheresis in patients with leukemic cutaneous T-cell lymphoma and graft-versus-host disease. <i>Journal of Investigative Dermatology</i> , 2013 , 133, 2098-100 | 4.3 | 18 |
| 198 | Primary cutaneous follicle center lymphoma following deep brain stimulation implants. <i>Leukemia and Lymphoma</i> , 2013 , 54, 1806-7 | 1.9 | |
| 197 | Cutaneous T-Cell Lymphomas: Mycosis Fungoides and Sezary Syndrome 2013 , 287-329 | | 2 |
| 196 | Cutaneous T-Cell Lymphomas 2013 , 155-177 | | |
| 195 | Increased Twist expression in advanced stage of mycosis fungoides and Sezary syndrome. <i>Journal of Cutaneous Pathology</i> , 2012 , 39, 500-7 | 1.7 | 23 |
| 194 | Prevalence and severity of pruritus in cutaneous T cell lymphoma. <i>International Journal of Dermatology</i> , 2012 , 51, 930-4 | 1.7 | 48 |
| 193 | Identification of an active, well-tolerated dose of pralatrexate in patients with relapsed or refractory cutaneous T-cell lymphoma. <i>Blood</i> , 2012 , 119, 4115-22 | 2.2 | 104 |
| 192 | Long-term follow-up and survival of cutaneous T-cell lymphoma patients treated with extracorporeal photopheresis. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2012 , 28, 250-7 | 2.4 | 32 |

| | | | |
|-----|---|------|-----|
| 191 | HDAC inhibitors for the treatment of cutaneous T-cell lymphomas. <i>Future Medicinal Chemistry</i> , 2012 , 4, 471-86 | 4.1 | 54 |
| 190 | Long-term outcomes of 1,263 patients with mycosis fungoides and Sary syndrome from 1982 to 2009. <i>Clinical Cancer Research</i> , 2012 , 18, 5051-60 | 12.9 | 188 |
| 189 | Incidence of spontaneous remission in patients with CD25-positive mycosis fungoides/Sary syndrome receiving placebo. <i>Journal of the American Academy of Dermatology</i> , 2012 , 67, 867-75 | 4.5 | 13 |
| 188 | Pilot study of denileukin diftitox alternate dosing regimen in patients with cutaneous peripheral T-cell lymphomas. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2012 , 12, 180-5 | 2 | 12 |
| 187 | Trends in the treatment of cutaneous T-cell lymphoma – critical evaluation and perspectives on vorinostat. <i>Blood and Lymphatic Cancer: Targets and Therapy</i> , 2012 , 17 | 2.6 | 1 |
| 186 | Cutaneous T-cell lymphomas: a spectrum of presentations with overlap with other cytotoxic lymphomas. <i>American Journal of Surgical Pathology</i> , 2012 , 36, 1656-65 | 6.7 | 119 |
| 185 | Multicenter photopheresis intervention trial in early-stage mycosis fungoides. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2011 , 11, 219-27 | 2 | 30 |
| 184 | Treatment of cutaneous lymphoid hyperplasia with the monoclonal anti-CD20 antibody rituximab. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2011 , 11, 286-8 | 2 | 12 |
| 183 | The spectrum of hair loss in patients with mycosis fungoides and Sary syndrome. <i>Journal of the American Academy of Dermatology</i> , 2011 , 64, 53-63 | 4.5 | 37 |
| 182 | Sary syndrome: immunopathogenesis, literature review of therapeutic options, and recommendations for therapy by the United States Cutaneous Lymphoma Consortium (USCLC). <i>Journal of the American Academy of Dermatology</i> , 2011 , 64, 352-404 | 4.5 | 133 |
| 181 | CO26. Results of a phase 1/2 study for KW-0761, a monoclonal antibody directed against CC chemokine receptor type 4 (CCR4), in CTCL patients. <i>Melanoma Research</i> , 2011 , 21, e15-e16 | 3.3 | 3 |
| 180 | Sary syndrome cells overexpress syndecan-4 bearing distinct heparan sulfate moieties that suppress T-cell activation by binding DC-HIL and trapping TGF-beta on the cell surface. <i>Blood</i> , 2011 , 117, 3382-90 | 2.2 | 28 |
| 179 | EORTC, ISCL, and USCLC consensus recommendations for the treatment of primary cutaneous CD30-positive lymphoproliferative disorders: lymphomatoid papulosis and primary cutaneous anaplastic large-cell lymphoma. <i>Blood</i> , 2011 , 118, 4024-35 | 2.2 | 295 |
| 178 | CD30+ neoplasms of the skin. <i>Current Hematologic Malignancy Reports</i> , 2011 , 6, 245-50 | 4.4 | 55 |
| 177 | Predictors of complete responses with denileukin diftitox in cutaneous T-cell lymphoma. <i>American Journal of Hematology</i> , 2011 , 86, 627-30 | 7.1 | 9 |
| 176 | Clinical end points and response criteria in mycosis fungoides and Sary syndrome: a consensus statement of the International Society for Cutaneous Lymphomas, the United States Cutaneous Lymphoma Consortium, and the Cutaneous Lymphoma Task Force of the European Organisation for Research and Treatment of Cancer. <i>Journal of Clinical Oncology</i> , 2011 , 29, 2598-607 | 2.2 | 407 |
| 175 | Phase II Multicenter Trial of Lenalidomide: Clinical and Immunomodulatory Effects in Patients with CTCL. <i>Blood</i> , 2011 , 118, 1638-1638 | 2.2 | 2 |
| 174 | Dendritic cells and cutaneous T-cell lymphomas. <i>Giornale Italiano Di Dermatologia E Venereologia</i> , 2011 , 146, 103-13 | 0.8 | 8 |

| | | | |
|-----|---|------|-----|
| 173 | Serum T helper 1 cytokine levels are greater in patients with alopecia areata regardless of severity or atopy. <i>Clinical and Experimental Dermatology</i> , 2010 , 35, 409-16 | 1.8 | 50 |
| 172 | Genome-wide association study in alopecia areata implicates both innate and adaptive immunity. <i>Nature</i> , 2010 , 466, 113-7 | 50.4 | 500 |
| 171 | Vorinostat for the treatment of cutaneous T-cell lymphoma. <i>Expert Review of Dermatology</i> , 2010 , 5, 393-401 | | 1 |
| 170 | Total skin electron beam and non-myeloablative allogeneic hematopoietic stem-cell transplantation in advanced mycosis fungoides and Sezary syndrome. <i>Journal of Clinical Oncology</i> , 2010 , 28, 2365-72 | 2.2 | 122 |
| 169 | Curcumin selectively induces apoptosis in cutaneous T-cell lymphoma cell lines and patients M PBMCs: potential role for STAT-3 and NF-kappaB signaling. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 2110-9 | 4.3 | 81 |
| 168 | Rational clinical trial design in cutaneous lymphoma. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2010 , 10 Suppl 2, S80-3 | 2 | |
| 167 | Treatment of Cutaneous T-cell Lymphomas 2010 , 449-457 | | |
| 166 | Phase I trial of a Toll-like receptor 9 agonist, PF-3512676 (CPG 7909), in patients with treatment-refractory, cutaneous T-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 2010 , 63, 975-83 | 4.5 | 79 |
| 165 | Final results from a multicenter, international, pivotal study of romidepsin in refractory cutaneous T-cell lymphoma. <i>Journal of Clinical Oncology</i> , 2010 , 28, 4485-91 | 2.2 | 502 |
| 164 | HR23B is a biomarker for tumor sensitivity to HDAC inhibitor-based therapy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 6532-7 | 11.5 | 129 |
| 163 | Phase III placebo-controlled trial of denileukin diftitox for patients with cutaneous T-cell lymphoma. <i>Journal of Clinical Oncology</i> , 2010 , 28, 1870-7 | 2.2 | 179 |
| 162 | Primary and metastatic malignant tumors of the scalp: an update. <i>American Journal of Clinical Dermatology</i> , 2010 , 11, 233-46 | 7.1 | 18 |
| 161 | A phase II study of SGN-30 in cutaneous anaplastic large cell lymphoma and related lymphoproliferative disorders. <i>Clinical Cancer Research</i> , 2009 , 15, 6217-24 | 12.9 | 73 |
| 160 | Detection of high-risk human papillomaviruses in verrucae of patients with mycosis fungoides and Sezary syndrome: a case series. <i>International Journal of Dermatology</i> , 2009 , 48, 598-602 | 1.7 | 7 |
| 159 | Overall survival in erythrodermic cutaneous T-cell lymphoma: an analysis of prognostic factors in a cohort of patients with erythrodermic cutaneous T-cell lymphoma. <i>International Journal of Dermatology</i> , 2009 , 48, 243-52 | 1.7 | 92 |
| 158 | Lymphomatoid papulosis and associated lymphomas: a retrospective case series of 84 patients. <i>Clinical and Experimental Dermatology</i> , 2009 , 34, 576-81 | 1.8 | 74 |
| 157 | Poor prognosis in non-Caucasian patients with early-onset mycosis fungoides. <i>Journal of the American Academy of Dermatology</i> , 2009 , 60, 231-5 | 4.5 | 55 |
| 156 | History of atopy or autoimmunity increases risk of alopecia areata. <i>Journal of the American Academy of Dermatology</i> , 2009 , 61, 581-91 | 4.5 | 120 |

| | | | |
|-----|--|-----|-----|
| 155 | Evaluation of the long-term tolerability and clinical benefit of vorinostat in patients with advanced cutaneous T-cell lymphoma. <i>Clinical Lymphoma and Myeloma</i> , 2009 , 9, 412-6 | | 74 |
| 154 | Efficacy and safety of topical tazarotene: a review. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2009 , 5, 195-210 | 5.5 | 18 |
| 153 | Molecular and Biological Characteristics of Acquired Vorinostat-Resistant Cutaneous T-Cell Lymphoma Cells.. <i>Blood</i> , 2009 , 114, 1732-1732 | 2.2 | |
| 152 | Primary Sezary Syndrome Commonly Involves Skin Without Epidermotropism. <i>Journal of Cutaneous Pathology</i> , 2008 , 32, 85-85 | 1.7 | |
| 151 | Induction of T-cell responses against cutaneous T-cell lymphomas ex vivo by autologous dendritic cells transfected with amplified tumor mRNA. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 2631-2639 | 4.3 | 22 |
| 150 | Avicin D selectively induces apoptosis and downregulates p-STAT-3, bcl-2, and survivin in cutaneous T-cell lymphoma cells. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 2728-2735 | 4.3 | 37 |
| 149 | Human leukocyte antigen class II alleles are associated with risk of alopecia areata. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 240-3 | 4.3 | 23 |
| 148 | Prevalence and treatment of Staphylococcus aureus colonization in patients with mycosis fungoides and Sezary syndrome. <i>British Journal of Dermatology</i> , 2008 , 159, 105-12 | 4 | 130 |
| 147 | Cutaneous gamma/delta T-cell lymphoma treated with radiation and denileukin diftitox. <i>Clinical Lymphoma and Myeloma</i> , 2008 , 8, 55-8 | | 16 |
| 146 | Increased malignancy risk in the cutaneous T-cell lymphoma patient population. <i>Clinical Lymphoma and Myeloma</i> , 2008 , 8, 100-5 | | 25 |
| 145 | Central nervous system involvement in patients with mycosis fungoides and cutaneous large-cell transformation. <i>Journal of the American Academy of Dermatology</i> , 2008 , 59, S16-22 | 4.5 | 18 |
| 144 | EGFR inhibitor-associated acneiform folliculitis: assessment and management. <i>American Journal of Clinical Dermatology</i> , 2008 , 9, 285-94 | 7.1 | 23 |
| 143 | Optimizing denileukin diftitox (Ontak) therapy. <i>Future Oncology</i> , 2008 , 4, 457-69 | 3.6 | 46 |
| 142 | Review of the treatment of mycosis fungoides and sezary syndrome: a stage-based approach. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2008 , 6, 436-42 | 7.3 | 97 |
| 141 | Skin directed therapy for mycosis fungoides: a review. <i>Journal of Drugs in Dermatology</i> , 2008 , 7, 655-66 | 2.2 | 27 |
| 140 | Pulmonary manifestations in patients with cutaneous T-cell lymphomas. <i>Cancer</i> , 2007 , 109, 1550-5 | 6.4 | 15 |
| 139 | The optimal use of bexarotene in cutaneous T-cell lymphoma. <i>British Journal of Dermatology</i> , 2007 , 157, 433-40 | 4 | 111 |
| 138 | Mycosis fungoides: pathophysiology and emerging therapies. <i>Seminars in Oncology</i> , 2007 , 34, S21-8 | 5.5 | 33 |

| | | | |
|-----|--|-----|------|
| 137 | Activation of HIV in Human Skin by Ultraviolet B Radiation and its Inhibition by NFB Blocking Agents¶. <i>Photochemistry and Photobiology</i> , 2007 , 74, 805-810 | 3.6 | 2 |
| 136 | Treatment of transformed mycosis fungoides with intermittent low-dose gemcitabine. <i>Oncology</i> , 2007 , 73, 130-5 | 3.6 | 8 |
| 135 | Phase IIb multicenter trial of vorinostat in patients with persistent, progressive, or treatment refractory cutaneous T-cell lymphoma. <i>Journal of Clinical Oncology</i> , 2007 , 25, 3109-15 | 2.2 | 835 |
| 134 | Phase 2 trial of oral vorinostat (suberoylanilide hydroxamic acid, SAHA) for refractory cutaneous T-cell lymphoma (CTCL). <i>Blood</i> , 2007 , 109, 31-9 | 2.2 | 924 |
| 133 | Lesional gene expression profiling in cutaneous T-cell lymphoma reveals natural clusters associated with disease outcome. <i>Blood</i> , 2007 , 110, 3015-27 | 2.2 | 96 |
| 132 | Clinical efficacy of zanolimumab (HuMax-CD4): two phase 2 studies in refractory cutaneous T-cell lymphoma. <i>Blood</i> , 2007 , 109, 4655-62 | 2.2 | 168 |
| 131 | Revisions to the staging and classification of mycosis fungoides and Sezary syndrome: a proposal of the International Society for Cutaneous Lymphomas (ISCL) and the cutaneous lymphoma task force of the European Organization of Research and Treatment of Cancer (EORTC). <i>Blood</i> , 2007 , 110, 1713-22 | 2.2 | 1012 |
| 130 | Vorinostat: a new oral histone deacetylase inhibitor approved for cutaneous T-cell lymphoma. <i>Expert Opinion on Investigational Drugs</i> , 2007 , 16, 1111-20 | 5.9 | 248 |
| 129 | Coexistent psoriasis and lupus erythematosus treated with alefacept. <i>American Journal of Clinical Dermatology</i> , 2007 , 8, 47-50 | 7.1 | 6 |
| 128 | Vorinostat in cutaneous T-cell lymphoma. <i>Drugs of Today</i> , 2007 , 43, 585-99 | 2.5 | 43 |
| 127 | Update on the treatment of cutaneous T-cell lymphoma (CTCL): Focus on vorinostat. <i>Biologics: Targets and Therapy</i> , 2007 , 1, 377-92 | 4.4 | 27 |
| 126 | Retinoids in Cutaneous T-Cell Lymphomas. <i>Basic and Clinical Dermatology</i> , 2007 , 183-196 | | 1 |
| 125 | Overall Survival in Erythrodermic Cutaneous T-Cell Lymphoma: An Analysis of Prognostic Factors in a Cohort of Erythrodermic Cutaneous T-Cell Lymphoma Patients.. <i>Blood</i> , 2007 , 110, 3574-3574 | 2.2 | |
| 124 | Systemic monotherapy vs combination therapy for CTCL: rationale and future strategies. <i>Oncology</i> , 2007 , 21, 33-40 | 1.8 | 8 |
| 123 | Thyrotoxicosis after denileukin diftitox therapy in patients with mycosis fungoides. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006 , 91, 2205-8 | 5.6 | 25 |
| 122 | Photopheresis versus transimmunization?. <i>Leukemia and Lymphoma</i> , 2006 , 47, 1449 | 1.9 | 2 |
| 121 | Treatment of mycosis fungoides with denileukin diftitox and oral bexarotene. <i>Clinical Lymphoma and Myeloma</i> , 2006 , 6, 488-92 | | 8 |
| 120 | Phase II evaluation of gemcitabine monotherapy for cutaneous T-cell lymphoma. <i>Clinical Lymphoma and Myeloma</i> , 2006 , 7, 51-8 | | 119 |

| | | | |
|-----|---|------|-----|
| 119 | Mycosis fungoides with coccidioidomycosis. <i>Clinical Lymphoma and Myeloma</i> , 2006 , 7, 148-50 | | 5 |
| 118 | A phase II open-label study of recombinant human interleukin-12 in patients with stage IA, IB, or IIA mycosis fungoides. <i>Journal of the American Academy of Dermatology</i> , 2006 , 55, 807-13 | 4.5 | 52 |
| 117 | High flotillin-2 expression is associated with lymph node metastasis and Breslow depth in melanoma. <i>Melanoma Research</i> , 2006 , 16, 461-3 | 3.3 | 26 |
| 116 | Treatment of cutaneous T-cell lymphoma with retinoids. <i>Dermatologic Therapy</i> , 2006 , 19, 264-71 | 2.2 | 64 |
| 115 | Clinical and laboratory experience of vorinostat (suberoylanilide hydroxamic acid) in the treatment of cutaneous T-cell lymphoma. <i>British Journal of Cancer</i> , 2006 , 95, S13-S19 | 8.7 | 27 |
| 114 | CD25 expression is correlated with histological grade and response to denileukin diftitox in cutaneous T-cell lymphoma. <i>Journal of Investigative Dermatology</i> , 2006 , 126, 575-83 | 4.3 | 87 |
| 113 | Oral Forodesine (Bcx-1777) Is Clinically Active in Refractory Cutaneous T-Cell Lymphoma: Results of a Phase I/II Study.. <i>Blood</i> , 2006 , 108, 2467-2467 | 2.2 | 10 |
| 112 | Vorinostat (suberoylanilide hydroxamic acid, SAHA) is clinically active in advanced cutaneous T-cell lymphoma (CTCL): Results of a phase IIb trial. <i>Journal of Clinical Oncology</i> , 2006 , 24, 7500-7500 | 2.2 | 19 |
| 111 | Induction of Polyclonal T-Cell Responses Against Cutaneous T-Cell Lymphomas by Autologous Dendritic Cells Transfected with Amplified Tumor RNA.. <i>Blood</i> , 2006 , 108, 5189-5189 | 2.2 | |
| 110 | Primary cutaneous B-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 2005 , 53, 479-84 | 4.5 | 46 |
| 109 | Resistance to activation-induced cell death and bystander cytotoxicity via the Fas/Fas ligand pathway are implicated in the pathogenesis of cutaneous T cell lymphomas. <i>Journal of Investigative Dermatology</i> , 2005 , 124, 741-50 | 4.3 | 38 |
| 108 | Selective induction of apoptosis by histone deacetylase inhibitor SAHA in cutaneous T-cell lymphoma cells: relevance to mechanism of therapeutic action. <i>Journal of Investigative Dermatology</i> , 2005 , 125, 1045-52 | 4.3 | 198 |
| 107 | Degree of CD25 expression in T-cell lymphoma is dependent on tissue site: implications for targeted therapy. <i>Clinical Cancer Research</i> , 2004 , 10, 5587-94 | 12.9 | 28 |
| 106 | Up-regulation of Flotillin-2 is associated with melanoma progression and modulates expression of the thrombin receptor protease activated receptor 1. <i>Cancer Research</i> , 2004 , 64, 7361-9 | 10.1 | 84 |
| 105 | Bone marrow transplantation in Jehovah's Witnesses. <i>Leukemia and Lymphoma</i> , 2004 , 45, 635-6 | 1.9 | 3 |
| 104 | The novel synthetic oleanane triterpenoid CDDO (2-cyano-3, 12-dioxoolean-1, 9-dien-28-oic acid) induces apoptosis in Mycosis fungoides/Szary syndrome cells. <i>Journal of Investigative Dermatology</i> , 2004 , 123, 380-7 | 4.3 | 24 |
| 103 | Phase II study of pentostatin in advanced T-cell lymphoid malignancies: update of an M.D. Anderson Cancer Center series. <i>Cancer</i> , 2004 , 100, 342-9 | 6.4 | 66 |
| 102 | Lymphomatoid papulosis from childhood with anaplastic large-cell lymphoma of the small bowel. <i>Clinical Lymphoma and Myeloma</i> , 2004 , 5, 190-3 | | 11 |

| | | | |
|-----|---|-----|-----|
| 101 | Pilot study of etanercept in patients with relapsed cutaneous T-cell lymphomas. <i>Journal of the American Academy of Dermatology</i> , 2004 , 51, 200-4 | 4.5 | 23 |
| 100 | Tazarotene 0.1% gel for refractory mycosis fungoides lesions: an open-label pilot study. <i>Journal of the American Academy of Dermatology</i> , 2004 , 50, 600-7 | 4.5 | 49 |
| 99 | Cutaneous T-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 2004 , 51, S43-5 | 4.5 | 10 |
| 98 | A systematic approach to diagnosis of mature T-cell leukemias reveals heterogeneity among WHO categories. <i>Blood</i> , 2004 , 104, 328-35 | 2.2 | 153 |
| 97 | Extracorporeal Photopheresis for the Treatment of Cutaneous T-Cell Lymphoma. <i>Journal of Cutaneous Medicine and Surgery</i> , 2003 , 7, 3-7 | 1.6 | 33 |
| 96 | Cytomegalovirus seropositivity is significantly associated with mycosis fungoides and Sézary syndrome. <i>Blood</i> , 2003 , 101, 2132-6 | 2.2 | 67 |
| 95 | Extracorporeal photopheresis for the treatment of cutaneous T-cell lymphoma. <i>Journal of Cutaneous Medicine and Surgery</i> , 2003 , 7, 3-7 | 1.6 | 25 |
| 94 | Mycosis fungoides associated with malignant melanoma and dysplastic nevus syndrome. <i>International Journal of Dermatology</i> , 2003 , 42, 116-22 | 1.7 | 29 |
| 93 | Long-term control of mycosis fungoides of the hands with topical bexarotene. <i>International Journal of Dermatology</i> , 2003 , 42, 238-41 | 1.7 | 6 |
| 92 | Tazarotene-induced gene 3 is suppressed in basal cell carcinomas and reversed in vivo by tazarotene application. <i>Journal of Investigative Dermatology</i> , 2003 , 121, 902-9 | 4.3 | 33 |
| 91 | Alopecia areata registry: an overview. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 2003 , 8, 219-21 | 1.1 | 18 |
| 90 | Retinoids: therapeutic applications and mechanisms of action in cutaneous T-cell lymphoma. <i>Dermatologic Therapy</i> , 2003 , 16, 322-30 | 2.2 | 59 |
| 89 | Topical and systemic retinoid therapy for cutaneous T-cell lymphoma. <i>Hematology/Oncology Clinics of North America</i> , 2003 , 17, 1405-19 | 3.1 | 30 |
| 88 | Analysis of long-term outcomes of combined modality therapy for cutaneous T-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 2003 , 49, 35-49 | 4.5 | 58 |
| 87 | Mycosis fungoides with follicular mucinosis displaying aggressive tumor-stage transformation : successful treatment using radiation therapy plus oral bexarotene combination therapy. <i>American Journal of Clinical Dermatology</i> , 2003 , 4, 429-33 | 7.1 | 11 |
| 86 | 003UVB dose and cell cycle conditions influence the outcome of UVB-induced HIV activation. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2002 , 18, 104-104 | 2.4 | |
| 85 | A stable aberrant immunophenotype characterizes nearly all cases of cutaneous T-cell lymphoma in blood and can be used to monitor response to therapy. <i>BMC Clinical Pathology</i> , 2002 , 2, 5 | 3 | 50 |
| 84 | Treatment of refractory peripheral T-cell lymphoma with denileukin diftitox (ONTAK). <i>Leukemia and Lymphoma</i> , 2002 , 43, 121-6 | 1.9 | 52 |

| | | | |
|----|--|-----|-----|
| 83 | Phase 1 and 2 trial of bexarotene gel for skin-directed treatment of patients with cutaneous T-cell lymphoma. <i>Archives of Dermatology</i> , 2002 , 138, 325-32 | | 134 |
| 82 | Clonal heterogeneity in mycosis fungoides and its relationship to clinical course. <i>Blood</i> , 2002 , 100, 3369-73 | | 113 |
| 81 | Quality-of-life improvements in cutaneous T-cell lymphoma patients treated with denileukin diftitox (ONTAK). <i>Clinical Lymphoma and Myeloma</i> , 2002 , 2, 222-8 | | 80 |
| 80 | Treatment of cutaneous T cell lymphoma: current status and future directions. <i>American Journal of Clinical Dermatology</i> , 2002 , 3, 193-215 | 7.1 | 54 |
| 79 | Update on erythrodermic cutaneous T-cell lymphoma: report of the International Society for Cutaneous Lymphomas. <i>Journal of the American Academy of Dermatology</i> , 2002 , 46, 95-106 | 4.5 | 374 |
| 78 | Cold urticaria in a patient with mycosis fungoides. <i>Journal of the American Academy of Dermatology</i> , 2002 , 47, 608-10 | 4.5 | 4 |
| 77 | Optimizing bexarotene therapy for cutaneous T-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 2002 , 47, 672-84 | 4.5 | 142 |
| 76 | Increased serum immunoglobulin levels are common in mycosis fungoides and Sjögren syndrome. <i>Journal of the American Academy of Dermatology</i> , 2002 , 47, 685-91 | 4.5 | 12 |
| 75 | The role for interleukin-12 therapy of cutaneous T cell lymphoma. <i>Annals of the New York Academy of Sciences</i> , 2001 , 941, 177-84 | 6.5 | 44 |
| 74 | Persistent atypical lymphocytic hyperplasia following tick bite in a child: report of a case and review of the literature. <i>Pediatric Dermatology</i> , 2001 , 18, 481-4 | 1.9 | 27 |
| 73 | Transient CD30+ nodal transformation of cutaneous T-cell lymphoma associated with cyclosporine treatment. <i>International Journal of Dermatology</i> , 2001 , 40, 505-11 | 1.7 | 21 |
| 72 | The T-cell chemokine receptor CXCR3 is expressed highly in low-grade mycosis fungoides. <i>American Journal of Clinical Pathology</i> , 2001 , 115, 413-21 | 1.9 | 70 |
| 71 | Absence of CD26 expression is a useful marker for diagnosis of T-cell lymphoma in peripheral blood. <i>American Journal of Clinical Pathology</i> , 2001 , 115, 885-92 | 1.9 | 153 |
| 70 | The genetics of alopecia areata. <i>Clinics in Dermatology</i> , 2001 , 19, 135-9 | 3 | 22 |
| 69 | Cutaneous T-cell lymphoma. New immunomodulators. <i>Dermatologic Clinics</i> , 2001 , 19, 737-48 | 4.2 | 10 |
| 68 | Therapy options in cutaneous T-cell lymphoma. <i>Expert Review of Anticancer Therapy</i> , 2001 , 1, 403-20 | 3.5 | 13 |
| 67 | Bexarotene is effective and safe for treatment of refractory advanced-stage cutaneous T-cell lymphoma: multinational phase II-III trial results. <i>Journal of Clinical Oncology</i> , 2001 , 19, 2456-71 | 2.2 | 562 |
| 66 | Pivotal phase III trial of two dose levels of denileukin diftitox for the treatment of cutaneous T-cell lymphoma. <i>Journal of Clinical Oncology</i> , 2001 , 19, 376-88 | 2.2 | 534 |

| | | | |
|----|--|------|-----|
| 65 | Phase 2 and 3 clinical trial of oral bexarotene (Targretin capsules) for the treatment of refractory or persistent early-stage cutaneous T-cell lymphoma. <i>Archives of Dermatology</i> , 2001 , 137, 581-93 | | 256 |
| 64 | Fas ligand expression by neoplastic T lymphocytes mediates elimination of CD8+ cytotoxic T lymphocytes in mycosis fungoides: a potential mechanism of tumor immune escape?. <i>Clinical Cancer Research</i> , 2001 , 7, 2682-92 | 12.9 | 41 |
| 63 | Topical treatment of cutaneous lesions of acquired immunodeficiency syndrome-related Kaposi sarcoma using alitretinoin gel: results of phase 1 and 2 trials. <i>Archives of Dermatology</i> , 2000 , 136, 1461-9 | | 51 |
| 62 | Cytokine expression patterns distinguish HIV associated skin diseases. <i>Experimental Dermatology</i> , 2000 , 9, 341-50 | 4 | 12 |
| 61 | Alitretinoin. <i>American Journal of Clinical Dermatology</i> , 2000 , 1, 315-316 | 7.1 | |
| 60 | Emerging new therapies for cutaneous T-cell lymphoma. <i>Dermatologic Clinics</i> , 2000 , 18, 147-56 | 4.2 | 50 |
| 59 | Enterococcal eschars in cutaneous T-cell lymphoma tumors: a distinct clinical entity. <i>Clinical Lymphoma and Myeloma</i> , 2000 , 1, 141-5 | | 9 |
| 58 | Treatment of cutaneous T-cell lymphoma from a dermatologist's perspective. <i>Clinical Lymphoma and Myeloma</i> , 2000 , 1 Suppl 1, S15-20 | | 6 |
| 57 | Bexarotene and DAB(389)IL-2 (denileukin diftitox, ONTAK) in treatment of cutaneous T-cell lymphomas: algorithms. <i>Clinical Lymphoma and Myeloma</i> , 2000 , 1 Suppl 1, S51-5 | | 19 |
| 56 | Expression of a retinoid-inducible tumor suppressor, Tazarotene-inducible gene-3, is decreased in psoriasis and skin cancer. <i>Clinical Cancer Research</i> , 2000 , 6, 3249-59 | 12.9 | 52 |
| 55 | Pentostatin therapy of T-cell lymphomas with cutaneous manifestations. <i>Journal of Clinical Oncology</i> , 1999 , 17, 3117-21 | 2.2 | 88 |
| 54 | Alopecia areata in families: association with the HLA locus. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 1999 , 4, 220-3 | 1.1 | 67 |
| 53 | Central hypothyroidism associated with retinoid X receptor-selective ligands. <i>New England Journal of Medicine</i> , 1999 , 340, 1075-9 | 59.2 | 237 |
| 52 | Flotillin 2 is distinct from epidermal surface antigen (ESA) and is associated with filopodia formation 1999 , 75, 147-159 | | 25 |
| 51 | Prognostic factor analysis in mycosis fungoides/Sézary syndrome. <i>Journal of the American Academy of Dermatology</i> , 1999 , 40, 914-24 | 4.5 | 133 |
| 50 | Dysregulated synthesis of intracellular type 1 and type 2 cytokines by T cells of patients with cutaneous T-cell lymphoma. <i>Vaccine Journal</i> , 1999 , 6, 79-84 | | 48 |
| 49 | Effectiveness of photopheresis in Sézary syndrome. <i>Archives of Dermatology</i> , 1999 , 135, 995-7 | | 24 |
| 48 | DAB389IL2 diphtheria fusion toxin produces clinical responses in tumor stage cutaneous T cell lymphoma. <i>American Journal of Hematology</i> , 1998 , 58, 87-90 | 7.1 | 52 |

| | | | |
|----|---|------|-----|
| 47 | Nerve growth factor in normal and psoriatic skin equivalent models. <i>Archives of Dermatological Research</i> , 1998 , 290, 610-4 | 3.3 | 4 |
| 46 | The pathogenesis of psoriasis and the mechanism of action of tazarotene. <i>Journal of the American Academy of Dermatology</i> , 1998 , 39, S129-33 | 4.5 | 41 |
| 45 | Alopecia areata and cytomegalovirus infection in twins: genes versus environment?. <i>Journal of the American Academy of Dermatology</i> , 1998 , 38, 418-25 | 4.5 | 101 |
| 44 | Tumor necrosis factor-alpha and interferon-gamma, but not HTLV-I tax, are likely factors in the epidermotropism of cutaneous T-cell lymphoma via induction of interferon-inducible protein-10. <i>Leukemia and Lymphoma</i> , 1998 , 29, 315-28 | 1.9 | 19 |
| 43 | Transformation of Mycosis Fungoides/Sezary Syndrome: Clinical Characteristics and Prognosis. <i>Blood</i> , 1998 , 92, 1150-1159 | 2.2 | 243 |
| 42 | Identification and characterization of a retinoid-induced class II tumor suppressor/growth regulatory gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 14811-5 | 11.5 | 132 |
| 41 | Transformation of mycosis fungoides/Sezary syndrome: clinical characteristics and prognosis. <i>Blood</i> , 1998 , 92, 1150-9 | 2.2 | 31 |
| 40 | Tazarotene: a review of its pharmacological profile and potential for clinical use in psoriasis. <i>Expert Opinion on Investigational Drugs</i> , 1997 , 6, 1537-51 | 5.9 | 13 |
| 39 | Tazarotene gel, a new retinoid, for topical therapy of psoriasis: vehicle-controlled study of safety, efficacy, and duration of therapeutic effect. <i>Journal of the American Academy of Dermatology</i> , 1997 , 37, 85-92 | 4.5 | 141 |
| 38 | Follicular mucinosis associated with scarring alopecia, oligoclonal T-cell receptor V beta expansion, and Staphylococcus aureus: when does follicular mucinosis become mycosis fungoides?. <i>Journal of the American Academy of Dermatology</i> , 1997 , 37, 828-31 | 4.5 | 49 |
| 37 | Molecular mechanisms of tazarotene action in psoriasis. <i>Journal of the American Academy of Dermatology</i> , 1997 , 37, S18-S24 | 4.5 | 78 |
| 36 | Association of Erythrodermic Cutaneous T-Cell Lymphoma, Superantigen-Positive Staphylococcus aureus, and Oligoclonal T-Cell Receptor V Gene Expansion. <i>Blood</i> , 1997 , 89, 32-40 | 2.2 | 191 |
| 35 | Tazarotene-induced gene 2 (TIG2), a novel retinoid-responsive gene in skin. <i>Journal of Investigative Dermatology</i> , 1997 , 109, 91-5 | 4.3 | 194 |
| 34 | The sign of Leser-Trélat. <i>Medical and Pediatric Oncology</i> , 1997 , 28, 234-7 | | 5 |
| 33 | Association of erythrodermic cutaneous T-cell lymphoma, superantigen-positive Staphylococcus aureus, and oligoclonal T-cell receptor V beta gene expansion. <i>Blood</i> , 1997 , 89, 32-40 | 2.2 | 61 |
| 32 | Interferon-inducible protein 10 as a possible factor in the pathogenesis of cutaneous T-cell lymphomas. <i>Clinical Cancer Research</i> , 1997 , 3, 169-77 | 12.9 | 15 |
| 31 | Interferon-inducible protein-10 and the pathogenesis of cutaneous T-cell lymphomas. <i>Leukemia and Lymphoma</i> , 1996 , 24, 103-10 | 1.9 | 9 |
| 30 | Combined modality therapy for cutaneous T-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 1996 , 34, 1022-9 | 4.5 | 80 |

| | | | |
|----|--|-----|-----|
| 29 | Photopheresis therapy for cutaneous T-cell lymphoma. <i>Journal of the American Academy of Dermatology</i> , 1996 , 35, 573-9 | 4.5 | 120 |
| 28 | Effects of fibroblast-keratinocyte interactions on the secretion of interleukin-8. <i>Bulletin of Experimental Biology and Medicine</i> , 1996 , 121, 167-170 | 0.8 | 2 |
| 27 | Tazarotene-induced gene 1 (TIG1), a novel retinoic acid receptor-responsive gene in skin. <i>Journal of Investigative Dermatology</i> , 1996 , 106, 269-74 | 4.3 | 96 |
| 26 | HLA-DR5 and DQB1*03 class II alleles are associated with cutaneous T-cell lymphoma. <i>Journal of Investigative Dermatology</i> , 1996 , 107, 373-6 | 4.3 | 71 |
| 25 | Amplification of genomic DNA demonstrates the presence of the t(2;5) (p23;q35) in anaplastic large cell lymphoma, but not in other non-Hodgkin lymphomas, Hodgkin disease, or lymphomatoid papulosis. <i>Blood</i> , 1996 , 88, 1771-1779 | 2.2 | 57 |
| 24 | Amplification of genomic DNA demonstrates the presence of the t(2;5) (p23;q35) in anaplastic large cell lymphoma, but not in other non-Hodgkin lymphomas, Hodgkin disease, or lymphomatoid papulosis. <i>Blood</i> , 1996 , 88, 1771-1779 | 2.2 | 2 |
| 23 | Amplification of genomic DNA demonstrates the presence of the t(2;5) (p23;q35) in anaplastic large cell lymphoma, but not in other non-Hodgkin lymphomas, Hodgkin disease, or lymphomatoid papulosis. <i>Blood</i> , 1996 , 88, 1771-9 | 2.2 | 12 |
| 22 | Cytokine loops involving interferon-gamma and IP-10, a cytokine chemotactic for CD4+ lymphocytes: an explanation for the epidermotropism of cutaneous T-cell lymphoma? [see comments]. <i>Blood</i> , 1995 , 86, 651-658 | 2.2 | 72 |
| 21 | Analysis of HLA-D locus alleles in alopecia areata patients and families. <i>Journal of Investigative Dermatology</i> , 1995 , 104, 5S-6S | 4.3 | 34 |
| 20 | Epidermal surface antigen (MS17S1) is highly conserved between mouse and human. <i>Genomics</i> , 1995 , 27, 251-8 | 4.3 | 12 |
| 19 | Werner syndrome. <i>Dermatologic Clinics</i> , 1995 , 13, 163-8 | 4.2 | 6 |
| 18 | 2-Chlorodeoxyadenosine therapy in patients with T-cell lymphoproliferative disorders. <i>Blood</i> , 1994 , 84, 733-738 | 2.2 | 53 |
| 17 | Keratinocyte transglutaminase expression varies in squamous cell carcinomas. <i>Journal of Investigative Dermatology</i> , 1994 , 102, 462-9 | 4.3 | 21 |
| 16 | In vivo fluorescence spectroscopy and imaging of human skin tumours. <i>Lasers in Medical Science</i> , 1994 , 9, 191-201 | 3.1 | 87 |
| 15 | Human leukocyte antigen-DQB1*03 alleles are associated with alopecia areata. <i>Journal of Investigative Dermatology</i> , 1994 , 103, 758-63 | 4.3 | 80 |
| 14 | High-frequency sonography of the skin. <i>Journal of the European Academy of Dermatology and Venereology</i> , 1994 , 3, 47-55 | 4.6 | 4 |
| 13 | Response of psoriasis to a new topical retinoid, AGN 190168. <i>Journal of the American Academy of Dermatology</i> , 1994 , 30, 581-90 | 4.5 | 112 |
| 12 | Emergence of leprosy in a patient with mycosis fungoides. <i>Journal of the American Academy of Dermatology</i> , 1994 , 30, 313-5 | 4.5 | 12 |

| | | | |
|----|---|-----|-----|
| 11 | 2-Chlorodeoxyadenosine therapy in patients with T-cell lymphoproliferative disorders. <i>Blood</i> , 1994 , 84, 733-8 | 2.2 | 7 |
| 10 | Cloning and characterization of a novel epidermal cell surface antigen (ESA). <i>Journal of Biological Chemistry</i> , 1994 , 269, 19983-91 | 5.4 | 25 |
| 9 | 2-Chlorodeoxyadenosine therapy in patients with T-cell lymphoproliferative disorders. <i>Blood</i> , 1994 , 84, 733-738 | 2.2 | 2 |
| 8 | Cloning and characterization of a novel epidermal cell surface antigen (ESA).. <i>Journal of Biological Chemistry</i> , 1994 , 269, 19983-19991 | 5.4 | 28 |
| 7 | The gene for a novel epidermal antigen maps near the neurofibromatosis 1 gene. <i>Genomics</i> , 1992 , 14, 369-76 | 4.3 | 9 |
| 6 | Blue rubber bleb nevus syndrome: laser photocoagulation of colonic hemangiomas in a child with microcytic anemia. <i>Pediatric Dermatology</i> , 1992 , 9, 91-4 | 1.9 | 23 |
| 5 | HLA-D Locus Associations in Alopecia Areata. <i>Archives of Dermatology</i> , 1991 , 127, 64 | | 29 |
| 4 | The human gene for an epidermal surface antigen (M17S1) is located at 17q11-12. <i>Genomics</i> , 1991 , 11, 481-2 | 4.3 | 7 |
| 3 | Human immunodeficiency virus-associated psoriasis, psoriatic arthritis, and Reiter's syndrome: a disease continuum?. <i>Arthritis and Rheumatism</i> , 1990 , 33, 1574-8 | | 94 |
| 2 | Expression of pemphigus vulgaris antigen in cultured human keratinocytes: effect of inhibitors, tunicamycin, and lectins. <i>Journal of Investigative Dermatology</i> , 1988 , 90, 708-15 | 4.3 | 10 |
| 1 | Use of monospecific antisera and cRNA probes to localize the major changes in keratin expression during normal and abnormal epidermal differentiation. <i>Journal of Cell Biology</i> , 1988 , 107, 427-46 | 7.3 | 326 |