

# Ulrich SchmÄjcke

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6689943/publications.pdf>

Version: 2024-02-01

38  
papers

979  
citations

567281

15  
h-index

454955

30  
g-index

38  
all docs

38  
docs citations

38  
times ranked

1484  
citing authors

#	ARTICLE	IF	CITATIONS
1	Finding Mesolithic Sites: A Multichannel Ground-Penetrating Radar (GPR) Investigation at the Ancient Lake Duvensee. <i>Remote Sensing</i> , 2022, 14, 781.	4.0	4
2	The Baltic grey seal: A 9000-year history of presence and absence. <i>Holocene</i> , 2022, 32, 569-577.	1.7	3
3	Cattle husbandry and aurochs hunting in the Neolithic of northern Central Europe and southern Scandinavia. A statistical approach to distinguish between domestic and wild forms. <i>International Journal of Osteoarchaeology</i> , 2021, 31, 108-118.	1.2	1
4	Winter temperature and forest cover have shaped red deer distribution in Europe and the Ural Mountains since the Late Pleistocene. <i>Journal of Biogeography</i> , 2021, 48, 147-159.	3.0	26
5	Reconstructing the ecological history of the extinct harp seal population of the Baltic Sea. <i>Quaternary Science Reviews</i> , 2021, 251, 106701.	3.0	10
6	Historical Demographic Processes Dominate Genetic Variation in Ancient Atlantic Cod Mitogenomes. <i>Frontiers in Ecology and Evolution</i> , 2021, 9, .	2.2	9
7	A 5,000-year-old hunter-gatherer already plagued by <i>Yersinia pestis</i> . <i>Cell Reports</i> , 2021, 35, 109278.	6.4	42
8	Duvensee WP 10 – an Early Mesolithic Site at Ancient Lake Duvensee, Germany. <i>Journal of Wetland Archaeology</i> , 2021, 21, 1-20.	1.2	6
9	Stone Age fishing strategies in a dynamic river landscape: Evidence from Veksa 3, Northwest Russia. <i>Quaternary International</i> , 2020, 541, 23-40.	1.5	2
10	Two burials in a unique freshwater shell midden: insights into transformations of Stone Age hunter-fisher daily life in Latvia. <i>Archaeological and Anthropological Sciences</i> , 2020, 12, 1.	1.8	13
11	Multiproxy palaeontological investigations of Holocene sediments in the harbour area of the Hanseatic town Stralsund, North-Eastern Germany, southern Baltic Sea coast. <i>Quaternary International</i> , 2019, 511, 22-42.	1.5	3
12	Adaptations and transformations of hunter-gatherers in forest environments: New archaeological and anthropological insights. <i>Holocene</i> , 2019, 29, 1531-1544.	1.7	21
13	Mittel- bis jungneolithische Siedlungshinterlassenschaften zwischen 3300–2600 v. Chr. Der Fundplatz Oldenburg LA 232 im Oldenburger Graben, Ostholstein. <i>Praehistorische Zeitschrift</i> , 2019, 93, 185-224.	0.4	6
14	Early Mesolithic activities at ancient Lake Duvensee, northern Germany. <i>Holocene</i> , 2019, 29, 197-208.	1.7	18
15	Lack of support for adaptation of post-glacial horses to woodlands. <i>Nature Ecology and Evolution</i> , 2018, 2, 582-583.	7.8	9
16	Stone-age subsistence strategies at Lake Burtnieks, Latvia. <i>Journal of Archaeological Science: Reports</i> , 2018, 17, 992-1006.	0.5	14
17	Stable isotopic ratios from Mesolithic and Neolithic canids as an indicator of human economic and ritual activity. <i>Journal of Archaeological Science: Reports</i> , 2018, 17, 346-357.	0.5	6
18	How Fishy was the Inland Mesolithic? New Data from Friesack, Brandenburg, Germany. <i>Radiocarbon</i> , 2018, 60, 1621-1636.	1.8	11

#	ARTICLE	IF	CITATIONS
19	Neolithic fish remains from the freshwater shell midden RiĀĀĀukalns in northern Latvia. <i>Environmental Archaeology</i> , 2016, 21, 325-333.	1.2	18
20	Archaeogenetic evidence for medieval occurrence of Atlantic sturgeon <i>Acipenser oxyrinchus</i> in the North Sea. <i>Environmental Archaeology</i> , 2016, 21, 137-143.	1.2	1
21	Reconstruction of the historical distribution of sturgeons (Acipenseridae) in the eastern North Atlantic based on ancient <sc>DNA</sc> and bone morphology of archaeological remains: implications for conservation and restoration programmes. <i>Diversity and Distributions</i> , 2016, 22, 1036-1044.	4.1	11
22	Dietary freshwater reservoir effects and the radiocarbon ages of prehistoric human bones from Zvejnieki, Latvia. <i>Journal of Archaeological Science: Reports</i> , 2016, 6, 678-689.	0.5	9
23	European Bison as a Refugee Species? Evidence from Isotopic Data on Early Holocene Bison and Other Large Herbivores in Northern Europe. <i>PLoS ONE</i> , 2015, 10, e0115090.	2.5	109
24	First archaeogenetic results verify the mid-Holocene occurrence of Dalmatian pelican <i>Pelecanus crispus</i> far out of present range. <i>Journal of Avian Biology</i> , 2015, 46, 344-351.	1.2	8
25	New research at RiĀĀĀukalns, a Neolithic freshwater shell midden in northern Latvia. <i>Antiquity</i> , 2014, 88, 715-732.	1.0	34
26	Depositional environment and climate changes during the late Pleistocene as recorded by the Netiesos section in southern Lithuania. <i>Quaternary International</i> , 2013, 292, 136-149.	1.5	22
27	Carbon and nitrogen isotope signals in eel bone collagen from Mesolithic and Neolithic sites in northern Europe. <i>Journal of Archaeological Science</i> , 2012, 39, 2003-2011.	2.4	31
28	Holocene survival of the wild horse in Europe: a matter of open landscape?. <i>Journal of Quaternary Science</i> , 2011, 26, 805-812.	2.1	54
29	Mesolithic Hunter-Fishers in a Changing World: , 2011, , 21-37.		36
30	A new method in palaeoecology: fish community structure indicates environmental changes. <i>International Journal of Earth Sciences</i> , 2010, 99, 1763-1772.	1.8	12
31	Quaternary history of the European roe deer <i>Capreolus capreolus</i>. <i>Mammal Review</i> , 2009, 39, 1-16.	4.8	65
32	Post-Glacial immigration of the harbour porpoise (<i>Phocoena phocoena</i>) into the Baltic Sea<sup>*</sup>. <i>Boreas</i> , 2008, 37, 458-464.	2.4	14
33	Holocene environmental changes and the seal (Phocidae) fauna of the Baltic Sea: coming, going and staying. <i>Mammal Review</i> , 2008, 38, 231-246.	4.8	15
34	Ancient DNA provides no evidence for independent domestication of cattle in Mesolithic Rosenhof, Northern Germany. <i>Journal of Archaeological Science</i> , 2008, 35, 1257-1264.	2.4	44
35	Mitochondrial DNA analysis shows a Near Eastern Neolithic origin for domestic cattle and no indication of domestication of European aurochs. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2007, 274, 1377-1385.	2.6	209
36	Changes of sea level, landscape and culture: A review of the south-western Baltic area between 8800 and 4000BC. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2006, 240, 423-438.	2.3	45

#	ARTICLE	IF	CITATIONS
37	Holocene distribution and extinction of the moose ( <i>Alces alces</i> , Cervidae) in Central Europe. <i>Mammalian Biology</i> , 2005, 70, 329-344.	1.5	37
38	Neolithic fish remains from the freshwater shell midden Riņķuakalns in northern Latvia. <i>Environmental Archaeology</i> , 0, , 1-14.	1.2	1