

Oliver Ktting

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6691440/oliver-kotting-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19
papers

1,420
citations

17
h-index

19
g-index

19
ext. papers

1,616
ext. citations

7.8
avg, IF

3.9
L-index

#	Paper	IF	Citations
19	LIKE SEX4 1 Acts as a β -Amylase-Binding Scaffold on Starch Granules during Starch Degradation. <i>Plant Cell</i> , 2019 , 31, 2169-2186	11.6	17
18	Plastidial NAD-Dependent Malate Dehydrogenase: A Moonlighting Protein Involved in Early Chloroplast Development through Its Interaction with an FtsH12-FtsHi Protease Complex. <i>Plant Cell</i> , 2018 , 30, 1745-1769	11.6	31
17	Mechanistic Insights into Glucan Phosphatase Activity against Polyglucan Substrates. <i>Journal of Biological Chemistry</i> , 2015 , 290, 23361-70	5.4	22
16	Phosphoglucan-bound structure of starch phosphatase Starch Excess4 reveals the mechanism for C6 specificity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 7272-7	11.5	50
15	Phosphoglucan phosphatase function sheds light on starch degradation. <i>Trends in Plant Science</i> , 2014 , 19, 471-8	13.1	53
14	Plastidial NAD-dependent malate dehydrogenase is critical for embryo development and heterotrophic metabolism in Arabidopsis. <i>Plant Physiology</i> , 2014 , 164, 1175-90	6.6	59
13	Structure of the Arabidopsis glucan phosphatase like sex four2 reveals a unique mechanism for starch dephosphorylation. <i>Plant Cell</i> , 2013 , 25, 2302-14	11.6	31
12	Comprehensive survey of redox sensitive starch metabolising enzymes in Arabidopsis thaliana. <i>Plant Physiology and Biochemistry</i> , 2012 , 58, 89-97	5.4	56
11	Mutagenesis of cysteine 81 prevents dimerization of the APS1 subunit of ADP-glucose pyrophosphorylase and alters diurnal starch turnover in Arabidopsis thaliana leaves. <i>Plant Journal</i> , 2012 , 70, 231-42	6.9	72
10	Analysis of starch metabolism in chloroplasts. <i>Methods in Molecular Biology</i> , 2011 , 775, 387-410	1.4	41
9	The phosphoglucan phosphatase like sex Four2 dephosphorylates starch at the C3-position in Arabidopsis. <i>Plant Cell</i> , 2011 , 23, 4096-111	11.6	95
8	The Laforin-like dual-specificity phosphatase SEX4 from Arabidopsis hydrolyzes both C6- and C3-phosphate esters introduced by starch-related dikinases and thereby affects phase transition of alpha-glucans. <i>Plant Physiology</i> , 2010 , 152, 711-22	6.6	65
7	A putative phosphatase, LSF1, is required for normal starch turnover in Arabidopsis leaves. <i>Plant Physiology</i> , 2010 , 152, 685-97	6.6	84
6	Regulation of starch metabolism: the age of enlightenment?. <i>Current Opinion in Plant Biology</i> , 2010 , 13, 321-9	9.9	149
5	STARCH-EXCESS4 is a laforin-like Phosphoglucan phosphatase required for starch degradation in Arabidopsis thaliana. <i>Plant Cell</i> , 2009 , 21, 334-46	11.6	180
4	Starch breakdown: recent discoveries suggest distinct pathways and novel mechanisms. <i>Functional Plant Biology</i> , 2007 , 34, 465-473	2.7	63
3	Phosphorylation of C6- and C3-positions of glucosyl residues in starch is catalysed by distinct dikinases. <i>FEBS Letters</i> , 2006 , 580, 4872-6	3.8	137

- 2 Identification of a novel enzyme required for starch metabolism in Arabidopsis leaves. The phosphoglucan, water dikinase. *Plant Physiology*, **2005**, 137, 242-52 6.6 209
- 1 Starch Biosynthesis and Degradation in Plants1-10 6