

Juan H Ayala DÃ-az

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6693709/publications.pdf>

Version: 2024-02-01

81
papers

2,440
citations

172386

29
h-index

223716

46
g-index

84
all docs

84
docs citations

84
times ranked

2290
citing authors

#	ARTICLE	IF	CITATIONS
1	A green miniaturized aqueous biphasic system prepared with cholinium chloride and a phosphate salt to extract and preconcentrate personal care products in wastewater samples. <i>Journal of Chromatography A</i> , 2021, 1648, 462219.	1.8	3
2	Headspace solid-phase microextraction based on the metal-organic framework CIM-80(Al) coating to determine volatile methylsiloxanes and musk fragrances in water samples using gas chromatography and mass spectrometry. <i>Talanta</i> , 2021, 232, 122440.	2.9	21
3	Insights into Paraben Adsorption by Metal-Organic Frameworks for Analytical Applications. <i>ACS Applied Materials & Interfaces</i> , 2021, 13, 45639-45650.	4.0	9
4	Evaluation of Structurally Different Ionic Liquid-Based Surfactants in a Green Microwave-Assisted Extraction for the Flavonoids Profile Determination of <i>Mangifera</i> sp. and <i>Passiflora</i> sp. Leaves from Canary Islands. <i>Molecules</i> , 2020, 25, 4734.	1.7	12
5	Green solid-phase microextraction fiber coating based on the metal-organic framework CIM-80(Al): Analytical performance evaluation in direct immersion and headspace using gas chromatography and mass spectrometry for the analysis of water, urine and brewed coffee. <i>Analytica Chimica Acta</i> , 2020, 1133, 137-149.	2.6	30
6	Evolution and current advances in sorbent-based microextraction configurations. <i>Journal of Chromatography A</i> , 2020, 1634, 461670.	1.8	44
7	Solid-phase microextraction coatings based on the metal-organic framework ZIF-8: Ensuring stable and reusable fibers. <i>Talanta</i> , 2020, 215, 120910.	2.9	36
8	Use of a pH-sensitive polymer in a microextraction and preconcentration method directly combined with high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2020, 1619, 460910.	1.8	10
9	Mixed Functionalization of Organic Ligands in UiO-66: A Tool to Design Metal-Organic Frameworks for Tailored Microextraction. <i>Molecules</i> , 2019, 24, 3656.	1.7	15
10	Ionic liquid-based miniaturized aqueous biphasic system to develop an environmental-friendly analytical preconcentration method. <i>Talanta</i> , 2019, 203, 305-313.	2.9	13
11	Application of a Pillared-Layer Zn-Triazolate Metal-Organic Framework in the Dispersive Miniaturized Solid-Phase Extraction of Personal Care Products from Wastewater Samples. <i>Molecules</i> , 2019, 24, 690.	1.7	20
12	A guanidinium ionic liquid-based surfactant as an adequate solvent to separate and preconcentrate cadmium and copper in water using <i>in situ</i> dispersive liquid-liquid microextraction. <i>Analytical Methods</i> , 2018, 10, 1529-1537.	1.3	11
13	Salt-induced ionic liquid-based microextraction using a low cytotoxic guanidinium ionic liquid and liquid chromatography with fluorescence detection to determine monohydroxylated polycyclic aromatic hydrocarbons in urine. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 4701-4713.	1.9	25
14	Guanidinium ionic liquid-based surfactants as low cytotoxic extractants: Analytical performance in an <i>in-situ</i> dispersive liquid-liquid microextraction method for determining personal care products. <i>Journal of Chromatography A</i> , 2018, 1559, 102-111.	1.8	31
15	Insights in the analytical performance of neat metal-organic frameworks in the determination of pollutants of different nature from waters using dispersive miniaturized solid-phase extraction and liquid chromatography. <i>Talanta</i> , 2018, 179, 775-783.	2.9	52
16	Influence of Ligand Functionalization of UiO-66-Based Metal-Organic Frameworks When Used as Sorbents in Dispersive Solid-Phase Analytical Microextraction for Different Aqueous Organic Pollutants. <i>Molecules</i> , 2018, 23, 2869.	1.7	40
17	A green metal-organic framework to monitor water contaminants. <i>RSC Advances</i> , 2018, 8, 31304-31310.	1.7	34
18	Vacuum-assisted headspace-solid phase microextraction for determining volatile free fatty acids and phenols. Investigations on the effect of pressure on competitive adsorption phenomena in a multicomponent system. <i>Analytica Chimica Acta</i> , 2017, 962, 41-51.	2.6	53

#	ARTICLE	IF	CITATIONS
19	Monitoring trihalomethanes and nitrogenous disinfection by-products in blending desalinated waters using solid-phase microextraction and gas chromatography. <i>Environmental Technology</i> (United Kingdom), 2017, 38, 911-922.	1.2	11
20	Effect of the inclusion of banana silage in the diet of goats on physicochemical and sensory characteristics of cheeses at different ripening times. <i>Small Ruminant Research</i> , 2017, 149, 52-61.	0.6	4
21	Influence of vegetable coagulant and ripening time on the lipolytic and sensory profile of cheeses made with raw goat milk from Canary breeds. <i>Food Science and Technology International</i> , 2017, 23, 254-264.	1.1	11
22	Monitoring trihalomethanes in chlorinated waters using a dispersive liquid-liquid microextraction method with a non-chlorinated organic solvent and gas chromatography-mass spectrometry. <i>Environmental Technology</i> (United Kingdom), 2017, 38, 718-729.	1.2	5
23	Ionic Liquid-based Surfactants: A Step Forward. <i>RSC Smart Materials</i> , 2017, , 53-78.	0.1	8
24	Utilization of highly robust and selective crosslinked polymeric ionic liquid-based sorbent coatings in direct-immersion solid-phase microextraction and high-performance liquid chromatography for determining polar organic pollutants in waters. <i>Talanta</i> , 2016, 158, 125-133.	2.9	60
25	Magnetic ionic liquids as non-conventional extraction solvents for the determination of polycyclic aromatic hydrocarbons. <i>Analytica Chimica Acta</i> , 2016, 934, 106-113.	2.6	64
26	A magnetic-based dispersive micro-solid-phase extraction method using the metal-organic framework HKUST-1 and ultra-high-performance liquid chromatography with fluorescence detection for determining polycyclic aromatic hydrocarbons in waters and fruit tea infusions. <i>Journal of Chromatography A</i> , 2016, 1436, 42-50.	1.8	100
27	The metal-organic framework HKUST-1 as efficient sorbent in a vortex-assisted dispersive micro solid-phase extraction of parabens from environmental waters, cosmetic creams, and human urine. <i>Talanta</i> , 2015, 139, 13-20.	2.9	144
28	A simplified vortex-assisted emulsification microextraction method for determining personal care products in environmental water samples by ultra-high-performance liquid chromatography. <i>Analytical Methods</i> , 2015, 7, 1825-1833.	1.3	12
29	Double salts of ionic-liquid-based surfactants in microextraction: application of their mixed hemimicelles as novel sorbents in magnetic-assisted micro-dispersive solid-phase extraction for the determination of phenols. <i>Analytical and Bioanalytical Chemistry</i> , 2015, 407, 8753-8764.	1.9	26
30	Vortex-assisted emulsification microextraction followed by in-syringe ultrasound-assisted back-microextraction to determine haloacetic acids in waters. <i>Analytical Methods</i> , 2014, 6, 4115-4123.	1.3	9
31	Multiple headspace solid-phase microextraction for quantifying volatile free fatty acids in cheeses. <i>Talanta</i> , 2014, 129, 183-190.	2.9	19
32	Utilization of an ionic liquid <i>in situ</i> preconcentration method for the determination of the 15 + 1 European Union polycyclic aromatic hydrocarbons in drinking water and fruit tea infusions. <i>Journal of Separation Science</i> , 2013, 36, 2496-2506.	1.3	13
33	An in-situ extraction-preconcentration method using ionic liquid-based surfactants for the determination of organic contaminants contained in marine sediments. <i>Talanta</i> , 2012, 99, 972-983.	2.9	57
34	A novel preconcentration strategy for extraction methods based on common cationic surfactants: An alternative to classical coacervative extraction. <i>Journal of Chromatography A</i> , 2012, 1257, 9-18.	1.8	18
35	Headspace-single drop microextraction (HS-SDME) in combination with high-performance liquid chromatography (HPLC) to evaluate the content of alkyl- and methoxy-phenolic compounds in biomass smoke. <i>Talanta</i> , 2011, 85, 1265-1273.	2.9	26
36	In-situ ionic liquid-dispersive liquid-liquid microextraction method to determine endocrine disrupting phenols in seawaters and industrial effluents. <i>Mikrochimica Acta</i> , 2011, 174, 213-222.	2.5	59

#	ARTICLE	IF	CITATIONS
37	Suitability of ionic liquids as mobile phase additives in HPLC with fluorescence and UV detection for the determination of heterocyclic aromatic amines. <i>Journal of Separation Science</i> , 2010, 33, 182-190.	1.3	22
38	Ionic liquids as desorption solvents and memory effect suppressors in heterocyclic aromatic amines determination by SPME-HPLC fluorescence. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 394, 937-946.	1.9	49
39	Ionic liquids as mobile phase additives in high-performance liquid chromatography with electrochemical detection: Application to the determination of heterocyclic aromatic amines in meat-based infant foods. <i>Talanta</i> , 2009, 79, 590-597.	2.9	67
40	Determination of carbonyl compounds in smoke samples: strategies for sampling and standardization. <i>Journal of Environmental Monitoring</i> , 2009, 11, 1043.	2.1	1
41	Analytical methods applied to the determination of heterocyclic aromatic amines in foods. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 862, 15-42.	1.2	92
42	Micelle-mediated extractions using nonionic surfactant mixtures and HPLC-UV to determine endocrine-disrupting phenols in seawaters. <i>Analytical and Bioanalytical Chemistry</i> , 2008, 391, 735-744.	1.9	27
43	The ionic liquid 1-hexadecyl-3-methylimidazolium bromide as novel extracting system for polycyclic aromatic hydrocarbons contained in sediments using focused microwave-assisted extraction. <i>Journal of Chromatography A</i> , 2008, 1182, 145-152.	1.8	87
44	Estimation of Uncertainty in the Analysis of Carbonyl Compounds by HPLC-UV Using DNPH Derivatization. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 31, 361-381.	0.5	8
45	Evaluation of the Uncertainty Associated to the Determination of Heavy Metals in Seawater Using Graphite Furnace Atomic Absorption Spectrometry. <i>Analytical Letters</i> , 2007, 40, 3322-3342.	1.0	4
46	Monitoring chlorophenols in industrial effluents by solid-phase microextraction-gas chromatography-mass spectrometry. <i>International Journal of Environmental Analytical Chemistry</i> , 2007, 87, 159-175.	1.8	12
47	Focused Microwave-Assisted Extraction and HPLC with Electrochemical Detection to Determine Heterocyclic Amines in Meat Extracts. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2007, 30, 27-42.	0.5	13
48	Determination of the alkyl- and methoxy-phenolic content in wood extractives by micellar solid-phase microextraction and gas chromatography-mass spectrometry. <i>Talanta</i> , 2007, 73, 505-513.	2.9	15
49	Focused microwave-assisted micellar extraction combined with solid-phase microextraction-gas chromatography/mass spectrometry to determine chlorophenols in wood samples. <i>Analytica Chimica Acta</i> , 2007, 582, 10-18.	2.6	24
50	Determination of less polar heterocyclic amines in meat extracts. <i>Analytica Chimica Acta</i> , 2007, 582, 259-266.	2.6	43
51	Micellar solid-phase microextraction for determining partition coefficients of substituted polycyclic aromatic hydrocarbons in micellar media: possible prediction of hydrocarbon-micelle behaviour. <i>Analytical and Bioanalytical Chemistry</i> , 2007, 387, 2271-2281.	1.9	5
52	Experimental Design Optimization of Solid-Phase Microextraction Conditions for the Determination of Heterocyclic Aromatic Amines by High-Performance Liquid Chromatography. <i>Analytical Letters</i> , 2006, 39, 405-423.	1.0	19
53	Correlations Between Phenols-Micelles Partition Coefficients and Several Molecular Descriptors. An Approach to Predict the Phenols Behaviour in MSPME. <i>Chromatographia</i> , 2006, 63, 167-174.	0.7	8
54	Optimization of an analytical methodology for the determination of alkyl- and methoxy-phenolic compounds by HS-SPME in biomass smoke. <i>Analytical and Bioanalytical Chemistry</i> , 2006, 385, 1162-1171.	1.9	17

#	ARTICLE	IF	CITATIONS
55	Biosynthesis of Antitumoral and Bactericidal Sanguinarine. <i>Journal of Biomedicine and Biotechnology</i> , 2006, 2006, 1-6.	3.0	9
56	Study of the interactions between phenolic compounds and micellar media using micellar solid-phase microextraction/gas chromatography. <i>Journal of Chromatography A</i> , 2005, 1099, 64-74.	1.8	18
57	Emissions of polycyclic aromatic hydrocarbons from combustion of agricultural and sylvicultural debris. <i>Atmospheric Environment</i> , 2005, 39, 6654-6663.	1.9	48
58	Polycyclic Aromatic Hydrocarbons in Smoke Used to Smoke Cheese Produced by the Combustion of Rock Rose (<i>Cistus monspeliensis</i>) and Tree Heather (<i>Erica arborea</i>) Wood. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 176-182.	2.4	35
59	Coupling micelle-mediated extraction using mixtures of surfactants and fluorescence measurements with a fiber-optic for the screening of PAHs in seawater. <i>Analyst</i> , The, 2005, 130, 571-577.	1.7	13
60	Solid-phase microextraction coupled with high-performance liquid chromatography for the analysis of heterocyclic aromatic amines. <i>Journal of Chromatography A</i> , 2004, 1030, 87-93.	1.8	52
61	Optimization of a sampling method to determine polycyclic aromatic hydrocarbons in smoke from incomplete biomass combustion. <i>Analytica Chimica Acta</i> , 2004, 524, 287-294.	2.6	15
62	Nonionic surfactant mixtures: a new cloud-point extraction approach for the determination of PAHs in seawater using HPLC with fluorimetric detection. <i>Analytica Chimica Acta</i> , 2004, 518, 165-172.	2.6	105
63	Solid-Phase Microextraction Coupled to Gas Chromatography/Mass Spectrometry for Determining Polycyclic Aromatic Hydrocarbon ^â Micelle Partition Coefficients. <i>Analytical Chemistry</i> , 2004, 76, 4572-4578.	3.2	31
64	Micellar microwave-assisted extraction combined with solid-phase microextraction for the determination of polycyclic aromatic hydrocarbons in a certified marine sediment. <i>Analytica Chimica Acta</i> , 2003, 477, 81-91.	2.6	75
65	DETERMINATION OF N-NITROSODIMETHYLAMINE BY HPLC, WITH FLUORESCENCE DETECTION. A SURVEY OF N-NITROSODIMETHYLAMINE IN COMMERCIAL BEERS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2002, 25, 977-984.	0.5	9
66	Fast microwave-assisted dansylation of N-nitrosamines. <i>Journal of Chromatography A</i> , 2002, 946, 133-140.	1.8	42
67	Determination of polycyclic aromatic hydrocarbons in seawater by high-performance liquid chromatography with fluorescence detection following micelle-mediated preconcentration. <i>Journal of Chromatography A</i> , 2002, 949, 291-299.	1.8	71
68	Ultrasonic micellar extraction of polycyclic aromatic hydrocarbons from marine sediments. <i>Talanta</i> , 2001, 54, 15-23.	2.9	36
69	Cloud-point preconcentration and HPLC determination of polycyclic aromatic hydrocarbons in marine sediments. <i>Fresenius' Journal of Analytical Chemistry</i> , 2001, 371, 526-531.	1.5	29
70	Micellar Extraction of Polycyclic Aromatic Hydrocarbons from Certified Marine Sediment. <i>International Journal of Environmental Analytical Chemistry</i> , 2001, 81, 281-294.	1.8	17
71	Determination of polycyclic aromatic hydrocarbons in marine sediments by high-performance liquid chromatography after microwave-assisted extraction with micellar media. <i>Journal of Chromatography A</i> , 2000, 869, 515-522.	1.8	93
72	NON-LINEAR CALIBRATION IN QUANTITATIVE ANALYSIS BY HPTLC UTILIZING A FIBRE OPTIC FLUORESCENCE DETECTOR. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2000, 23, 2653-2668.	0.5	4

#	ARTICLE	IF	CITATIONS
73	Rapid microwave-assisted dansylation of biogenic amines. <i>Journal of Chromatography A</i> , 1998, 808, 87-93.	1.8	26
74	Selective Analysis of Fluorene by Quenched Fluorescence in Cetylpyridinium Bromide Micelles. <i>Microchemical Journal</i> , 1998, 60, 101-109.	2.3	13
75	Effect of non-ionic surfactants as mobile phase additives on the fluorescence intensity of dansyl derivatives of biogenic amines in high-performance thin-layer chromatography. <i>Analyst</i> , The, 1998, 123, 725-729.	1.7	17
76	Fluorescence Quenching of Polycyclic Aromatic Hydrocarbons by Cetylpyridinium Bromide: Discrimination between Alternant and Nonalternant Hydrocarbons. <i>Applied Spectroscopy</i> , 1997, 51, 380-386.	1.2	17
77	Effects of cetylpyridinium bromide micelles on the spectrofluorimetric characteristics of polycyclic aromatic hydrocarbons. <i>Talanta</i> , 1997, 44, 257-267.	2.9	18
78	Selective determination of acenaphthene in mixtures of three-ring polycyclic aromatic hydrocarbons by fluorescence quenching in micellar medium of cetylpyridinium bromide. <i>Journal of Fluorescence</i> , 1997, 7, 147-153.	1.3	5
79	Effects of cationic micelles on fluorescence of indole and indolecarboxylic acids. Analytical determinations. <i>Mikrochimica Acta</i> , 1995, 118, 153-162.	2.5	5
80	Degradation of carbaryl in natural waters: Enhanced hydrolysis rate in micellar solution. <i>Bulletin of Environmental Contamination and Toxicology</i> , 1992, 48, 171-8.	1.3	4
81	Spectrofluorimetric determination of carbaryl and 1-naphthol in micellar media. <i>Mikrochimica Acta</i> , 1991, 103, 171-179.	2.5	5