Mao Xu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/670107/publications.pdf

Version: 2024-02-01

1181555 1464605 18 198 7 14 citations h-index g-index papers 25 25 25 235 docs citations citing authors all docs times ranked

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | The effect of continuous intercostal nerve block vs. single shot on analgesic outcomes and hospital stays in minimally invasive direct coronary artery bypass surgery: a retrospective cohort study. BMC Anesthesiology, 2022, 22, 64. | 0.7 | 2 |
| 2 | Application of the new â€~XU-line' in the dynamic ultrasonographic evaluation of the spatial displacement of anterior cervical tissues for airway management: protocol for a prospective observational study. BMJ Open, 2022, 12, e058896. | 0.8 | 1 |
| 3 | Carbon dioxide: the cause of devastating stroke without hemodynamic compromise during laparoscopic nephrectomy with injury of the inferior vena cava. Medicine (United States), 2021, 100, e24892. | 0.4 | 3 |
| 4 | Preoperative X-ray C2C6AR is applicable for prediction of difficult laryngoscopy in patients with cervical spondylosis. BMC Anesthesiology, 2021, 21, 111. | 0.7 | 5 |
| 5 | Preemptive analgesia with a single low dose of intrathecal morphine in multilevel posterior lumbar interbody fusion surgery: a double-blind, randomized, controlled trial. Spine Journal, 2020, 20, 989-997. | 0.6 | 13 |
| 6 | A study of optimal concentration range and time window of sevoflurane preconditioning for brain protection in MCAO rats. BMC Anesthesiology, 2020, 20, 78. | 0.7 | 2 |
| 7 | Anterior neck soft tissue thickness for airway evaluation measured by MRI in patients with cervical spondylosis: prospective cohort study. BMJ Open, 2019, 9, e029987. | 0.8 | 5 |
| 8 | Reply to "Use of radiologic indicators to predict difficult laryngoscopy in patients with cervical spondylosis― Acta Anaesthesiologica Scandinavica, 2018, 62, 1033-1034. | 0.7 | 0 |
| 9 | Radiologic indicators for prediction of difficult laryngoscopy in patients with cervical spondylosis. Acta Anaesthesiologica Scandinavica, 2018, 62, 474-482. | 0.7 | 27 |
| 10 | P808Association between genetic variants in PCSK9/APOB/LDLR and premature myocardial infarction in Han Chinese. European Heart Journal, 2018, 39, . | 1.0 | 0 |
| 11 | Calcineurin/nuclear factor-l̂ºB signaling mediates isoflurane-induced hippocampal neuroinflammation and subsequent cognitive impairment in aged rats. Molecular Medicine Reports, 2017, 15, 201-209. | 1.1 | 45 |
| 12 | A cross-sectional study in a tertiary care hospital in China: noise or silence in the operating room. BMJ Open, 2017, 7, e016316. | 0.8 | 15 |
| 13 | Neck circumference to inter-incisor gap ratio: a new predictor of difficult laryngoscopy in cervical spondylosis patients. BMC Anesthesiology, 2017, 17, 55. | 0.7 | 17 |
| 14 | Shikani Optical Stylet versus Macintosh Laryngoscope for Intubation in Patients Undergoing Surgery for Cervical Spondylosis. Chinese Medical Journal, 2017, 130, 297-302. | 0.9 | 17 |
| 15 | Reply to "Comparing Shikani Optical Stylet and Macintosh Laryngoscope for Orotracheal Intubation― Chinese Medical Journal, 2017, 130, 1630. | 0.9 | 1 |
| 16 | Massive Pulmonary Embolism in Recovery Period of General Anesthesia: Rapid Diagnosis and Successful Rescue by the Guidance of Transthoracic Echocardiography. Chinese Medical Journal, 2017, 130, 245-246. | 0.9 | 1 |
| 17 | Regional anesthesia and lipid resuscitation for local anesthetic systemic toxicity in China: results of a survey by the orthopedic anesthesia group of the chinese society of anesthesiology. BMC Anesthesiology, 2015, 16 , 1 . | 0.7 | 35 |
| 18 | Application of a new combined model including radiological indicators to predict difficult airway in patients undergoing surgery for cervical spondylosis. Chinese Medical Journal, 2014, 127, 4043-8. | 0.9 | 6 |