

Theodore Pincus

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6701702/publications.pdf>

Version: 2024-02-01

202
papers

20,385
citations

26610

56
h-index

10152

140
g-index

206
all docs

206
docs citations

206
times ranked

14517
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | 2010 Rheumatoid arthritis classification criteria: An American College of Rheumatology/European League Against Rheumatism collaborative initiative. <i>Arthritis and Rheumatism</i> , 2010, 62, 2569-2581. | 6.7 | 6,781 |
| 2 | Assessment of patient satisfaction in activities of daily living using a modified stanford health assessment questionnaire. <i>Arthritis and Rheumatism</i> , 1983, 26, 1346-1353. | 6.7 | 1,223 |
| 3 | American College of Rheumatology/European League Against Rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. <i>Arthritis and Rheumatism</i> , 2011, 63, 573-586. | 6.7 | 864 |
| 4 | The american college of rheumatology 1991 revised criteria for the classification of global functional status in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1992, 35, 498-502. | 6.7 | 762 |
| 5 | Severe functional declines, work disability, and increased mortality in seventy-five rheumatoid arthritis patients studied over nine years. <i>Arthritis and Rheumatism</i> , 1984, 27, 864-872. | 6.7 | 657 |
| 6 | Measurement of Serum DNA-Binding Activity in Systemic Lupus Erythematosus. <i>New England Journal of Medicine</i> , 1969, 281, 701-705. | 13.9 | 468 |
| 7 | Toward a multidimensional health assessment questionnaire (MDHAQ): Assessment of advanced activities of daily living and psychological status in the patient-friendly health assessment questionnaire format. <i>Arthritis and Rheumatism</i> , 1999, 42, 2220-2230. | 6.7 | 335 |
| 8 | Increased expression of FcγRI/CD64 on circulating monocytes parallels ongoing inflammation and nephritis in lupus. <i>Arthritis Research and Therapy</i> , 2009, 11, R6. | 1.6 | 297 |
| 9 | Physical inactivity in patients with rheumatoid arthritis: Data from twenty-one countries in a cross-sectional, international study. <i>Arthritis and Rheumatism</i> , 2008, 59, 42-50. | 6.7 | 277 |
| 10 | RAPID3 (Routine Assessment of Patient Index Data 3), a Rheumatoid Arthritis Index Without Formal Joint Counts for Routine Care: Proposed Severity Categories Compared to Disease Activity Score and Clinical Disease Activity Index Categories. <i>Journal of Rheumatology</i> , 2008, 35, 2136-2147. | 1.0 | 267 |
| 11 | Prediction of Long-Term Mortality in Patients with Rheumatoid Arthritis according to Simple Questionnaire and Joint Count Measures. <i>Annals of Internal Medicine</i> , 1994, 120, 26. | 2.0 | 262 |
| 12 | A MAJOR GENETIC LOCUS AFFECTING RESISTANCE TO INFECTION WITH MURINE LEUKEMIA VIRUSES. <i>Journal of Experimental Medicine</i> , 1971, 133, 1234-1241. | 4.2 | 244 |
| 13 | Development and validation of the health assessment questionnaire II: A revised version of the health assessment questionnaire. <i>Arthritis and Rheumatism</i> , 2004, 50, 3296-3305. | 6.7 | 231 |
| 14 | Shouldn't standard rheumatology clinical care be evidence-based rather than eminence-based, eloquence-based, or elegance-based?. <i>Journal of Rheumatology</i> , 2007, 34, 1-4. | 1.0 | 229 |
| 15 | A simplified twenty-eight° joint quantitative articular index in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1989, 32, 531-537. | 6.7 | 214 |
| 16 | Social Conditions and Self-Management Are More Powerful Determinants of Health Than Access to Care. <i>Annals of Internal Medicine</i> , 1998, 129, 406. | 2.0 | 202 |
| 17 | Premature mortality in patients with rheumatoid arthritis: Evolving concepts. <i>Arthritis and Rheumatism</i> , 2001, 44, 1234-1236. | 6.7 | 201 |
| 18 | Self-Report Questionnaire Scores in Rheumatoid Arthritis Compared with Traditional Physical, Radiographic, and Laboratory Measures. <i>Annals of Internal Medicine</i> , 1989, 110, 259. | 2.0 | 194 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | QUEST-RA: quantitative clinical assessment of patients with rheumatoid arthritis seen in standard rheumatology care in 15 countries. <i>Annals of the Rheumatic Diseases</i> , 2007, 66, 1491-1496. | 0.5 | 172 |
| 20 | Patients seen for standard rheumatoid arthritis care have significantly better articular, radiographic, laboratory, and functional status in 2000 than in 1985. <i>Arthritis and Rheumatism</i> , 2005, 52, 1009-1019. | 6.7 | 170 |
| 21 | Formal education level as a significant marker of clinical status in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1988, 31, 1346-1357. | 6.7 | 164 |
| 22 | WHAT IS THE NATURAL HISTORY OF RHEUMATOID ARTHRITIS?. <i>Rheumatic Disease Clinics of North America</i> , 1993, 19, 123-151. | 0.8 | 163 |
| 23 | Eligibility of patients in routine care for major clinical trials of anti-tumor necrosis factor ? agents in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2003, 48, 313-318. | 6.7 | 156 |
| 24 | Remission and rheumatoid arthritis: Data on patients receiving usual care in twenty-four countries. <i>Arthritis and Rheumatism</i> , 2008, 58, 2642-2651. | 6.7 | 156 |
| 25 | Elevated MMPI scores for hypochondriasis, depression, and hysteria in patients with rheumatoid arthritis reflect disease rather than psychological status. <i>Arthritis and Rheumatism</i> , 1986, 29, 1456-1466. | 6.7 | 143 |
| 26 | Identification of work disability in rheumatoid arthritis: Physical, radiographic and laboratory variables do not add explanatory power to demographic and functional variables. <i>Journal of Clinical Epidemiology</i> , 1992, 45, 127-138. | 2.4 | 140 |
| 27 | Treatment of the anemia of rheumatoid arthritis with recombinant human erythropoietin: Clinical and in vitro studies. <i>Arthritis and Rheumatism</i> , 1989, 32, 638-642. | 6.7 | 131 |
| 28 | Listening to the patient: A practical guide to self-report questionnaires in clinical care. <i>Arthritis and Rheumatism</i> , 1999, 42, 1797-1808. | 6.7 | 125 |
| 29 | Measures of activity and damage in rheumatoid arthritis: Depiction of changes and prediction of mortality over five years. <i>Arthritis and Rheumatism</i> , 1997, 10, 381-394. | 6.7 | 116 |
| 30 | RAPID3, an Index to Assess and Monitor Patients with Rheumatoid Arthritis, Without Formal Joint Counts: Similar Results to DAS28 and CDAl in Clinical Trials and Clinical Care. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 773-778. | 0.8 | 116 |
| 31 | Self-report questionnaires in five rheumatic diseases comparisons of health status constructs and associations with formal education level. <i>Arthritis and Rheumatism</i> , 1989, 2, 122-131. | 6.7 | 111 |
| 32 | The importance of age, education, and comorbidity in the substantial earnings losses of individuals with symmetric polyarthritis. <i>Arthritis and Rheumatism</i> , 1988, 31, 348-357. | 6.7 | 110 |
| 33 | Immunologic studies of rheumatoid arthritis patients treated with methotrexate. <i>Arthritis and Rheumatism</i> , 1987, 30, 481-488. | 6.7 | 108 |
| 34 | Hla-dr alleles with naturally occurring amino acid substitutions and risk for development of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1990, 33, 939-946. | 6.7 | 108 |
| 35 | Quantitative pain assessment for routine care of rheumatoid arthritis patients, using a pain scale based on activities of daily living and a visual analog pain scale. <i>Arthritis and Rheumatism</i> , 1987, 30, 630-636. | 6.7 | 105 |
| 36 | Further development of a physical function scale on a MDHAQ [corrected] for standard care of patients with rheumatic diseases. <i>Journal of Rheumatology</i> , 2005, 32, 1432-9. | 1.0 | 105 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Erythrocyte Sedimentation Rate, C-Reactive Protein, or Rheumatoid Factor Are Normal at Presentation in 35%–45% of Patients with Rheumatoid Arthritis Seen Between 1980 and 2004: Analyses from Finland and the United States. <i>Journal of Rheumatology</i> , 2009, 36, 1387-1390. | 1.0 | 104 |
| 38 | Reduced joint counts in controlled clinical trials in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1994, 37, 470-475. | 6.7 | 101 |
| 39 | Making an impact on mortality in rheumatoid arthritis: Targeting cardiovascular comorbidity. <i>Arthritis and Rheumatism</i> , 2004, 50, 1734-1739. | 6.7 | 95 |
| 40 | Formal education and five-year mortality in rheumatoid arthritis: Mediation by helplessness scale scores. <i>Arthritis and Rheumatism</i> , 1996, 9, 463-472. | 6.7 | 94 |
| 41 | RAPID3 (Routine Assessment of Patient Index Data) on an MDHAQ (Multidimensional Health Assessment) Tj ETQq1 1 0.784314 rgBT Research. 2010. 62. 181-189. | 1.5 | 93 |
| 42 | Most patients receiving routine care for rheumatoid arthritis in 2001 did not meet inclusion criteria for most recent clinical trials or american college of rheumatology criteria for remission. <i>Journal of Rheumatology</i> , 2003, 30, 1138-46. | 1.0 | 89 |
| 43 | A composite disease activity scale for clinical practice, observational studies, and clinical trials: the patient activity scale (PAS/PAS-II). <i>Journal of Rheumatology</i> , 2005, 32, 2410-5. | 1.0 | 88 |
| 44 | Radiographic and joint count findings of the hand in rheumatoid arthritis. related and unrelated findings. <i>Arthritis and Rheumatism</i> , 1988, 31, 44-51. | 6.7 | 85 |
| 45 | Approaches for identifying and defining environmentally associated rheumatic disorders. <i>Arthritis and Rheumatism</i> , 2000, 43, 243. | 6.7 | 82 |
| 46 | Learned helplessness in systemic lupus erythematosus: analysis using the rheumatology attitudes index. <i>Arthritis and Rheumatism</i> , 1990, 33, 281-286. | 6.7 | 81 |
| 47 | Urgent care and tight control of rheumatoid arthritis as in diabetes and hypertension: Better treatments but a shortage of rheumatologists. <i>Arthritis and Rheumatism</i> , 2002, 46, 851-854. | 6.7 | 80 |
| 48 | The beck depression inventory, center for epidemiological studies depression scale (CES-D), and general well-being schedule depression subscale in rheumatoid arthritis criterion contamination of responses. <i>Arthritis and Rheumatism</i> , 1991, 4, 3-11. | 6.7 | 74 |
| 49 | The sense of coherence scale in patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1995, 8, 28-35. | 6.7 | 73 |
| 50 | Mortality in the rheumatic diseases. <i>Arthritis and Rheumatism</i> , 1995, 8, 229-241. | 6.7 | 71 |
| 51 | Quantitative measures for assessing rheumatoid arthritis in clinical trials and clinical care. <i>Best Practice and Research in Clinical Rheumatology</i> , 2003, 17, 753-781. | 1.4 | 69 |
| 52 | RAPID3 (Routine Assessment of Patient Index Data 3) severity categories and response criteria: Similar results to DAS28 (Disease Activity Score) and CDAI (Clinical Disease Activity Index) in the RAPID 1 (Rheumatoid Arthritis Prevention of Structural Damage) clinical trial of certolizumab pegol. <i>Arthritis Care and Research</i> , 2011, 63, 1142-1149. | 1.5 | 67 |
| 53 | Assessing dyspnea and its impact on patients with connective tissue disease-related interstitial lung disease. <i>Respiratory Medicine</i> , 2010, 104, 1350-1355. | 1.3 | 60 |
| 54 | The Disease Activity Score is not suitable as the sole criterion for initiation and evaluation of anti-tumor necrosis factor therapy in the clinic: Discordance between assessment measures and limitations in questionnaire use for regulatory purposes. <i>Arthritis and Rheumatism</i> , 2005, 52, 3873-3879. | 6.7 | 58 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Morning stiffness in patients with early rheumatoid arthritis is associated more strongly with functional disability than with joint swelling and erythrocyte sedimentation rate. <i>Journal of Rheumatology</i> , 2004, 31, 1723-6. | 1.0 | 57 |
| 56 | Visual analog scales in formats other than a 10 centimeter horizontal line to assess pain and other clinical data. <i>Journal of Rheumatology</i> , 2008, 35, 1550-8. | 1.0 | 57 |
| 57 | A clue from a self-report questionnaire to distinguish rheumatoid arthritis from noninflammatory diffuse musculoskeletal pain. <i>Arthritis and Rheumatism</i> , 1990, 33, 1317-1322. | 6.7 | 56 |
| 58 | A proposed continuous quality improvement approach to assessment and management of patients with rheumatoid arthritis without formal joint counts, based on quantitative routine assessment of patient index data (RAPID) scores on a multidimensional health assessment questionnaire (MDHAQ). <i>Best Practice and Research in Clinical Rheumatology</i> , 2007, 21, 789-804. | 1.4 | 56 |
| 59 | Discordance of Global Assessments by Patient and Physician Is Higher in Female than in Male Patients Regardless of the Physician's Sex: Data on Patients with Rheumatoid Arthritis, Axial Spondyloarthritis, and Psoriatic Arthritis from the DANBIO Registry. <i>Journal of Rheumatology</i> , 2015, 42, 1781-1785. | 1.0 | 56 |
| 60 | The Underestimated Long Term Medical and Economic Consequences of Rheumatoid Arthritis. <i>Drugs</i> , 1995, 50, 1-14. | 4.9 | 55 |
| 61 | Laboratory Tests to Assess Patients with Rheumatoid Arthritis: Advantages and Limitations. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 731-734. | 0.8 | 55 |
| 62 | The assessment of rheumatoid arthritis and the acceptability of self-report questionnaires in clinical practice. <i>Arthritis and Rheumatism</i> , 2003, 49, 59-63. | 6.7 | 49 |
| 63 | Patient questionnaires and formal education level as prospective predictors of mortality over 10 years in 97% of 1416 patients with rheumatoid arthritis from 15 United States private practices. <i>Journal of Rheumatology</i> , 2004, 31, 229-34. | 1.0 | 49 |
| 64 | Time to score quantitative rheumatoid arthritis measures: 28-Joint Count, Disease Activity Score, Health Assessment Questionnaire (HAQ), Multidimensional HAQ (MDHAQ), and Routine Assessment of Patient Index Data (RAPID) scores. <i>Journal of Rheumatology</i> , 2008, 35, 603-9. | 1.0 | 48 |
| 65 | MDHAQ/RAPID3 to Recognize Improvement Over 2 Months in Usual Care of Patients With Osteoarthritis, Systemic Lupus Erythematosus, Spondyloarthropathy, and Gout, as Well as Rheumatoid Arthritis. <i>Journal of Clinical Rheumatology</i> , 2013, 19, 169-174. | 0.5 | 45 |
| 66 | Depression scales in rheumatoid arthritis: criterion contamination in interpretation of patient responses. <i>Patient Education and Counseling</i> , 1993, 20, 133-143. | 1.0 | 43 |
| 67 | Treatment of rheumatoid arthritis: a global perspective on the use of antirheumatic drugs. <i>Modern Rheumatology</i> , 2008, 18, 228-239. | 0.9 | 43 |
| 68 | ACR 20: Clinical or statistical significance?. <i>Arthritis and Rheumatism</i> , 1999, 42, 1572-1576. | 6.7 | 42 |
| 69 | Diagnosis of Fibromyalgia: Disagreement Between Fibromyalgia Criteria and Clinician-Based Fibromyalgia Diagnosis in a University Clinic. <i>Arthritis Care and Research</i> , 2019, 71, 343-351. | 1.5 | 41 |
| 70 | A Multi-Dimensional Health Assessment Questionnaire (MDHAQ) and Routine Assessment of Patient Index Data (RAPID3) Scores are Informative in Patients with All Rheumatic Diseases. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 819-827. | 0.8 | 40 |
| 71 | Continuous indices of core data set measures in rheumatoid arthritis clinical trials: Lower responses to placebo than seen with categorical responses with the American College of Rheumatology 20% criteria. <i>Arthritis and Rheumatism</i> , 2005, 52, 1031-1036. | 6.7 | 39 |
| 72 | Observer Variation in Quantitative Assessment of Rheumatoid Arthritis Part II. A Simplified Scoring System. <i>Investigative Radiology</i> , 1987, 22, 41-46. | 3.5 | 38 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | An index of patient reported outcomes (PRO-Index) discriminates effectively between active and control treatment in 4 clinical trials of adalimumab in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2006, 33, 2146-52. | 1.0 | 38 |
| 74 | A practical guide to scoring a Multi-Dimensional Health Assessment Questionnaire (MDHAQ) and Routine Assessment of Patient Index Data (RAPID) scores in 10–20 seconds for use in standard clinical care, without rulers, calculators, websites or computers. <i>Best Practice and Research in Clinical Rheumatology</i> , 2007, 21, 755-787. | 1.4 | 37 |
| 75 | Education, self-care, and outcomes of rheumatic diseases: Further challenges to the “biomedical model” paradigm. <i>Arthritis and Rheumatism</i> , 1997, 10, 283-288. | 6.7 | 36 |
| 76 | A Biopsychosocial Model to Complement a Biomedical Model: Patient Questionnaire Data and Socioeconomic Status Usually Are More Significant than Laboratory Tests and Imaging Studies in Prognosis of Rheumatoid Arthritis. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 699-712. | 0.8 | 36 |
| 77 | Rheumatoid Arthritis. <i>Annals of Internal Medicine</i> , 2010, 153, ITC1-1. | 2.0 | 36 |
| 78 | Implementation of multiple outpatient formularies: Undesirable effects*. <i>Clinical Pharmacology and Therapeutics</i> , 1997, 61, 1-7. | 2.3 | 35 |
| 79 | Quantitative measures of rheumatic diseases for clinical research versus standard clinical care: differences, advantages and limitations. <i>Best Practice and Research in Clinical Rheumatology</i> , 2007, 21, 601-628. | 1.4 | 35 |
| 80 | Decline of Mean Initial Prednisone Dosage From 10.3 to 3.6 mg/day to Treat Rheumatoid Arthritis Between 1980 and 2004 in One Clinical Setting, With Long-Term Effectiveness of Dosages Less Than 5 mg/day. <i>Arthritis Care and Research</i> , 2013, 65, 729-736. | 1.5 | 34 |
| 81 | Observer Variation in Quantitative Assessment of Rheumatoid Arthritis Part I. Scoring Erosions and Joint Space Narrowing. <i>Investigative Radiology</i> , 1986, 21, 922-927. | 3.5 | 33 |
| 82 | Clinical evidence for osteoarthritis as an inflammatory disease. <i>Current Rheumatology Reports</i> , 2001, 3, 524-534. | 2.1 | 33 |
| 83 | Quantitative Data for Care of Patients with Systemic Lupus Erythematosus in Usual Clinical Settings: A Patient Multidimensional Health Assessment Questionnaire and Physician Estimate of Noninflammatory Symptoms. <i>Journal of Rheumatology</i> , 2011, 38, 1309-1316. | 1.0 | 32 |
| 84 | Proposed Severity and Response Criteria for Routine Assessment of Patient Index Data (RAPID3): Results for Categories of Disease Activity and Response Criteria in Abatacept Clinical Trials. <i>Journal of Rheumatology</i> , 2011, 38, 2565-2571. | 1.0 | 31 |
| 85 | Comparison of the dapson recovery ratio and the erythromycin breath test as in vivo probes of CYP3A activity in patients with rheumatoid arthritis receiving cyclosporine*. <i>Clinical Pharmacology and Therapeutics</i> , 1996, 59, 47-51. | 2.3 | 30 |
| 86 | Screening for Low Literacy in a Rheumatology Setting. <i>Journal of Clinical Rheumatology</i> , 2010, 16, 359-364. | 0.5 | 30 |
| 87 | The importance of the patient history and physical examination in rheumatoid arthritis in contrast to other chronic diseases: Results of a physician survey. <i>Arthritis Care and Research</i> , 2012, 64, n/a-n/a. | 1.5 | 30 |
| 88 | Relative efficiencies of physician/assessor global estimates and patient questionnaire measures are similar to or greater than joint counts to distinguish adalimumab from control treatments in rheumatoid arthritis clinical trials. <i>Journal of Rheumatology</i> , 2008, 35, 201-5. | 1.0 | 30 |
| 89 | Are Long-Term Very Low Doses of Prednisone for Patients with Rheumatoid Arthritis as Helpful as High Doses Are Harmful?. <i>Annals of Internal Medicine</i> , 2002, 136, 76. | 2.0 | 29 |
| 90 | Value of the Routine Assessment of Patient Index Data 3 in Patients With Psoriatic Arthritis: Results From a Tight-Control Clinical Trial and an Observational Cohort. <i>Arthritis Care and Research</i> , 2018, 70, 1198-1205. | 1.5 | 29 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Disease Burden in Osteoarthritis Is Similar to That of Rheumatoid Arthritis at Initial Rheumatology Visit and Significantly Greater Six Months Later. <i>Arthritis and Rheumatology</i> , 2019, 71, 1276-1284. | 2.9 | 29 |
| 92 | Bony Ankylosis in Rheumatoid Arthritis. <i>Investigative Radiology</i> , 1987, 22, 303-309. | 3.5 | 28 |
| 93 | HLA-DPB1*0301 is a Major Risk Factor for Rheumatoid Factor-Negative Adult Rheumatoid Arthritis. <i>Arthritis and Rheumatism</i> , 1991, 34, 1310-1312. | 6.7 | 28 |
| 94 | Discordance of Global Estimates by Patients and Their Physicians in Usual Care of Many Rheumatic Diseases: Association With 5 Scores on a Multidimensional Health Assessment Questionnaire (MDHAQ) That Are Not Found on the Health Assessment Questionnaire (HAQ). <i>Arthritis Care and Research</i> , 2014, 66, 934-942. | 1.5 | 28 |
| 95 | RAPID3-an index of physical function, pain, and global status as "vital signs" to improve care for people with chronic rheumatic diseases. <i>Bulletin of the NYU Hospital for Joint Diseases</i> , 2009, 67, 211-25. | 0.7 | 28 |
| 96 | Can a Multi-Dimensional Health Assessment Questionnaire (MDHAQ) and Routine Assessment of Patient Index Data (RAPID) scores be informative in patients with all rheumatic diseases?. <i>Best Practice and Research in Clinical Rheumatology</i> , 2007, 21, 733-753. | 1.4 | 27 |
| 97 | Patient self-report RADAI (Rheumatoid Arthritis Disease Activity Index) joint counts on an MDHAQ (Multidimensional Health Assessment Questionnaire) in usual care of consecutive patients with rheumatic diseases other than rheumatoid arthritis. <i>Arthritis Care and Research</i> , 2013, 65, 288-293. | 1.5 | 27 |
| 98 | A PRAGMATIC APPROACH TO COST-EFFECTIVE USE OF LABORATORY TESTS AND IMAGING PROCEDURES IN PATIENTS WITH MUSCULOSKELETAL SYMPTOMS. <i>Primary Care - Clinics in Office Practice</i> , 1993, 20, 795-814. | 0.7 | 27 |
| 99 | A multidimensional health assessment questionnaire (MDHAQ) for all patients with rheumatic diseases to complete at all visits in standard clinical care. <i>Bulletin of the NYU Hospital for Joint Diseases</i> , 2007, 65, 150-60. | 0.7 | 27 |
| 100 | Limitations of a quantitative swollen and tender joint count to assess and monitor patients with rheumatoid arthritis. <i>Bulletin of the NYU Hospital for Joint Diseases</i> , 2008, 66, 216-23. | 0.7 | 27 |
| 101 | The case for early intervention in rheumatoidarthritis. <i>Journal of Autoimmunity</i> , 1992, 5, 209-226. | 3.0 | 26 |
| 102 | Documenting quality management in rheumatic disease: Are patient questionnaires the best (and only) method?. <i>Arthritis and Rheumatism</i> , 1996, 9, 339-348. | 6.7 | 26 |
| 103 | Prediction of Remission in a French Early Arthritis Cohort by RAPID3 and other Core Data Set Measures, but Not by the Absence of Rheumatoid Factor, Anticitrullinated Protein Antibodies, or Radiographic Erosions. <i>Journal of Rheumatology</i> , 2016, 43, 1285-1291. | 1.0 | 26 |
| 104 | Patient questionnaires for clinical research and improved standard patient care: is it better to have 80% of the information in 100% of patients or 100% of the information in 5% of patients?. <i>Journal of Rheumatology</i> , 2005, 32, 575-7. | 1.0 | 26 |
| 105 | Joint Counts to Assess Rheumatoid Arthritis for Clinical Research and Usual Clinical Care: Advantages and Limitations. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 713-722. | 0.8 | 25 |
| 106 | Can Remission in Rheumatoid Arthritis Be Assessed Without Laboratory Tests or a Formal Joint Count? Possible Remission Criteria Based on a Self-report RAPID3 Score and Careful Joint Examination in the ESPOIR Cohort. <i>Journal of Rheumatology</i> , 2013, 40, 386-393. | 1.0 | 25 |
| 107 | Clinical Trials Documenting the Efficacy of Low-Dose Glucocorticoids in Rheumatoid Arthritis. <i>NeuroImmunoModulation</i> , 2015, 22, 46-50. | 0.9 | 25 |
| 108 | Further evidence for the lack of association between acetylator phenotype and systemic lupus erythematosus. <i>Arthritis and Rheumatism</i> , 1986, 29, 508-514. | 6.7 | 24 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Evidence-based practice and practice-based evidence. <i>Nature Clinical Practice Rheumatology</i> , 2006, 2, 114-115. | 3.2 | 24 |
| 110 | Information technology in rheumatology. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, 1. | 0.4 | 24 |
| 111 | The HAQ Compared with the MDHAQ: "Keep It Simple, Stupid" (KISS), with Feasibility and Clinical Value as Primary Criteria for Patient Questionnaires in Usual Clinical Care. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 787-798. | 0.8 | 23 |
| 112 | Premature Mortality: A Neglected Outcome in Rheumatic Diseases?. <i>Arthritis Care and Research</i> , 2015, 67, 1043-1046. | 1.5 | 23 |
| 113 | WHY SHOULD RHEUMATOLOGISTS COLLECT PATIENT SELF-REPORT QUESTIONNAIRES IN ROUTINE RHEUMATOLOGIC CARE?. <i>Rheumatic Disease Clinics of North America</i> , 1995, 21, 271-319. | 0.8 | 23 |
| 114 | The DAS is the most specific measure, but a patient questionnaire is the most informative measure to assess rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2006, 33, 834-7. | 1.0 | 23 |
| 115 | Treat-to-target: not as simple as it appears. <i>Clinical and Experimental Rheumatology</i> , 2012, 30, S10-20. | 0.4 | 23 |
| 116 | Limitations of clinical trials in chronic diseases: is the efficacy of methotrexate (MTX) underestimated in polyarticular psoriatic arthritis on the basis of limitations of clinical trials more than on limitations of MTX, as was seen in rheumatoid arthritis?. <i>Clinical and Experimental Rheumatology</i> , 2015, 33, S82-93. | 0.4 | 23 |
| 117 | How useful are combinations of blood tests in "Rheumatic panels" in diagnosis of rheumatic diseases?. <i>Journal of General Internal Medicine</i> , 1988, 3, 435-442. | 1.3 | 22 |
| 118 | Patient Questionnaires in Rheumatoid Arthritis: Advantages and Limitations as "Quantitative, Standardized Scientific Medical History". <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 735-743. | 0.8 | 22 |
| 119 | Criterion Contamination of Depression Scales in Patients with Rheumatoid Arthritis: The Need for Interpretation of Patient Questionnaires (as All Clinical Measures) in the Context of All Information About the Patient. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 861-864. | 0.8 | 22 |
| 120 | Associations between clinical status questionnaire scores and formal education level in persons with systemic lupus erythematosus. <i>Arthritis and Rheumatism</i> , 1990, 33, 407-411. | 6.7 | 21 |
| 121 | Clinical trials in rheumatic diseases: designs and limitations. <i>Rheumatic Disease Clinics of North America</i> , 2004, 30, 701-724. | 0.8 | 20 |
| 122 | Declines in Erythrocyte Sedimentation Rates in Patients with Rheumatoid Arthritis Over the Second Half of the 20th Century. <i>Journal of Rheumatology</i> , 2009, 36, 1596-1599. | 1.0 | 20 |
| 123 | Poor physical function, pain and limited exercise: risk factors for premature mortality in the range of smoking or hypertension, identified on a simple patient self-report questionnaire for usual care. <i>BMJ Open</i> , 2011, 1, e000070-e000070. | 0.8 | 20 |
| 124 | Minimal Clinically Important Improvement of Routine Assessment of Patient Index Data 3 in Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2019, 46, 27-30. | 1.0 | 20 |
| 125 | In vitro rheumatoid factor synthesis in patients taking second-line drugs for rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1988, 31, 1090-1096. | 6.7 | 19 |
| 126 | Routine Assessment of Patient Index Data (RAPID3) and Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) Scores Yield Similar Information in 85 Korean Patients With Ankylosing Spondylitis Seen in Usual Clinical Care. <i>Journal of Clinical Rheumatology</i> , 2015, 21, 300-304. | 0.5 | 19 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | MDHAQ/RAPID3 scores in patients with osteoarthritis are similar to or higher than in patients with rheumatoid arthritis: a cross-sectional study from current routine rheumatology care at four sites. <i>RMD Open</i> , 2017, 3, e000391. | 1.8 | 18 |
| 128 | Use of self-report activities of daily living questionnaires in osteoarthritis. <i>Arthritis and Rheumatism</i> , 1988, 1, 23-32. | 6.7 | 17 |
| 129 | Complexities in Assessment of Rheumatoid Arthritis: Absence of a Single Gold Standard Measure. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 687-697. | 0.8 | 17 |
| 130 | How to Collect an MDHAQ to Provide Rheumatology Vital Signs (Function, Pain, Global Status, and) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 Regarding the MDHAQ. <i>Rheumatic Disease Clinics of North America</i> , 2009, 35, 799-812. | 0.8 | 17 |
| 131 | Circadian Use of Glucocorticoids in Rheumatoid Arthritis. <i>NeuroImmunoModulation</i> , 2015, 22, 33-39. | 0.9 | 17 |
| 132 | Fibromyalgia Assessment Screening Tools (FAST) Based on Only Multidimensional Health Assessment Questionnaire (MDHAQ) Scores as Clues to Fibromyalgia. <i>ACR Open Rheumatology</i> , 2019, 1, 516-525. | 0.9 | 16 |
| 133 | Are Patient Self-Report Questionnaires as "Scientific" as Biomarkers in "Treat-totarget" and Prognosis in Rheumatoid Arthritis?. <i>Current Pharmaceutical Design</i> , 2014, 21, 241-256. | 0.9 | 16 |
| 134 | Pain, function, and RAPID scores: vital signs in chronic diseases, analogous to pulse and temperature in acute diseases and blood pressure and cholesterol in long-term health. <i>Bulletin of the NYU Hospital for Joint Diseases</i> , 2008, 66, 155-65. | 0.7 | 15 |
| 135 | Association of scleroderma with a t cell antigen receptor β gene restriction fragment length polymorphism. <i>Arthritis and Rheumatism</i> , 1990, 33, 569-573. | 6.7 | 14 |
| 136 | Quantitative measures and indices to assess rheumatoid arthritis in clinical trials and clinical care. <i>Rheumatic Disease Clinics of North America</i> , 2004, 30, 725-751. | 0.8 | 14 |
| 137 | Responsiveness of a simple RAPID-3-like index compared to disease-specific BASDAI and ASDAS indices in patients with axial spondyloarthritis. <i>RMD Open</i> , 2016, 2, e000235. | 1.8 | 14 |
| 138 | Can RAPID3, an index without formal joint counts or laboratory tests, serve to guide rheumatologists in tight control of rheumatoid arthritis in usual clinical care?. <i>Bulletin of the NYU Hospital for Joint Diseases</i> , 2009, 67, 254-66. | 0.7 | 13 |
| 139 | GUERARD treat-to-target strategy is significantly more efficacious than ESPOIR routine care in early rheumatoid arthritis according to patient-reported outcomes and physician global estimate. <i>Rheumatology</i> , 2013, 52, 1890-1897. | 0.9 | 12 |
| 140 | The Past versus the Present, 1980-2004: Reduction of Mean Initial Low-Dose, Long-Term Glucocorticoid Therapy in Rheumatoid Arthritis from 10.3 to 3.6 mg/Day, Concomitant with Early Methotrexate, with Long-Term Effectiveness and Safety of Less than 5 mg/Day. <i>NeuroImmunoModulation</i> , 2015, 22, 89-103. | 0.9 | 12 |
| 141 | Fibromyalgia Assessment Screening Tool: Clues to Fibromyalgia on a Multidimensional Health Assessment Questionnaire for Routine Care. <i>Journal of Rheumatology</i> , 2020, 47, 761-769. | 1.0 | 12 |
| 142 | Patient preference in a crossover clinical trial of patients with osteoarthritis of the knee or hip: face validity of self-report questionnaire ratings. <i>Journal of Rheumatology</i> , 2005, 32, 533-9. | 1.0 | 11 |
| 143 | Pragmatic and scientific advantages of MDHAQ/ RAPID3 completion by all patients at all visits in routine clinical care. <i>Bulletin of the NYU Hospital for Joint Diseases</i> , 2012, 70 Suppl 1, 30-6. | 0.7 | 11 |
| 144 | Saving time and improving care with a multidimensional health assessment questionnaire: 10 practical considerations. <i>Journal of Rheumatology</i> , 2006, 33, 448-54. | 1.0 | 10 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Are patient questionnaire scores as "scientific" as laboratory tests for rheumatology clinical care? Bulletin of the NYU Hospital for Joint Diseases, 2010, 68, 130-9. | 0.7 | 10 |
| 146 | MDHAQ/RAPID3 scores: quantitative patient history data in a standardized "scientific" format for optimal assessment of patient status and quality of care in rheumatic diseases. Bulletin of the NYU Hospital for Joint Diseases, 2011, 69, 201-14. | 0.7 | 10 |
| 147 | MDHAQ/RAPID3 can provide a roadmap or agenda for all rheumatology visits when the entire MDHAQ is completed at all patient visits and reviewed by the doctor before the encounter. Bulletin of the NYU Hospital for Joint Diseases, 2012, 70, 177-86. | 0.7 | 10 |
| 148 | Flowsheets That Include MDHAQ Physical Function, Pain, Global, and RAPID3 Scores, Laboratory Tests, and Medications to Monitor Patients with all Rheumatic Diseases: An Electronic Database for an Electronic Medical Record. Rheumatic Disease Clinics of North America, 2009, 35, 829-842. | 0.8 | 9 |
| 149 | Multidimensional Health Assessment Questionnaire as an Effective Tool to Screen for Depression in Routine Rheumatology Care. Arthritis Care and Research, 2021, 73, 120-129. | 1.5 | 9 |
| 150 | Electronic multidimensional health assessment questionnaire (eMDHAQ): past, present and future of a proposed single data management system for clinical care, research, quality improvement, and monitoring of long-term outcomes. Clinical and Experimental Rheumatology, 2016, 34, S17-S33. | 0.4 | 9 |
| 151 | Reliability, Feasibility, and Patient Acceptance of an Electronic Version of a Multidimensional Health Assessment Questionnaire for Routine Rheumatology Care: Validation and Patient Preference Study. JMIR Formative Research, 2020, 4, e15815. | 0.7 | 8 |
| 152 | Advantages and limitations of quantitative measures to assess rheumatoid arthritis: joint counts, radiographs, laboratory tests, and patient questionnaires. Bulletin of the NYU Hospital for Joint Diseases, 2006, 64, 32-9. | 0.7 | 8 |
| 153 | Patient questionnaires and formal education as more significant prognostic markers than radiographs or laboratory tests for rheumatoid arthritis mortality—limitations of a biomedical model to predict long-term outcomes. Bulletin of the NYU Hospital for Joint Diseases, 2007, 65 Suppl 1, S29-36. | 0.7 | 7 |
| 154 | Assessment of fatigue in routine care on a Multidimensional Health Assessment Questionnaire (MDHAQ): a cross-sectional study of associations with RAPID3 and other variables in different rheumatic diseases. Clinical and Experimental Rheumatology, 2016, 34, 901-909. | 0.4 | 7 |
| 155 | Clues on the MDHAQ to Identify Patients with Fibromyalgia and Similar Chronic Pain Conditions. Rheumatic Disease Clinics of North America, 2009, 35, 865-869. | 0.8 | 6 |
| 156 | Complex Measures and Indices for Clinical Research Compared with Simple Patient Questionnaires to Assess Function, Pain, and Global Estimates as Rheumatology "Vital Signs" for Usual Clinical Care. Rheumatic Disease Clinics of North America, 2009, 35, 779-786. | 0.8 | 6 |
| 157 | Is a Patient Questionnaire Without a Joint Examination as Undesirable as a Joint Examination Without a Patient Questionnaire?. Journal of Rheumatology, 2014, 41, 619-621. | 1.0 | 6 |
| 158 | Special Issue on Glucocorticoid Therapy in Rheumatic Diseases: Introduction. NeuroImmunoModulation, 2015, 22, 3-5. | 0.9 | 6 |
| 159 | A Self-Report Multidimensional Health Assessment Questionnaire (MDHAQ) for Face-To-Face or Telemedicine Encounters to Assess Clinical Severity (RAPID3) and Screen for Fibromyalgia (FAST) and Depression (DEP). Current Treatment Options in Rheumatology, 2021, 7, 161-181. | 0.6 | 6 |
| 160 | Will Racial and Ethnic Disparities in Health Be Resolved Primarily Outside of Standard Medical Care?. Annals of Internal Medicine, 2004, 141, 224. | 2.0 | 5 |
| 161 | A three-page Standard Protocol to Evaluate Rheumatoid Arthritis (SPERA) for efficient capture of essential data from patients and health professionals in standard clinical care and clinical research. Best Practice and Research in Clinical Rheumatology, 2007, 21, 677-685. | 1.4 | 5 |
| 162 | Quantitative Clinical Rheumatology: "Keep It Simple, Stupid" MDHAQ Function, Pain, Global, and RAPID3 Quantitative Scores to Improve and Document the Quality of Rheumatologic Care. Journal of Rheumatology, 2009, 36, 1099-1100. | 1.0 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Is a Self-Report RAPID3 Score a Reasonable Alternative to a DAS28 in Usual Clinical Care?. Journal of Clinical Rheumatology, 2009, 15, 215-217. | 0.5 | 5 |
| 164 | Documenting the Value of Care for Rheumatoid Arthritis, Analogous to Hypertension, Diabetes, and Hyperlipidemia: Is Control of Individual Patient Self-Report Measures of Global Estimate and Physical Function More Valuable Than Laboratory Tests, Radiographs, Indices, or Remission Criteria?. Journal of Rheumatology, 2013, 40, 1469-1474. | 1.0 | 5 |
| 165 | Hotel-based medicine. Journal of Rheumatology, 2008, 35, 1487-8. | 1.0 | 5 |
| 166 | Electronic eRAPID3 (Routine Assessment of Patient Index Data): opportunities and complexities. Clinical and Experimental Rheumatology, 2016, 34, S49-S53. | 0.4 | 5 |
| 167 | Osteoarthritis is as severe as rheumatoid arthritis: evidence over 40 years according to the same measure in each disease. Clinical and Experimental Rheumatology, 2019, 37 Suppl 120, 7-17. | 0.4 | 5 |
| 168 | Dietary and allergic associations with rheumatoid arthritis: Self-report of 704 patients. Arthritis and Rheumatism, 1990, 3, 189-195. | 6.7 | 4 |
| 169 | Quantitative Measures to Assess Patients with Rheumatic Diseases: 2006 Update. Rheumatic Disease Clinics of North America, 2006, 32, 29-36. | 0.8 | 4 |
| 170 | Are excellent systematic reviews of clinical trials useful for patient care?. Nature Clinical Practice Rheumatology, 2008, 4, 294-295. | 3.2 | 4 |
| 171 | Quality Control of a Medical History: Improving Accuracy with Patient Participation, Supported by a Four-Page Version of the Multidimensional Health Assessment Questionnaire (MDHAQ). Rheumatic Disease Clinics of North America, 2009, 35, 851-860. | 0.8 | 4 |
| 172 | Beyond RAPID3 - practical use of the MDHAQ to improve doctor-patient communication. Bulletin of the NYU Hospital for Joint Diseases, 2010, 68, 223-31. | 0.7 | 4 |
| 173 | Low socioeconomic status and patient questionnaires in osteoarthritis: challenges to a "biomedical model" and value of a complementary "biopsychosocial model". Clinical and Experimental Rheumatology, 2019, 37 Suppl 120, 18-23. | 0.4 | 4 |
| 174 | Quantitative Recording of Physician Clinical Estimates, Beyond a Global Estimate and Formal Joint Count, in Usual Care: Applying the Scientific Method, Using a Simple One-Page Worksheet. Rheumatic Disease Clinics of North America, 2009, 35, 813-817. | 0.8 | 3 |
| 175 | Long-term Glucocorticoid Use in Rheumatoid Arthritis. Journal of Rheumatology, 2021, 48, 1342-1342. | 1.0 | 3 |
| 176 | Comment on: "It can't be zero!" Difficulties in completing patient global assessment in rheumatoid arthritis: a mixed methods study. Rheumatology, 2021, 60, e28-e29. | 0.9 | 3 |
| 177 | Should Quantitative Measures and Management of Rheumatoid Arthritis Include More Than Control of Inflammatory Activity?. Journal of Rheumatology, 2021, , jrheum.210953. | 1.0 | 3 |
| 178 | Assessment of pain and other patient symptoms in routine clinical care as quantitative, standardised, "scientific" data. Clinical and Experimental Rheumatology, 2017, 35 Suppl 107, 13-20. | 0.4 | 3 |
| 179 | Pain and other self-report scores in patients with osteoarthritis indicate generally similar disease burden to patients with rheumatoid arthritis. Clinical and Experimental Rheumatology, 2017, 35 Suppl 107, 88-93. | 0.4 | 3 |
| 180 | Quantitative Clinical Rheumatology: Why Is a Test for Anti-CCP Antibodies Included in Most Routine Care for Rheumatoid Arthritis While a HAQ/MDHAQ Remains Largely a Research Tool?. Journal of Rheumatology, 2009, 36, 1563-1564. | 1.0 | 2 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | Predicting mortality in RA: the quest for useful information. Nature Reviews Rheumatology, 2015, 11, 507-509. | 3.5 | 2 |
| 182 | Pragmatic assessment of exercise in routine care using an MDHAQ: associations with changes in RAPID3 and other clinical variables. Arthritis Research and Therapy, 2016, 18, 199. | 1.6 | 2 |
| 183 | PROMs (MDHAQ/RAPID3) and Physician RheuMetric Measures. , 2016, , 59-99. | | 2 |
| 184 | An evidence-based medical visit for patients with rheumatoid arthritis based on standard, quantitative scientific data from a patient MDHAQ and physician report. Bulletin of the NYU Hospital for Joint Diseases, 2012, 70, 73-94. | 0.7 | 2 |
| 185 | A RheuMetric physician checklist to quantitate levels of inflammation, damage and distress on 0-10 visual analogue scales. Clinical and Experimental Rheumatology, 2017, 35 Suppl 107, 21-25. | 0.4 | 2 |
| 186 | A Standard Protocol to Evaluate Rheumatoid Arthritis (SPERA) for Efficient Capture of Essential Data from a Patient and a Health Professional in a Uniform "Scientific" Format. Rheumatic Disease Clinics of North America, 2009, 35, 843-850. | 0.8 | 1 |
| 187 | Evidence from a Multidimensional Health Assessment Questionnaire (MDHAQ) of the Value of a Biopsychosocial Model to Complement a Traditional Biomedical Model in Care of Patients with Rheumatoid Arthritis. Journal of Rheumatic Diseases, 2016, 23, 212. | 0.4 | 1 |
| 188 | Patient physical function in rheumatoid arthritis. , 2020, , 221-250. | | 1 |
| 189 | Implementing a "Treat-to-Target" Approach for Rheumatoid Arthritis During the COVID-19 Pandemic: Results of a Virtual Learning Collaborative Program. Arthritis Care and Research, 2022, 74, 572-578. | 1.5 | 1 |
| 190 | HAQ and DAS28 for clinical trials over months and MDHAQ, RheuMetric and psycho-socio-economic measures for long-term observations over years?. Rheumatology, 2022, 61, 3884-3886. | 0.9 | 1 |
| 191 | Improving survival in inflammatory rheumatic diseases: a neglected goal?. Current Rheumatology Reports, 2006, 8, 401-403. | 2.1 | 0 |
| 192 | Dr. Pincus, et al reply. Journal of Rheumatology, 2009, 36, 456.2-456. | 1.0 | 0 |
| 193 | Dr. Pincus replies. Journal of Rheumatology, 2009, 36, 443.2-444. | 1.0 | 0 |
| 194 | Limited Value of the Multi-Biomarker Disease Activity Assay Compared to the Routine Assessment of Patient Index Data 3 (RAPID3) Score in the Prognosis of Important Clinical Outcomes in Rheumatoid Arthritis: Comment on the Article by Fleischmann et al and Accompanying Editorial by Davis. Arthritis and Rheumatology, 2017, 69, 866-867. | 2.9 | 0 |
| 195 | 069. "CONSTRUCT VALIDITY, RESPONSIVENESS AND MINIMALLY IMPORTANT DIFFERENCE OF THE ROUTINE ASSESSMENT OF PATIENT INDEX DATA 3 IN PSORIATIC ARTHRITIS. Rheumatology, 2017, 56, . | 0.9 | 0 |
| 196 | THU0091 "DEPRESSION IN PATIENTS WITH RHEUMATOID ARTHRITIS: UNDER RECOGNIZED, UNDERTREATED AND ASSOCIATED WITH POORER CLINICAL STATUS AND LOWER RATES OF REMISSION IN ROUTINE CARE. , 2019, , . | | 0 |
| 197 | THU0477 "A FIBROMYALGIA ASSESSMENT SCREENING TOOL ON A MULTIDIMENSIONAL HEALTH ASSESSMENT QUESTIONNAIRE (MDHAQ) WHICH DOES NOT INCLUDE A SELF-REPORT PAINFUL JOINT COUNT (PAINFUL JC), FAST3NJC, RECOGNIZES FIBROMYALGIA SIMILARLY TO OTHER FAST3 INDICES WHICH INCLUDE A PAINFUL JC. , 2019, , . | | 0 |
| 198 | Functional Status Measures and Indices in Rheumatoid Arthritis: Comment on the Articles by Barber et al and England et al. Arthritis Care and Research, 2020, 72, 1185-1186. | 1.5 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 199 | Limitations of Traditional Randomized Controlled Clinical Trials in Rheumatology. , 2014, , 179-207. | | 0 |
| 200 | Evidence-Based Medicine in Rheumatology: How Does It Differ from Other Diseases?. , 2014, , 1-12. | | 0 |
| 201 | Pain in rheumatic diseases. Clinical and Experimental Rheumatology, 2017, 35 Suppl 107, 1. | 0.4 | 0 |
| 202 | Patient questionnaires in osteoarthritis: what patients teach doctors about their osteoarthritis on a multidimensional health assessment questionnaire (MDHAQ) in clinical trials and clinical care. Clinical and Experimental Rheumatology, 2019, 37 Suppl 120, 100-111. | 0.4 | 0 |