

# Myriam Valero

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/6702323/myriam-valero-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

120  
papers

3,954  
citations

38  
h-index

57  
g-index

122  
ext. papers

4,605  
ext. citations

4  
avg, IF

5.11  
L-index

#	Paper	IF	Citations
120	Development of tools to rapidly identify cryptic species and characterize their genetic diversity in different European kelp species. <i>Journal of Applied Phycology</i> , <b>2021</b> , 33, 4169	3.2	
119	Genomic signatures of clonality in the deep water kelp <i>Laminaria rodriguezii</i> . <i>Molecular Ecology</i> , <b>2021</b> , 30, 1806-1822	5.7	4
118	Evolution of life cycles and reproductive traits: Insights from the brown algae. <i>Journal of Evolutionary Biology</i> , <b>2021</b> , 34, 992-1009	2.3	5
117	Exploring the Genetic Consequences of Clonality in Haplodiplontic Taxa. <i>Journal of Heredity</i> , <b>2021</b> , 112, 92-107	2.4	3
116	After a catastrophe, a little bit of sex is better than nothing: Genetic consequences of a major earthquake on asexual and sexual populations. <i>Evolutionary Applications</i> , <b>2020</b> , 13, 2086-2100	4.8	4
115	Genetic structure of ampho-Atlantic <i>Laminaria digitata</i> (Laminariales, Phaeophyceae) reveals a unique range-edge gene pool and suggests post-glacial colonization of the NW Atlantic. <i>European Journal of Phycology</i> , <b>2020</b> , 55, 517-528	2.2	5
114	How do microbiota associated with an invasive seaweed vary across scales?. <i>Molecular Ecology</i> , <b>2020</b> , 29, 2094-2108	5.7	13
113	Parallelisable non-invasive biomass, fitness and growth measurement of macroalgae and other protists with nephelometry. <i>Algal Research</i> , <b>2020</b> , 46, 101762	5	1
112	Genetic diversity of a marine foundation species, <i>Laminaria hyperborea</i> (Gunnerus) Foslie, along the coast of Ireland. <i>European Journal of Phycology</i> , <b>2020</b> , 55, 310-326	2.2	3
111	Better off alone? Compared performance of monoclonal and polyclonal stands of a cultivated red alga growth. <i>Evolutionary Applications</i> , <b>2020</b> , 13, 905-917	4.8	1
110	Congruence between fine-scale genetic breaks and dispersal potential in an estuarine seaweed across multiple transition zones. <i>ICES Journal of Marine Science</i> , <b>2020</b> , 77, 371-378	2.7	5
109	Phylogeny and Evolution of the Brown Algae. <i>Critical Reviews in Plant Sciences</i> , <b>2020</b> , 39, 281-321	5.6	31
108	Heat stress responses and population genetics of the kelp (Phaeophyceae) across latitudes reveal differentiation among North Atlantic populations. <i>Ecology and Evolution</i> , <b>2020</b> , 10, 9144-9177	2.8	10
107	Seascape Genomics of the Sugar Kelp along the North Eastern Atlantic Latitudinal Gradient. <i>Genes</i> , <b>2020</b> , 11,	4.2	4
106	Dual influence of terrestrial and marine historical processes on the phylogeography of the Brazilian intertidal red alga <i>Gracilaria caudata</i> . <i>Journal of Phycology</i> , <b>2019</b> , 55, 1096-1114	3	9
105	Local Coastal Configuration Rather Than Latitudinal Gradient Shape Clonal Diversity and Genetic Structure of <i>Phymatolithon calcareum</i> Maerl Beds in North European Atlantic. <i>Frontiers in Marine Science</i> , <b>2019</b> , 6,	4.5	9
104	Comparative phylogeography of six red algae along the Antarctic Peninsula: extreme genetic depletion linked to historical bottlenecks and recent expansion. <i>Polar Biology</i> , <b>2018</b> , 41, 827-837	2	12

103	Combining niche shift and population genetic analyses predicts rapid phenotypic evolution during invasion. <i>Evolutionary Applications</i> , <b>2018</b> , 11, 781-793	4.8	38
102	Genetic Diversity in the UV Sex Chromosomes of the Brown Alga. <i>Genes</i> , <b>2018</b> , 9,	4.2	10
101	Past climate changes and strong oceanographic barriers structured low-latitude genetic relics for the golden kelp <i>Laminaria ochroleuca</i> . <i>Journal of Biogeography</i> , <b>2018</b> , 45, 2326-2336	4.1	26
100	Entangled fates of holobiont genomes during invasion: nested bacterial and host diversities in <i>Caulerpa taxifolia</i> . <i>Molecular Ecology</i> , <b>2017</b> , 26, 2379-2391	5.7	25
99	Population dynamics of temperate kelp forests near their low-latitude limit. <i>Aquatic Botany</i> , <b>2017</b> , 139, 8-18	1.8	6
98	Multi-scale drivers of community diversity and composition across tidal heights: an example on temperate seaweed communities. <i>Journal of Ecology</i> , <b>2017</b> , 105, 1791-1805	6	6
97	High-density genetic map and identification of QTLs for responses to temperature and salinity stresses in the model brown alga <i>Ectocarpus</i> . <i>Scientific Reports</i> , <b>2017</b> , 7, 43241	4.9	25
96	Hybridization between two cryptic filamentous brown seaweeds along the shore: analysing pre- and postzygotic barriers in populations of individuals with varying ploidy levels. <i>Molecular Ecology</i> , <b>2017</b> , 26, 3497-3512	5.7	16
95	ClonEstiMate, a Bayesian method for quantifying rates of clonality of populations genotyped at two-time steps. <i>Molecular Ecology Resources</i> , <b>2017</b> , 17, e251-e267	8.4	11
94	Species delimitation and phylogeographic analyses in the <i>Ectocarpus</i> subgroup <i>siliculosi</i> (Ectocarpales, Phaeophyceae). <i>Journal of Phycology</i> , <b>2017</b> , 53, 17-31	3	35
93	Reproductive strategies and population genetic structure of <i>Fucus</i> spp. across a northeast Atlantic biogeographic transition. <i>Aquatic Living Resources</i> , <b>2017</b> , 30, 16	1.5	4
92	Perspectives on domestication research for sustainable seaweed aquaculture. <i>Perspectives in Phycology</i> , <b>2017</b> , 4, 33-46	3.1	41
91	Development and characterization of microsatellite markers in two agarophyte species, <i>Gracilaria birdiae</i> and <i>Gracilaria caudata</i> (Gracilariaceae, Rhodophyta), using next-generation sequencing. <i>Journal of Applied Phycology</i> , <b>2016</b> , 28, 653-662	3.2	9
90	Evidence for parasite-mediated selection during short-lasting toxic algal blooms. <i>Proceedings of the Royal Society B: Biological Sciences</i> , <b>2016</b> , 283,	4.4	8
89	Status, trends and drivers of kelp forests in Europe: an expert assessment. <i>Biodiversity and Conservation</i> , <b>2016</b> , 25, 1319-1348	3.4	72
88	Climate Oscillations, Range Shifts and Phylogeographic Patterns of North Atlantic Fucaceae <b>2016</b> , 279-308		20
87	Phylogeography of Seaweeds in the South East Pacific: Complex Evolutionary Processes Along a Latitudinal Gradient <b>2016</b> , 251-277		14
86	Lack of fine-scale genetic structure and distant mating in natural populations of <i>Fucus vesiculosus</i> . <i>Marine Ecology - Progress Series</i> , <b>2016</b> , 544, 131-142	2.6	7

85	Microsatellite markers and cytoplasmic sequences reveal contrasting pattern of spatial genetic structure in the red algae species complex <i>Mazzaella laminarioides</i> . <i>Journal of Phycology</i> , <b>2016</b> , 52, 806-816	3	3
84	Deep reefs are climatic refugia for genetic diversity of marine forests. <i>Journal of Biogeography</i> , <b>2016</b> , 43, 833-844	4.1	62
83	Characterization of newly developed expressed sequence tag-derived microsatellite markers revealed low genetic diversity within and low connectivity between European <i>Saccharina latissima</i> populations. <i>Journal of Applied Phycology</i> , <b>2016</b> , 28, 3057-3070	3.2	14
82	How does molecular-assisted identification affect our estimation of $\pi$ and $\theta$ biodiversity? An example from understory red seaweeds (Rhodophyta) of <i>Laminaria</i> kelp forests in Brittany, France. <i>Genetica</i> , <b>2015</b> , 143, 207-23	1.5	8
81	Contrasting timing of life stages across latitudes: a case study of a marine forest-forming species. <i>European Journal of Phycology</i> , <b>2015</b> , 50, 361-369	2.2	5
80	Barcoding of Cryptic Stages of Marine Brown Algae Isolated from Incubated Substratum Reveals High Diversity in Acinetosporaceae (Ectocarpales, Phaeophyceae)1. <i>Cryptogamie, Algologie</i> , <b>2015</b> , 36, 3	0.7	35
79	Species are hypotheses: avoid connectivity assessments based on pillars of sand. <i>Molecular Ecology</i> , <b>2015</b> , 24, 525-44	5.7	138
78	Response of kelps from different latitudes to consecutive heat shock. <i>Journal of Experimental Marine Biology and Ecology</i> , <b>2015</b> , 463, 57-62	2.1	20
77	Evolution and maintenance of haploid-diploid life cycles in natural populations: The case of the marine brown alga <i>Ectocarpus</i> . <i>Evolution; International Journal of Organic Evolution</i> , <b>2015</b> , 69, 1808-22	3.8	36
76	O father where art thou? Paternity analyses in a natural population of the haploid-diploid seaweed <i>Chondrus crispus</i> . <i>Heredity</i> , <b>2015</b> , 114, 185-94	3.6	29
75	Contrasting genetic diversity patterns in two sister kelp species co-distributed along the coast of Brittany, France. <i>Molecular Ecology</i> , <b>2014</b> , 23, 2669-85	5.7	42
74	A haploid system of sex determination in the brown alga <i>Ectocarpus</i> sp. <i>Current Biology</i> , <b>2014</b> , 24, 1945-573	5.7	93
73	Twenty years of observed and predicted changes in subtidal red seaweed assemblages along a biogeographical transition zone: inferring potential causes from environmental data. <i>Journal of Biogeography</i> , <b>2014</b> , 41, 2293-2306	4.1	50
72	Development and multiplexing of the first microsatellite markers in a coralline red alga ( <i>Phymatolithon calcareum</i> , Rhodophyta). <i>Phycologia</i> , <b>2014</b> , 53, 474-479	2.7	7
71	<i>Kallymenia crouaniorum</i> (Kallymeniaceae, Rhodophyta), a new red algal species from the <i>Laminaria</i> hyperborea understory community. <i>European Journal of Phycology</i> , <b>2014</b> , 49, 493-507	2.2	8
70	Spatiotemporal changes in the genetic diversity of harmful algal blooms caused by the toxic dinoflagellate <i>Alexandrium minutum</i> . <i>Molecular Ecology</i> , <b>2014</b> , 23, 549-60	5.7	41
69	Examining the bank of microscopic stages in kelps using culturing and barcoding. <i>European Journal of Phycology</i> , <b>2014</b> , 49, 128-133	2.2	12
68	Alteration of sexual reproduction and genetic diversity in the kelp species <i>Laminaria digitata</i> at the southern limit of its range. <i>PLoS ONE</i> , <b>2014</b> , 9, e102518	3.7	30

67	Tracing the trans-pacific evolutionary history of a domesticated Seaweed ( <i>Gracilaria chilensis</i> ) with archaeological and genetic data. <i>PLoS ONE</i> , <b>2014</b> , 9, e114039	3.7	31
66	Intergametophytic selfing and microgeographic genetic structure shape populations of the intertidal red seaweed <i>Chondrus crispus</i> . <i>Molecular Ecology</i> , <b>2013</b> , 22, 3242-60	5.7	54
65	Management and conservation of the kelp species <i>Laminaria digitata</i> : using genetic tools to explore the potential exporting role of the MPA Parc naturel marin d'Iroise. <i>Aquatic Living Resources</i> , <b>2013</b> , 26, 197-205	1.5	16
64	Decline in Kelp in West Europe and Climate. <i>PLoS ONE</i> , <b>2013</b> , 8, e66044	3.7	106
63	High and distinct range-edge genetic diversity despite local bottlenecks. <i>PLoS ONE</i> , <b>2013</b> , 8, e68646	3.7	66
62	Drifting fronds and drifting alleles: range dynamics, local dispersal and habitat isolation shape the population structure of the estuarine seaweed <i>Fucus ceranoides</i> . <i>Journal of Biogeography</i> , <b>2012</b> , 39, 1167-1178	4.4	44
61	Fine-scale genetic breaks driven by historical range dynamics and ongoing density-barrier effects in the estuarine seaweed <i>Fucus ceranoides</i> L. <i>BMC Evolutionary Biology</i> , <b>2012</b> , 12, 78	3	40
60	Temperature effects on gametophyte life-history traits and geographic distribution of two cryptic kelp species. <i>PLoS ONE</i> , <b>2012</b> , 7, e39289	3.7	37
59	SEX RATIO VARIATION IN THE LESSONIA NIGRESCENS COMPLEX (LAMINARIALES, PHAEOPHYCEAE): EFFECT OF LATITUDE, TEMPERATURE, AND MARGINALITY(1). <i>Journal of Phycology</i> , <b>2011</b> , 47, 5-12	3	31
58	GENETIC POPULATION STRUCTURE AND MATING SYSTEM IN CHONDRUS CRISPUS (RHODOPHYTA). <i>Journal of Phycology</i> , <b>2011</b> , 47, 440-450	3	42
57	THE LESSONIA NIGRESCENS SPECIES COMPLEX (LAMINARIALES, PHAEOPHYCEAE) SHOWS STRICT PARAPATRY AND COMPLETE REPRODUCTIVE ISOLATION IN A SECONDARY CONTACT ZONE(1). <i>Journal of Phycology</i> , <b>2011</b> , 47, 894-903	3	44
56	MICROSATELLITE DEVELOPMENT IN RHODOPHYTA USING HIGH-THROUGHPUT SEQUENCE DATA(1). <i>Journal of Phycology</i> , <b>2011</b> , 47, 1258-65	3	10
55	Evolution and diversification within the intertidal brown macroalgae <i>Fucus spiralis</i> /F. <i>vesiculosus</i> species complex in the North Atlantic. <i>Molecular Phylogenetics and Evolution</i> , <b>2011</b> , 58, 283-96	4.1	61
54	Dinucleotide microsatellite markers in the genus <i>Caulerpa</i> . <i>Journal of Applied Phycology</i> , <b>2011</b> , 23, 715-719	3.2	5
53	Waterborne signaling primes the expression of elicitor-induced genes and buffers the oxidative responses in the brown alga <i>Laminaria digitata</i> . <i>PLoS ONE</i> , <b>2011</b> , 6, e21475	3.7	19
52	Surfing the wave on a borrowed board: range expansion and spread of introgressed organellar genomes in the seaweed <i>Fucus ceranoides</i> L. <i>Molecular Ecology</i> , <b>2010</b> , 19, 4812-22	5.7	56
51	The phylogeographic architecture of the furoid seaweed <i>Ascophyllum nodosum</i> : an intertidal marine tree-land survivor of more than one glacial-interglacial cycle. <i>Journal of Biogeography</i> , <b>2010</b> , 37, 842-856	4.1	83
50	DEFENSE EVOLUTION IN THE GRACILARIACEAE (RHODOPHYTA): SUBSTRATE-REGULATED OXIDATION OF AGAR OLIGOSACCHARIDES IS MORE ANCIENT THAN THE OLIGOAGAR-ACTIVATED OXIDATIVE BURST1. <i>Journal of Phycology</i> , <b>2010</b> , 46, 958-968	3	14

49	Delineation of Two Sibling Red Algal Species, <i>Gracilaria Gracilis</i> and <i>Gracilaria Dura</i> (Gracilariales, Rhodophyta), Using Multiple DNA Markers: Resurrection of the Species <i>G. Dura</i> Previously Described in the Northern Atlantic 200 Years Ago1. <i>Journal of Phycology</i> , <b>2010</b> , 46, 720-727	3	30
48	<i>Fucus vesiculosus</i> and <i>spiralis</i> species complex: a nested model of local adaptation at the shore level. <i>Marine Ecology - Progress Series</i> , <b>2010</b> , 405, 163-174	2.6	36
47	Phylogeographic analyses of the 30 degrees S south-east Pacific biogeographic transition zone establish the occurrence of a sharp genetic discontinuity in the kelp <i>Lessonia nigrescens</i> : vicariance or parapatry?. <i>Molecular Phylogenetics and Evolution</i> , <b>2009</b> , 53, 679-93	4.1	84
46	PERMANENT GENETIC RESOURCES: Isolation of microsatellite loci from the kelp, <i>Saccorhiza polyschides</i> (Heterokontophyta, incertae sedis). <i>Molecular Ecology Resources</i> , <b>2008</b> , 8, 406-8	8.4	4
45	Molecular characterisation and development of rapid molecular methods to identify species of Gracilariaceae from the Atlantic coast of Morocco. <i>Aquatic Botany</i> , <b>2008</b> , 89, 324-330	1.8	35
44	Genetic variation in wild and cultivated populations of the haploid-diploid red alga <i>Gracilaria chilensis</i> : how farming practices favor asexual reproduction and heterozygosity. <i>Evolution; International Journal of Organic Evolution</i> , <b>2008</b> , 62, 1500-19	3.8	103
43	Complex life cycles of multicellular eukaryotes: new approaches based on the use of model organisms. <i>Gene</i> , <b>2007</b> , 406, 152-70	3.8	92
42	Development of microsatellites DNA markers in the cultivated seaweed, <i>Gracilaria chilensis</i> (Gracilariales, Rhodophyta). <i>Molecular Ecology Notes</i> , <b>2005</b> , 5, 155-157		20
41	GENETIC ISOLATION BETWEEN THREE CLOSELY RELATED TAXA: <i>FUCUS VESICULOSUS</i> , <i>F. SPIRALIS</i> , AND <i>F. CERANOIDES</i> (PHAOPHYCEAE)1. <i>Journal of Phycology</i> , <b>2005</b> , 41, 900-905	3	36
40	Analysis of sexual phenotype and prezygotic fertility in natural populations of <i>Fucus spiralis</i> , <i>F. vesiculosus</i> (Fucaceae, Phaeophyceae) and their putative hybrids. <i>European Journal of Phycology</i> , <b>2005</b> , 40, 397-407	2.2	30
39	Analysis of rDNA ITS1 indels in <i>Caulerpa taxifolia</i> (Chlorophyta) supports a derived, incipient species status for the invasive strain. <i>European Journal of Phycology</i> , <b>2004</b> , 39, 83-92	2.2	41
38	REDUCED GENETIC DIVERSITY AND INCREASED POPULATION DIFFERENTIATION IN PERIPHERAL AND OVERHARVESTED POPULATIONS OF <i>GIGARTINA SKOTTSBERGII</i> (RHODOPHYTA, GIGARTINALES) IN SOUTHERN CHILE1. <i>Journal of Phycology</i> , <b>2004</b> , 40, 454-462	3	35
37	MOLECULAR IDENTIFICATION OF TWO SIBLING SPECIES UNDER THE NAME <i>GRACILARIA CHILENSIS</i> (RHODOPHYTA, GRACILARIALES)1,3. <i>Journal of Phycology</i> , <b>2004</b> , 40, 742-747	3	44
36	Mating system and gene flow in the red seaweed <i>Gracilaria gracilis</i> : effect of haploid-diploid life history and intertidal rocky shore landscape on fine-scale genetic structure. <i>Heredity</i> , <b>2004</b> , 92, 289-98	3.6	70
35	Current patterns, habitat discontinuities and population genetic structure: the case of the kelp <i>Laminaria digitata</i> in the English Channel. <i>Marine Ecology - Progress Series</i> , <b>2003</b> , 253, 111-121	2.6	85
34	Polymerase chain reaction-single strand conformation polymorphism analyses of nuclear and chloroplast DNA provide evidence for recombination, multiple introductions and nascent speciation in the <i>Caulerpa taxifolia</i> complex. <i>Molecular Ecology</i> , <b>2002</b> , 11, 2317-25	5.7	36
33	Non-random mating in controlled multiple-donor crosses in <i>Gracilaria gracilis</i> (Gracilariaceae, Rhodophyta). <i>European Journal of Phycology</i> , <b>2002</b> , 37, 179-190	2.2	7
32	HIERARCHICAL SPATIAL STRUCTURE AND DISCRIMINANT ANALYSIS OF GENETIC DIVERSITY IN THE RED ALGA <i>MAZZAELLA LAMINARIOIDES</i> (GIGARTINALES, RHODOPHYTA). <i>Journal of Phycology</i> , <b>2001</b> , 37, 705-716	3	49

31	Population dynamics and stage structure in a haploid-diploid red seaweed, <i>Gracilaria gracilis</i> . <i>Journal of Ecology</i> , <b>2001</b> , 89, 436-450	6	49
30	Phylogenetic analyses of <i>Caulerpa taxifolia</i> (Chlorophyta) and of its associated bacterial microflora provide clues to the origin of the Mediterranean introduction. <i>Molecular Ecology</i> , <b>2001</b> , 10, 931-46	5.7	81
29	The spatial structure of sexual and cytonuclear polymorphism in the gynodioecious <i>Beta vulgaris</i> ssp. <i>maritima</i> : I/ at a local scale. <i>Genetics</i> , <b>2001</b> , 157, 1699-710	4	44
28	A model for the evolution of high frequencies of males in an androdioecious plant based on a cross-compatibility advantage of males. <i>Heredity</i> , <b>2000</b> , 85 Pt 5, 413-22	3.6	19
27	Identification of random amplified polymorphic DNA (RAPD) markers highly linked to sex determination in the red alga <i>Gracilaria gracilis</i> . <i>Molecular Ecology</i> , <b>1999</b> , 8, 1533-8	5.7	28
26	Performance of nonhotile male gametes in the sea: analysis of paternity and fertilization success in a natural population of a red seaweed, <i>Gracilaria gracilis</i> . <i>Proceedings of the Royal Society B: Biological Sciences</i> , <b>1999</b> , 266, 1879-1886	4.4	52
25	Characterization of microsatellite markers in the red alga <i>Gracilaria gracilis</i> . <i>Molecular Ecology</i> , <b>1999</b> , 8, 700-2	5.7	10
24	Genetic structure of natural populations of <i>Gelidium</i> species: A re-evaluation of results. <i>Journal of Applied Phycology</i> , <b>1998</b> , 10, 279-284	3.2	23
23	MEDITERRANEAN CAULERPA TAXIFOLIA AND C. MEXICANA (CHLOROPHYTA) ARE NOT CONSPECIFIC. <i>Journal of Phycology</i> , <b>1998</b> , 34, 850-856	3	33
22	Distribution of chloroplast DNA diversity within and among populations in gynodioecious <i>Beta vulgaris</i> ssp. <i>maritima</i> (Chenopodiaceae). <i>Molecular Ecology</i> , <b>1998</b> , 7, 1193-1204	5.7	25
21	Short allele dominance as a source of heterozygote deficiency at microsatellite loci: experimental evidence at the dinucleotide locus Gv1CT in <i>Gracilaria gracilis</i> (Rhodophyta). <i>Molecular Ecology</i> , <b>1998</b> , 7, 1569-1573	5.7	186
20	Isolation and characterization of microsatellite markers in the nuclear genome of the brown alga <i>Laminaria digitata</i> (Phaeophyceae). <i>Molecular Ecology</i> , <b>1998</b> , 7, 1778-80	5.7	35
19	Variation in Sexual and Asexual Reproduction among Young and Old Populations of the Perennial Macrophyte <i>Sparganium erectum</i> . <i>Oikos</i> , <b>1998</b> , 82, 139	4	97
18	SINGLE LOCUS MICROSATELLITES IN GRACILARIALES (RHODOPHYTA): HIGH LEVEL OF GENETIC VARIABILITY WITHIN GRACILARIA GRACILIS AND CONSERVATION IN RELATED SPECIES1. <i>Journal of Phycology</i> , <b>1997</b> , 33, 868-880	3	54
17	The Evolving Genetic History of a Population of <i>Lathyrus sylvestris</i> : Evidence From Temporal and Spatial Genetic Structure. <i>Evolution; International Journal of Organic Evolution</i> , <b>1996</b> , 50, 1808	3.8	14
16	THE EVOLVING GENETIC HISTORY OF A POPULATION OF LATHYRUS SYLVESTRIS: EVIDENCE FROM TEMPORAL AND SPATIAL GENETIC STRUCTURE. <i>Evolution; International Journal of Organic Evolution</i> , <b>1996</b> , 50, 1808-1821	3.8	26
15	Variation of reproductive success in a haplo-diploid Red Alga, <i>Gracilaria Verrucosa</i> : effects of parental identities and crossing distance. <i>American Journal of Botany</i> , <b>1993</b> , 80, 1379	2.7	7
14	Variation of reproductive success in a haplo-diploid Red Alga, <i>Gracilaria Verrucosa</i> : effects of parental identities and crossing distance. <i>American Journal of Botany</i> , <b>1993</b> , 80, 1379-1391	2.7	7

13	Evolution of the alternation of haploid and diploid phases in life cycles. II. Maintenance of the haplo-diplontic cycle. <i>Journal of Evolutionary Biology</i> , <b>1993</b> , 6, 263-280	2.3	26
12	Differences in response between haploid and diploid isomorphic phases of <i>Gracilaria verrucosa</i> (Rhodophyta: Gigartinales) exposed to artificial environmental conditions. <i>Hydrobiologia</i> , <b>1993</b> , 260-261, 131-137	2.4	51
11	Differences in response between haploid and diploid isomorphic phases of <i>Gracilaria verrucosa</i> (Rhodophyta: Gigartinales) exposed to artificial environmental conditions <b>1993</b> , 131-137		2
10	Evolution of alternation of haploid and diploid phases in life cycles. <i>Trends in Ecology and Evolution</i> , <b>1992</b> , 7, 25-9	10.9	82
9	Cole, K. M. and R. G. Sheath (eds.) 1990. Biology of the red algae. Cambridge University Press (Cambridge, New York, Port Chester, Melbourne, Sydney), USA, 517 pp. \$65.. <i>Journal of Evolutionary Biology</i> , <b>1992</b> , 5, 533-536	2.3	0
8	Transition from haploidy to diploidy. <i>Nature</i> , <b>1991</b> , 351, 315-7	50.4	115
7	Discriminant alleles and discriminant analysis: efficient characters to separate closely related species: the example of <i>Lathyrus latifolius</i> L. and <i>Lathyrus sylvestris</i> L. (Leguminosae). <i>Botanical Journal of the Linnean Society</i> , <b>1991</b> , 107, 139-161	2.2	1
6	Genetic variation between and within populations of a perennial grass: <i>Arrhenatherum elatius</i> . <i>Heredity</i> , <b>1990</b> , 65, 179-188	3.6	39
5	What controls haploid-diploid ratio in the red alga, <i>Gracilaria verrucosa</i> ?. <i>Journal of Evolutionary Biology</i> , <b>1989</b> , 2, 317-338	2.3	66
4	Spatio-temporal variation of male sterile frequencies in two natural populations of <i>Beta maritima</i> . <i>Heredity</i> , <b>1989</b> , 63, 395-400	3.6	19
3	Effect of Ovule Position in the Pod on Patterns of Seed Formation in Two Species of <i>Lathyrus</i> (Leguminosae: Papilionoideae). <i>American Journal of Botany</i> , <b>1988</b> , 75, 1714	2.7	20
2	EFFECT OF OVULE POSITION IN THE POD ON PATTERNS OF SEED FORMATION IN TWO SPECIES OF LATHYRUS (LEGUMINOSAE: PAPILIONOIDEAE). <i>American Journal of Botany</i> , <b>1988</b> , 75, 1714-1731	2.7	17
1	Evolution of life cycles and reproductive traits: insights from the brown algae		5