

# Jun-Ei Obata

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/670782/publications.pdf>

Version: 2024-02-01

28  
papers

291  
citations

933447

10  
h-index

940533

16  
g-index

28  
all docs

28  
docs citations

28  
times ranked

575  
citing authors

#	ARTICLE	IF	CITATIONS
1	A Propensity Score Matched Analysis of Statin Effects on Major Adverse Cardiac Events after Percutaneous Coronary Intervention in Patients Over 75 Years Old. <i>Internal Medicine</i> , 2022, , .	0.7	2
2	Impact of persistent endothelial dysfunction in an infarct-related coronary artery on future major adverse cardiovascular event occurrence in STEMI survivors. <i>Heart and Vessels</i> , 2021, 36, 472-482.	1.2	5
3	Echoluency of carotid plaque is useful for selecting high-risk patients with chronic coronary artery disease who benefit from intensive lipid-lowering therapy. <i>Journal of Cardiology</i> , 2021, 77, 590-598.	1.9	2
4	Improvement in Brachial Endothelial Vasomotor Function and Brachial-Ankle Pulse Wave Velocity Reduces the Residual Risk for Cardiovascular Events after Optimal Medical Treatment in Patients with Coronary Artery Disease. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 1133-1144.	2.0	12
5	Takotsubo Cardiomyopathy With Leaflet Valve Thrombosis 2 Years After Transcatheter Aortic Valve Replacement. <i>JACC: Cardiovascular Interventions</i> , 2021, 14, e331-e331.	2.9	0
6	Stratification Analysis of Statin Effect on Major Adverse Cardiac Events after Percutaneous Coronary Intervention in Patients on Hemodialysis. <i>Journal of Cardiovascular Pharmacology</i> , 2021, Publish Ahead of Print, .	1.9	0
7	Sustained endothelial dysfunction in the infarct-related coronary artery is associated with left ventricular adverse remodeling in survivors of ST-segment elevation myocardial infarction. <i>Journal of Cardiology</i> , 2020, 75, 261-269.	1.9	5
8	Long-Term Follow-up of a Coronary Artery Aneurysm After Biodegradable-Polymer Sirolimus-Eluting Stent Implantation. <i>Circulation Journal</i> , 2020, 84, 1196.	1.6	0
9	Persistent Myocardial Production of Follistatin-like 1 Is Associated With Left Ventricular Adverse Remodeling in Patients With Myocardial Infarction. <i>Journal of Cardiac Failure</i> , 2020, 26, 733-738.	1.7	6
10	Failure of drug-coated balloon angioplasty to treat bare metal in-stent restenosis accompanied by late stent thrombosis but successful treatment of binary in-stent restenosis. <i>Journal of Cardiology Cases</i> , 2019, 20, 84-87.	0.5	2
11	Remnant Lipoproteins Are Residual Risk Factor for Future Cardiovascular Events in Patients With Stable Coronary Artery Disease and On-Statins Low-Density Lipoprotein Cholesterol Levels <math>\leq 70</math> mg/dL. <i>Circulation Journal</i> , 2019, 83, 1302-1308.	1.6	34
12	Persistent Dysfunction of Coronary Endothelial Vasomotor Responses is Related to Atheroma Plaque Progression in the Infarct-Related Coronary Artery of AMI Survivors. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 1062-1074.	2.0	12
13	Pulmonary Vascular Resistance Is Associated With Brachial-Ankle Pulse-Wave Velocity and Adverse Clinical Outcomes in Patients With Heart Failure With Preserved Ejection Fraction. <i>Journal of Cardiac Failure</i> , 2019, 25, 725-732.	1.7	4
14	High levels of stromal cell-derived factor-1 predict short-term progression of renal dysfunction in patients with coronary artery disease. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 920-927.	1.6	3
15	Effect of coronary artery spasm on long-term outcomes in survivors of acute myocardial infarction. <i>International Journal of Cardiology</i> , 2018, 257, 7-11.	1.7	11
16	Angiotensin II receptor blockers suppress the release of stromal cell-derived factor-1 from infarcted myocardium in patients with acute myocardial infarction. <i>Journal of Cardiology</i> , 2018, 71, 367-374.	1.9	4
17	Human soluble phospholipase A <sub>2</sub> receptor is an inhibitor of the integrin-mediated cell migratory response to collagen-I. <i>American Journal of Physiology - Cell Physiology</i> , 2018, 315, C398-C408.	4.6	10
18	Remnant lipoproteinemia predicts cardiovascular events in patients with type 2 diabetes and chronic kidney disease. <i>Journal of Cardiology</i> , 2017, 69, 529-535.	1.9	24

#	ARTICLE	IF	CITATIONS
19	Echolucency of the carotid artery is associated with short-term plaque progression and positive remodeling in the culprit coronary artery in AMI survivors. <i>Journal of Cardiology</i> , 2017, 70, 438-445.	1.9	5
20	Impact of target vessel on long-term cardiac mortality after successful chronic total occlusion percutaneous coronary intervention: Insights from a Japanese multicenter registry. <i>International Journal of Cardiology</i> , 2017, 245, 77-82.	1.7	13
21	High levels of stromal cell-derived factor-1 $\pm$ predict secondary cardiac events in stable patients with a history of myocardial infarction. <i>Journal of Cardiology</i> , 2017, 69, 320-325.	1.9	12
22	A Case of Ruptured Vertebrobasilar Junction Aneurysm Associated with Subclavian Steal Phenomenon. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017, 26, e160-e164.	1.6	5
23	Assessment of carotid plaque neovascularization using quantitative analysis of contrast-enhanced ultrasound imaging is useful for risk stratification in patients with coronary artery disease. <i>International Journal of Cardiology</i> , 2015, 195, 113-119.	1.7	30
24	Sustained myocardial production of stromal cell-derived factor-1 $\pm$ was associated with left ventricular adverse remodeling in patients with myocardial infarction. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2015, 309, H1764-H1771.	3.2	16
25	C $\alpha$ -type lectin $\alpha$ -like domain and fibronectin $\alpha$ -like type II domain of phospholipase A <sub>2</sub> receptor 1 modulate binding and migratory responses to collagen. <i>FEBS Letters</i> , 2015, 589, 829-835.	2.8	22
26	The combined assessment of flow-mediated dilation of the brachial artery and brachial-ankle pulse wave velocity improves the prediction of future coronary events in patients with chronic coronary artery disease. <i>Journal of Cardiology</i> , 2014, 64, 179-184.	1.9	34
27	Abstract 11733: Assessment of Carotid Plaque Neovascularization Using Quantitative Analysis of Contrast-Enhanced Ultrasound Imaging is Useful for Risk Stratification in Patients With Coronary Artery Disease. <i>Circulation</i> , 2014, 130, .	1.6	0
28	Insulin resistance negatively affects long-term outcome in non-diabetic patients with coronary artery disease after therapies to reduce atherosclerotic risk factors. <i>Journal of Cardiology</i> , 2013, 62, 348-353.	1.9	18