

Stefan Theodor Schwarz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6711493/publications.pdf>

Version: 2024-02-01

14
papers

1,226
citations

933447

10
h-index

1058476

14
g-index

14
all docs

14
docs citations

14
times ranked

1655
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | <sc>Neuromelaninâ€MRI</sc> to Quantify and Track Nigral Depigmentation in Parkinson's Disease: A Multicenter Longitudinal Study Using Templateâ€Based Standardized Analysis. Movement Disorders, 2022, 37, 1028-1039. | 3.9 | 12 |
| 2 | Coordinate based meta-analysis of motor functional imaging in Parkinsonâ€™s: disease-specific patterns and modulation by dopamine replacement and deep brain stimulation. Brain Imaging and Behavior, 2020, 14, 1263-1280. | 2.1 | 7 |
| 3 | Relationship between neuromelanin and dopamine terminals within the Parkinsonâ€™s nigrostriatal system. Brain, 2019, 142, 2023-2036. | 7.6 | 48 |
| 4 | Parkinson's disease related signal change in the nigrosomes 1â€™5 and the substantia nigra using T2* weighted 7T MRI. NeuroImage: Clinical, 2018, 19, 683-689. | 2.7 | 39 |
| 5 | Limbic grey matter changes in early Parkinson's disease. Human Brain Mapping, 2017, 38, 3566-3578. | 3.6 | 25 |
| 6 | MRI of the Swallow Tail Sign: A Useful Marker in the Diagnosis of Lewy Body Dementia?. American Journal of Neuroradiology, 2017, 38, 1737-1741. | 2.4 | 50 |
| 7 | In Vivo Assessment of Brainstem Depigmentation in Parkinson Disease: Potential as a Severity Marker for Multicenter Studies. Radiology, 2017, 283, 789-798. | 7.3 | 84 |
| 8 | Protocol of a single group prospective observational study on the diagnostic value of 3T susceptibility weighted MRI of nigrosome-1 in patients with parkinsonian symptoms: the N3<i>i>PD study (nigrosomalironimagingin Parkinsonâ€™s disease). BMJ Open, 2017, 7, e016904. | 1.9 | 5 |
| 9 | MR Imaging of the Substantia Nigra for the Diagnosis of Parkinson Disease. Radiology, 2014, 273, 627-628. | 7.3 | 1 |
| 10 | The â€™Swallow Tailâ€™ Appearance of the Healthy Nigrosome â€™ A New Accurate Test of Parkinson's Disease: A Case-Control and Retrospective Cross-Sectional MRI Study at 3T. PLoS ONE, 2014, 9, e93814. | 2.5 | 252 |
| 11 | Diffusion tensor imaging of nigral degeneration in Parkinson's disease: A region-of-interest and voxel-based study at 3T and systematic review with meta-analysis. NeuroImage: Clinical, 2013, 3, 481-488. | 2.7 | 148 |
| 12 | Visualization of nigrosome 1 and its loss in PD. Neurology, 2013, 81, 534-540. | 1.1 | 208 |
| 13 | High resolution magnetic susceptibility mapping of the substantia nigra in Parkinson's disease. Journal of Magnetic Resonance Imaging, 2012, 35, 48-55. | 3.4 | 189 |
| 14 | T1-Weighted MRI shows stage-dependent substantia nigra signal loss in Parkinson's disease. Movement Disorders, 2011, 26, 1633-1638. | 3.9 | 158 |