## Shin Yonehara

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6729325/publications.pdf

Version: 2024-02-01

686830 996533 3,377 15 13 15 citations h-index g-index papers 16 16 16 2528 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. Cell, 1991, 66, 233-243.	13.5	2,763
2	Affinity labeling displays the stepwise activation of ICE-related proteases by Fas, staurosporine, and CrmA-sensitive caspase-8. Oncogene, 1997, 14, 2741-2752.	2.6	118
3	Human T-cell leukemia virus type-I oncoprotein Tax inhibits Fas-mediated apoptosis by inducing cellular FLIP through activation of NF-1ºB. Genes To Cells, 2006, 11, 177-191.	0.5	72
4	Phosphorylation and Dimerization Regulate Nucleocytoplasmic Shuttling of Mammalian STE20-like Kinase (MST). Journal of Biological Chemistry, 2002, 277, 12351-12358.	1.6	71
5	Interaction of FLASH with Arsenite Resistance Protein 2 Is Involved in Cell Cycle Progression at S Phase. Molecular and Cellular Biology, 2009, 29, 4729-4741.	1.1	65
6	Molecular cloning and characterization of mouse caspase-8. FEBS Journal, 1998, 253, 399-405.	0.2	49
7	Protein-driven RNA nanostructured devices that function in vitro and control mammalian cell fate. Nature Communications, 2017, 8, 540.	5.8	40
8	Protease Activity of Procaspase-8 Is Essential for Cell Survival by Inhibiting Both Apoptotic and Nonapoptotic Cell Death Dependent on Receptor-interacting Protein Kinase 1 (RIP1) and RIP3. Journal of Biological Chemistry, 2012, 287, 41165-41173.	1.6	39
9	Inflammasome and Fas-Mediated IL- $\hat{\Pi}^2$ Contributes to Th17/Th1 Cell Induction in Pathogenic Bacterial Infection In Vivo. Journal of Immunology, 2017, 199, 1122-1130.	0.4	38
10	Loss of Somatostatin Receptor Subtype 2 Promotes Growth of KRAS-Induced Pancreatic Tumors in Mice by Activating PI3K Signaling and Overexpression of CXCL16. Gastroenterology, 2015, 148, 1452-1465.	0.6	36
11	Molecular Mechanism of Apoptosis by Amyloid $\hat{l}^2$ -Protein Fibrils Formed on Neuronal Cells. ACS Chemical Neuroscience, 2020, 11, 796-805.	1.7	28
12	Caspase-8, receptor-interacting protein kinase 1 (RIPK1), and RIPK3 regulate retinoic acid-induced cell differentiation and necroptosis. Cell Death and Differentiation, 2020, 27, 1539-1553.	5.0	25
13	Interferon- $\hat{l}^3$ induces the cell surface exposure of phosphatidylserine by activating the protein MLKL in the absence of caspase-8 activity. Journal of Biological Chemistry, 2019, 294, 11994-12006.	1.6	15
14	Fas deficiency in mice with the Balb/c background induces blepharitis with allergic inflammation and hyper-IgE production in conjunction with severe autoimmune disease. International Immunology, 2013, 25, 287-293.	1.8	9
15	FLASH/casp8ap2 Is Indispensable for Early Embryogenesis but Dispensable for Proliferation and Differentiation of ES Cells. PLoS ONE, 2014, 9, e108032.	1.1	9