

Mehrdad Rakaee

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6740413/publications.pdf>

Version: 2024-02-01

18
papers

691
citations

566801

15
h-index

839053

18
g-index

18
all docs

18
docs citations

18
times ranked

1332
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Prognostic Value of Macrophage Phenotypes in Resectable Non-Small Cell Lung Cancer Assessed by Multiplex Immunohistochemistry. <i>Neoplasia</i> , 2019, 21, 282-293. | 2.3 | 117 |
| 2 | CTLA-4 expression in the non-small cell lung cancer patient tumor microenvironment: diverging prognostic impact in primary tumors and lymph node metastases. <i>Cancer Immunology, Immunotherapy</i> , 2017, 66, 1449-1461. | 2.0 | 69 |
| 3 | Prognostic effect of intratumoral neutrophils across histological subtypes of non-small cell lung cancer. <i>Oncotarget</i> , 2016, 7, 72184-72196. | 0.8 | 54 |
| 4 | High expression of PDGFR- β in prostate cancer stroma is independently associated with clinical and biochemical prostate cancer recurrence. <i>Scientific Reports</i> , 2017, 7, 43378. | 1.6 | 51 |
| 5 | Evaluation of tumor-infiltrating lymphocytes using routine H&E slides predicts patient survival in resected non-small cell lung cancer. <i>Human Pathology</i> , 2018, 79, 188-198. | 1.1 | 49 |
| 6 | LAG-3 in Non-Small-cell Lung Cancer: Expression in Primary Tumors and Metastatic Lymph Nodes Is Associated With Improved Survival. <i>Clinical Lung Cancer</i> , 2018, 19, 249-259.e2. | 1.1 | 48 |
| 7 | MicroRNA 141 is associated to outcome and aggressive tumor characteristics in prostate cancer. <i>Scientific Reports</i> , 2019, 9, 386. | 1.6 | 43 |
| 8 | Evaluation of the proliferation marker Ki-67 in a large prostatectomy cohort. <i>PLoS ONE</i> , 2017, 12, e0186852. | 1.1 | 39 |
| 9 | Tertiary lymphoid structure score: a promising approach to refine the TNM staging in resected non-small cell lung cancer. <i>British Journal of Cancer</i> , 2021, 124, 1680-1689. | 2.9 | 37 |
| 10 | Tissue analyses reveal a potential immune-adjuvant function of FAP-1 positive fibroblasts in non-small cell lung cancer. <i>PLoS ONE</i> , 2018, 13, e0192157. | 1.1 | 35 |
| 11 | Interplay of immune and kidney resident cells in the formation of tertiary lymphoid structures in lupus nephritis. <i>Autoimmunity Reviews</i> , 2021, 20, 102980. | 2.5 | 35 |
| 12 | Low Expression of miR-424-3p is Highly Correlated with Clinical Failure in Prostate Cancer. <i>Scientific Reports</i> , 2019, 9, 10662. | 1.6 | 32 |
| 13 | High expression of miR-17-5p in tumor epithelium is a predictor for poor prognosis for prostate cancer patients. <i>Scientific Reports</i> , 2021, 11, 13864. | 1.6 | 21 |
| 14 | Differential prognostic impact of platelet-derived growth factor receptor expression in NSCLC. <i>Scientific Reports</i> , 2019, 9, 10163. | 1.6 | 20 |
| 15 | Digitally quantified CD8+ cells: the best candidate marker for an immune cell score in non-small cell lung cancer?. <i>Carcinogenesis</i> , 2020, 41, 1671-1681. | 1.3 | 18 |
| 16 | Overexpression of miR-20a-5p in Tumor Epithelium Is an Independent Negative Prognostic Indicator in Prostate Cancer—A Multi-Institutional Study. <i>Cancers</i> , 2021, 13, 4096. | 1.7 | 11 |
| 17 | Proteomic analyses identify major vault protein as a prognostic biomarker for fatal prostate cancer. <i>Carcinogenesis</i> , 2021, 42, 685-693. | 1.3 | 10 |
| 18 | Expression of phosphatase of regenerating liver (PRL)-3, is independently associated with biochemical failure, clinical failure and death in prostate cancer. <i>PLoS ONE</i> , 2017, 12, e0189000. | 1.1 | 2 |