

# Damir A Safin

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/6740772/damir-a-safin-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

175  
papers

2,441  
citations

25  
h-index

33  
g-index

185  
ext. papers

2,805  
ext. citations

3.3  
avg, IF

5.19  
L-index

#	Paper	IF	Citations
175	Computational Analysis of Molnupiravir.. <i>International Journal of Molecular Sciences</i> , <b>2022</b> , 23,	6.3	5
174	A readily available structural analogue of integrastatins A and B: Insight into the crystal structure, Hirshfeld surface analysis and computational study. <i>Tetrahedron</i> , <b>2022</b> , 109, 132671	2.4	1
173	Spodium bonds and metal-halogen-halogen-metal interactions in propagation of monomeric units to dimeric or polymeric architectures. <i>Journal of Molecular Structure</i> , <b>2022</b> , 1252, 132144	3.4	1
172	Supramolecular aggregation of lead(II) perchlorate and a thiosemicarbazide derivative linked by a myriad of non-covalent interactions. <i>Inorganica Chimica Acta</i> , <b>2022</b> , 120974	2.7	0
171	Computational analysis of aspirin. <i>Journal of Molecular Structure</i> , <b>2021</b> , 131975	3.4	4
170	A chiral (1R,2R)-N,N'-bis-(salicylidene)-1,2-diphenyl-1,2-ethanediamine Schiff base dye: synthesis, crystal structure, Hirshfeld surface analysis, computational study, photophysical properties and in silico antifungal activity. <i>Journal of the Iranian Chemical Society</i> , <b>2021</b> , 18, 2897	2	4
169	Solvent-Induced Formation of Novel Ni(II) Complexes Derived from Bis-Thiosemicarbazone Ligand: An Insight from Experimental and Theoretical Investigations. <i>International Journal of Molecular Sciences</i> , <b>2021</b> , 22,	6.3	3
168	Spodium bonding and other non-covalent interactions assisted supramolecular aggregation in a new mercury(II) complex of a nicotinohydrazide derivative. <i>Inorganica Chimica Acta</i> , <b>2021</b> , 519, 120279	2.7	11
167	Supramolecular structures of Ni and Cu with the sterically demanding Schiff base dyes driven by cooperative action of preagostic and other non-covalent interactions. <i>IUCrJ</i> , <b>2021</b> , 8, 351-361	4.7	
166	A new coordination polymer constructed from Pb(NO <sub>3</sub> ) <sub>2</sub> and a benzylideneisonicotinohydrazide derivative: Coordination-induced generation of a hole towards a tetrel-bonding stabilized structure. <i>Journal of Molecular Structure</i> , <b>2021</b> , 1234, 130139	3.4	4
165	A novel paramagnetic coordination polymer, fabricated from Co(NCS) <sub>2</sub> and 2-pyridinecarbaldehyde isonicotinoylhydrazone. <i>Inorganica Chimica Acta</i> , <b>2021</b> , 522, 120335	2.7	0
164	Zinc(II) complexes derived from 2-formylpyridine nicotinoyl hydrazone as organic blocker: Syntheses, crystal architectures, Hirshfeld surface analyses and DFT studies. <i>Journal of Molecular Structure</i> , <b>2021</b> , 1229, 129614	3.4	1
163	Ligand structure-driven self-assembly of Zn(NCS) <sub>2</sub> with a carbohydrazone ligand: A possible intermediate towards a [2D] metallic grid. <i>Journal of Molecular Structure</i> , <b>2021</b> , 1225, 129269	3.4	2
162	A supramolecular 3D structure constructed from a new metal chelate self-assembled from Sn(NCS) <sub>2</sub> and phenyl(pyridin-2-yl)methylenepicolinohydrazide. <i>Journal of Molecular Structure</i> , <b>2021</b> , 1224, 129188	3.4	4
161	-Aminophosphonates 4-XCH-NH-CH(4-BrCH)-P(O)(OPr) (X = H, Br, MeO): Crystal structures, Hirshfeld surface analysis, computational studies and in silico molecular docking with the SARS-CoV-2 proteins. <i>Tetrahedron</i> , <b>2021</b> , 97, 132376	2.4	2
160	Evaluation of the antitumor activity of a series of the pincer-type metal complexes produced from isonicotinohydrazide derivative. <i>Journal of Inorganic Biochemistry</i> , <b>2021</b> , 223, 111525	4.2	3
159	Novel sterically demanding Schiff base dyes: An insight from experimental and theoretical calculations. <i>Journal of Luminescence</i> , <b>2021</b> , 238, 118264	3.8	3

158	Antitumor effects of novel nickelhydrazone complexes in lung cancer cells. <i>New Journal of Chemistry</i> , <b>2020</b> , 44, 9064-9072	3.6	16
157	Resonance Assisted Hydrogen Bonding Phenomenon Unveiled through Both Experiments and Theory: A New Family of Ethyl N-Salicylidene-glycinate Dyes. <i>Chemistry - A European Journal</i> , <b>2020</b> , 26, 12987-12995	4.8	11
156	Complexes of BiCl <sub>3</sub> with hydrazone derived ligands: a Möbius-like discrete metal chelate versus a salt-like porous polymeric structure. <i>New Journal of Chemistry</i> , <b>2020</b> , 44, 9429-9437	3.6	4
155	Photophysical properties of ethyl N-(5-bromosalicylidene)glycinate and ethyl N-(5-nitrosalicylidene)glycinate in CH <sub>2</sub> Cl <sub>2</sub> . <i>Journal of Luminescence</i> , <b>2020</b> , 226, 117454	3.8	6
154	Supramolecular lead(II) architectures engineered by tetrel bonds. <i>CrystEngComm</i> , <b>2020</b> , 22, 2389-2396	3.3	20
153	New metal chelate constructed from Ni(NCS) <sub>2</sub> and 1,2-diphenyl-1,2-bis((phenyl(pyridin-2-yl)methylene)hydrazono)ethane. <i>Inorganica Chimica Acta</i> , <b>2020</b> , 509, 119707	2.7	6
152	Novel lanthanide(III) complex [LaL <sub>2</sub> (NO <sub>3</sub> )(H <sub>2</sub> O) <sub>2</sub> ][5H <sub>2</sub> O] with 2-pyridine carboxaldehyde isonicotinoyl hydrazine exhibiting a 3D supramolecular topology 3,6T49. <i>Journal of Molecular Structure</i> , <b>2020</b> , 1212, 128151	3.4	9
151	Non-covalent interactions induced supramolecular architecture of Hg(NCS) <sub>2</sub> with 3-pyridinecarbaldehyde nicotinoylhydrazine. <i>Inorganica Chimica Acta</i> , <b>2020</b> , 509, 119700	2.7	5
150	On the importance of σ-hole spodium bonding in tricoordinated Hg complexes. <i>Dalton Transactions</i> , <b>2020</b> , 49, 17547-17551	4.3	18
149	Supramolecular architecture constructed from the hemidirected lead(II) complex with N'-(4-hydroxybenzylidene)isonicotinohydrazide. <i>Inorganica Chimica Acta</i> , <b>2020</b> , 502, 119350	2.7	19
148	Supramolecular architectures of Mn(NCS) <sub>2</sub> complexes with N'-(1-(pyridin-4-yl)ethylidene)picolinohydrazide and N'-(phenyl(pyridin-4-yl)methylene)isonicotinohydrazide. <i>Polyhedron</i> , <b>2020</b> , 190, 114776	2.7	6
147	Lead(ii) coordination polymers driven by pyridine-hydrazine donors: from anion-guided self-assembly to structural features. <i>Dalton Transactions</i> , <b>2020</b> , 49, 11238-11248	4.3	11
146	A new spodium bond driven coordination polymer constructed from mercury(II) azide and 1,2-bis(pyridin-2-ylmethylene)hydrazine. <i>New Journal of Chemistry</i> , <b>2020</b> , 44, 21100-21107	3.6	14
145	Tetrel Bonding and Other Non-Covalent Interactions Assisted Supramolecular Aggregation in a New Pb(II) Complex of an Isonicotinohydrazide. <i>Molecules</i> , <b>2020</b> , 25,	4.8	13
144	Effect of Solvent on the Structural Diversity of Quasi-Aromatic Möbius Cadmium(II) Complexes Fabricated from the Bulky N <sub>6</sub> Tetradentate Helical Ligand. <i>Crystal Growth and Design</i> , <b>2019</b> , 19, 1649-1659	3.5	10
143	Chameleon-like Nature of Anagostic Interactions and Its Impact on Metalloaromaticity in Square-Planar Nickel Complexes. <i>Organometallics</i> , <b>2019</b> , 38, 1973-1981	3.8	13
142	Structural versatility of the quasi-aromatic Möbius type zinc(ii)-pseudohalide complexes - experimental and theoretical investigations.. <i>RSC Advances</i> , <b>2019</b> , 9, 23764-23773	3.7	9
141	A Highly Stable All-Inorganic CsPbBr <sub>3</sub> Perovskite Solar Cell. <i>European Journal of Inorganic Chemistry</i> , <b>2019</b> , 2019, 3699-3703	2.3	19

140	Metal chelates constructed from CdHal <sub>2</sub> (Hal = Cl, Br, I) and 1,2-diphenyl-1,2-bis((phenyl(pyridin-2-yl)methylene)hydrazono)ethane. <i>Journal of Molecular Structure</i> , <b>2019</b> , 1176, 743-750	3.4	8
139	Copper(II) acetate structures with benzimidazole derivatives. <i>Inorganica Chimica Acta</i> , <b>2019</b> , 488, 238-245.7		4
138	N-Thiophosphorylthioureas RNHC(S)NHP(S)(OiPr) <sub>2</sub> as an Excellent Platform for Studying the Synergy between Hydrogen-Hydrogen Bonding and Other Families of Non-Covalent Interactions. <i>European Journal of Organic Chemistry</i> , <b>2019</b> , 2019, 493-503	3.2	12
137	M <sup>B</sup> ius-like metal chelates constructed from CdHal <sub>2</sub> (Hal = Cl, Br, I) and benzilbis(pyridin-2-yl)methylidenehydrazone. <i>Inorganica Chimica Acta</i> , <b>2019</b> , 484, 481-490	2.7	7
136	Extended lead(II) architectures engineered via tetrel bonding interactions. <i>New Journal of Chemistry</i> , <b>2018</b> , 42, 4959-4971	3.6	63
135	Quasi-aromatic M <sup>B</sup> ius Metal Chelates. <i>Inorganic Chemistry</i> , <b>2018</b> , 57, 4395-4408	5.1	29
134	London Dispersion Forces in Crystal Packing of Thiourea Derivatives. <i>Crystal Growth and Design</i> , <b>2018</b> , 18, 5385-5397	3.5	13
133	Azide-rich complexes of cobalt(III) with the rare 5-phenyl-2,2'-bipyridine ligand. <i>Inorganica Chimica Acta</i> , <b>2017</b> , 459, 63-72	2.7	2
132	Alternative Route Toward Nitrones: Experimental and Theoretical Findings. <i>Journal of Organic Chemistry</i> , <b>2017</b> , 82, 1666-1675	4.2	2
131	Stepwise crystallographic visualization of dynamic guest binding in a nanoporous framework. <i>Chemical Science</i> , <b>2017</b> , 8, 3171-3177	9.4	49
130	Polar protic solvent-trapping polymorphism of the Hg(II)-hydrazone coordination polymer: experimental and theoretical findings. <i>CrystEngComm</i> , <b>2017</b> , 19, 3017-3025	3.3	22
129	Confinement effects of a crystalline sponge on ferrocene and ferrocene carboxaldehyde. <i>Chemical Communications</i> , <b>2017</b> , 53, 5645-5648	5.8	21
128	Anion-driven tetrel bond-induced engineering of lead(II) architectures with N <sup>2</sup> -(1-(2-pyridyl)ethylidene)nicotinohydrazide: experimental and theoretical findings. <i>Inorganic Chemistry Frontiers</i> , <b>2017</b> , 4, 171-182	6.8	36
127	Complexes and salts of the nitrogen-rich triazole-tetrazole hybrid ligand with alkali and alkaline earth metal cations: experimental and theoretical findings. <i>New Journal of Chemistry</i> , <b>2017</b> , 41, 6210-6218	3.6	6
126	Single-molecule magnetism arising from cobalt(II) nodes of a crystalline sponge. <i>Journal of Materials Chemistry C</i> , <b>2017</b> , 5, 835-841	7.1	51
125	Solvent-driven azide-induced mononuclear discrete versus one-dimensional polymeric aromatic M <sup>B</sup> ius cadmium(ii) complexes of an N tetradentate helical ligand. <i>Dalton Transactions</i> , <b>2017</b> , 46, 14888-14896	4.3	29
124	Solid Aluminum Borohydrides for Prospective Hydrogen Storage. <i>ChemSusChem</i> , <b>2017</b> , 10, 4725-4734	8.3	17
123	Organic matrix-induced formation of a discrete cyclic [Cl <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> cluster. <i>New Journal of Chemistry</i> , <b>2017</b> , 41, 8263-8269	3.6	6

122	Ligand-Driven Coordination Sphere-Induced Engineering of Hybride Materials Constructed from PbCl and Bis-Pyridyl Organic Linkers for Single-Component Light-Emitting Phosphors. <i>Inorganic Chemistry</i> , <b>2017</b> , 56, 9698-9709	5.1	49
121	CBr <sub>2</sub> Br vs. CBr <sub>2</sub> Br vs. CBr <sub>2</sub> N bonding in molecular self-assembly of pyridine-containing dyes. <i>RSC Advances</i> , <b>2016</b> , 6, 53669-53678	3.7	15
120	1,2,4-Triazole-based molecular switches: crystal structures, Hirshfeld surface analysis and optical properties. <i>CrystEngComm</i> , <b>2016</b> , 18, 7284-7296	3.3	43
119	Polymorphism driven optical properties of an anil dye. <i>CrystEngComm</i> , <b>2016</b> , 18, 7249-7259	3.3	23
118	Detailed studies of the interaction of 3-chloroaniline with O,O'-diphenylphosphorylthiocyanate. <i>New Journal of Chemistry</i> , <b>2016</b> , 40, 1230-1236	3.6	18
117	Reduction of CO <sub>2</sub> with KBH <sub>4</sub> in solvent-free conditions. <i>International Journal of Hydrogen Energy</i> , <b>2016</b> , 41, 14377-14386	6.7	29
116	Ligand-Driven Anion-Interaction-Induced Silver(I) Coordination Chemistry. <i>Crystal Growth and Design</i> , <b>2016</b> , 16, 3763-3770	3.5	17
115	Hidden Transformations of a Crystalline Sponge: Elucidating the Stability of a Highly Porous Three-Dimensional Metal-Organic Framework. <i>Crystal Growth and Design</i> , <b>2016</b> , 16, 4043-4050	3.5	17
114	CdS Nanoparticles Fabricated from the Single-Source Precursor [Cd{Et <sub>2</sub> NC(S)NP(S)(OiPr) <sub>2</sub> } <sub>2</sub> ]: In Depth Experimental and Theoretical Studies. <i>Crystal Growth and Design</i> , <b>2016</b> , 16, 3287-3296	3.5	5
113	Mononuclear heteroleptic complexes of copper(I) with 5-phenyl-2,2'-bipyridine and triphenylphosphine: crystal structures, Hirshfeld surface analysis and luminescence properties. <i>New Journal of Chemistry</i> , <b>2016</b> , 40, 6156-6163	3.6	22
112	An intermolecular pyrene excimer in the pyrene-labeled N-thiophosphorylated thiourea and its nickel(II) complex. <i>Inorganic Chemistry Frontiers</i> , <b>2016</b> , 3, 1419-1431	6.8	13
111	Unprecedented trinuclear Ag(I) complex with 2,4,6-tris(2-pyrimidyl)-1,3,5-triazine as an efficient catalyst for the aziridination of olefins. <i>Chemistry - A European Journal</i> , <b>2015</b> , 21, 6144-9	4.8	36
110	Interaction of 2,4,6-tris(2-pyrimidyl)-1,3,5-triazine (TPyMT) with CoX <sub>2</sub> (X = Cl, Br) in water: trapping of new self-assembled water-chloride/bromide clusters in a [Co(bpca) <sub>2</sub> ] <sup>+</sup> host (bpca = bis(2-pyrimidylcarbonyl)amidate anion). <i>New Journal of Chemistry</i> , <b>2015</b> , 39, 7147-7152	3.6	16
109	Anion-induced Ag(I) self-assemblies with electron deficient aromatic ligands: anion-system interactions as a driving force for templated coordination networks. <i>Chemical Communications</i> , <b>2015</b> , 51, 9547-50	5.8	36
108	The renaissance of 2,4,6-tris(2-pyrimidyl)-1,3,5-triazine (TPyMT) coordination chemistry. <i>Dalton Transactions</i> , <b>2015</b> , 44, 20287-94	4.3	23
107	Luminescent mononuclear mixed ligand complexes of copper(I) with 5-phenyl-2,2'-bipyridine and triphenylphosphine. <i>Dalton Transactions</i> , <b>2015</b> , 44, 16824-32	4.3	36
106	Hydrazine selective dual signaling chemodosimetric probe in physiological conditions and its application in live cells. <i>Analytica Chimica Acta</i> , <b>2015</b> , 893, 84-90	6.6	28
105	A smart rhodamine-pyridine conjugate for bioimaging of thiocyanate in living cells. <i>RSC Advances</i> , <b>2015</b> , 5, 103350-103357	3.7	8

104	Hybrid Material Constructed from Hg(NCS) <sub>2</sub> and 2,4,6-Tris(2-pyrimidyl)-1,3,5-triazine (TPymT): Coordination of TPymT in a 2,2'-Bipyridine-Like Mode. <i>European Journal of Inorganic Chemistry</i> , <b>2015</b> , 2015, 441-446	2.3	14
103	A composite of complex and chemical hydrides yields the first Al-based amidoborane with improved hydrogen storage properties. <i>Chemistry - A European Journal</i> , <b>2015</b> , 21, 14562-70	4.8	25
102	Influence of the Homopolar Dihydrogen Bonding C-H...H-C on Coordination Geometry: Experimental and Theoretical Studies. <i>Chemistry - A European Journal</i> , <b>2015</b> , 21, 16679-87	4.8	33
101	A Neutral 1D Coordination Polymer Constructed from the Ni(II) Complex of the N-Phosphorylated Thiourea PhNHC(S)NHP(O)(OPh) <sub>2</sub> and Pyrazine: A Single-Source Precursor for Nickel Nanoparticles. <i>European Journal of Inorganic Chemistry</i> , <b>2015</b> , 2015, 1160-1166	2.3	14
100	Elucidating the elusive crystal structure of 2,4,6-tris(2-pyrimidyl)-1,3,5-triazine. <i>CrystEngComm</i> , <b>2015</b> , 17, 2190-2195	3.3	15
99	Renaissance of the coordination chemistry of 2,4,6-tris(2-pyrimidyl)-1,3,5-triazine (TPymT). Part II: new insights into the reaction of TPymT with Pb(NO <sub>3</sub> ) <sub>2</sub> . <i>CrystEngComm</i> , <b>2014</b> , 16, 3466-3469	3.3	19
98	N-Salicylidene aniline derivatives based on the N <sup>+</sup> -thiophosphorylated thiourea scaffold. <i>CrystEngComm</i> , <b>2014</b> , 16, 7053-7061	3.3	14
97	Solid-state photochromism and thermochromism of N-salicylidene pyrene derivatives. <i>CrystEngComm</i> , <b>2014</b> , 16, 8786	3.3	15
96	Photoreversible solid state negative photochromism of N-(3,5-dichlorosalicylidene)-1-aminopyrene. <i>CrystEngComm</i> , <b>2014</b> , 16, 5524	3.3	12
95	Supramolecular Coordination Complexes of the N-Thiophosphorylated 2,5-Dithiobiurea [NHC(S)NHP(S)(OiPr) <sub>2</sub> ] <sub>2</sub> with Zn(II) and Cd(II) Ions [Cation-Induced Dinuclear Mesocate Structure versus Tetranuclear Nanoscaled Aggregate. <i>European Journal of Inorganic Chemistry</i> , <b>2014</b> , 2014, 5522-5529	2.3	10
94	Novel facile single source precursor N-phosphorylated bis-thiourea [NHC(S)NHP(O)(OPh) <sub>2</sub> ] <sub>2</sub> for the $\pi$ - $\pi$ cross-link formation of s-triazolidino[1,2-a]-s-triazolidine-3,7-dithiones: First structurally characterized representative. <i>Polyhedron</i> , <b>2013</b> , 62, 133-137	2.7	5
93	Experimental and theoretical investigations of the Ni(II) complex with N-phosphorylated thiourea iPrNHC(S)NHP(O)(OPh) <sub>2</sub> . <i>CrystEngComm</i> , <b>2013</b> , 15, 7845	3.3	16
92	From a mononuclear Ni(II) precursor to antiferromagnetically coupled trinuclear double-stranded helicates. <i>Dalton Transactions</i> , <b>2013</b> , 42, 16470-3	4.3	10
91	Crown ether-containing Schiff base as a highly efficient "turn-on" fluorescent sensor for determination and separation of Zn <sup>2+</sup> in water. <i>Dalton Transactions</i> , <b>2013</b> , 42, 1969-72	4.3	27
90	Crucial Influence of the Intramolecular Hydrogen Bond on the Coordination Mode of RC(S)NHP(S)(OiPr) <sub>2</sub> in Homoleptic Complexes with Ni(II). <i>European Journal of Inorganic Chemistry</i> , <b>2013</b> , 2013, 545-555	2.3	17
89	Complexation properties of N-thiophosphorylated thiourea 2-PyNHC(S)NHP(S)(OiPr) <sub>2</sub> towards Ni(II). <i>Dalton Transactions</i> , <b>2013</b> , 42, 5252-7	4.3	9
88	First evidence of thermo- and two-step photochromism of tris-anils. <i>RSC Advances</i> , <b>2013</b> , 3, 6466	3.7	22
87	Routes to novel mono- and bis-tetrazole compounds: synthesis, spectroscopic and structural characterization. <i>New Journal of Chemistry</i> , <b>2013</b> , 37, 1174	3.6	4



86	First structurally characterized self-assembly of bipodal N-thiophosphorylated bis-thiourea with Co(II): magnetic properties and thermal decomposition. <i>Dalton Transactions</i> , <b>2013</b> , 42, 5532-8	4.3	6
85	First paramagnetic Pd(II) complex with a PdN <sub>4</sub> S <sub>2</sub> coordination core. <i>Dalton Transactions</i> , <b>2013</b> , 42, 902-54.3	4.3	5
84	Renaissance of the coordination chemistry of 2,4,6-tris(2-pyrimidyl)-1,3,5-triazine (TPymT). Part I: First crystal structure of a TPymT complex with a d-metal cation. <i>CrystEngComm</i> , <b>2013</b> , 15, 10419	3.3	24
83	New mononuclear Cu(II) complexes and 1D chains with 4-amino-4H-1,2,4-triazole. <i>International Journal of Molecular Sciences</i> , <b>2013</b> , 14, 23597-613	6.3	9
82	Decomposition of the mixed-ligand silver(I) complex [Ag(PPh <sub>3</sub> ) <sub>2</sub> {Et <sub>2</sub> NC(S)NP(S)(OiPr) <sub>2</sub> }] with the formation of [Ag(PPh <sub>3</sub> ) <sub>2</sub> {Et <sub>2</sub> NC(S)NP(S)(OiPr) <sub>2</sub> }], [Ag(PPh <sub>3</sub> ) <sub>3</sub> NCS][C(CH <sub>3</sub> ) <sub>2</sub> C=O] and Et <sub>2</sub> NP(S)(OiPr) <sub>2</sub> . <i>Inorganic Chemistry Communication</i> , <b>2012</b> , 15, 117-120	3.1	1
81	Influence of polymorphism on N-thiophosphorylated thiourea 4-Me <sub>2</sub> NC <sub>6</sub> H <sub>4</sub> NHC(S)NHP(S)(OiPr) <sub>2</sub> crystal design. <i>Inorganic Chemistry Communication</i> , <b>2012</b> , 18, 34-37	3.1	2
80	Synthesis of tetraphenylphosphonium (thio)phosphorylthio benzamides and hydrolytic degradation to Ph <sub>3</sub> P=O, H <sub>2</sub> S and phosphoryl(thio)benzamides: A route for crystal design of PhC(X)NHP(O)(OiPr) <sub>2</sub> (X=S, O). <i>Inorganic Chemistry Communication</i> , <b>2012</b> , 20, 162-166	3.1	
79	Heteroleptic Cu(I) and Ag(I) complexes of N-thiophosphorylated tris(thioureas and triphenylphosphane. <i>Polyhedron</i> , <b>2012</b> , 33, 114-118	2.7	9
78	Solid-state thermo- and photochromism in N,N'-bis(5-X-salicylidene)diamines (X = H, Br). <i>RSC Advances</i> , <b>2012</b> , 2, 11379	3.7	34
77	Halogen anion-induced formation of [(PdLX) <sub>2</sub> ] (X = Cl-, Br-, I-) vs. [PdL <sub>2</sub> ] (L = [6-MeO(O)CC <sub>6</sub> H <sub>4</sub> NHC(S)NP(S)(OiPr) <sub>2</sub> ]-): reversible photoinduced cis/trans isomerization of [PdL <sub>2</sub> ]. <i>Dalton Transactions</i> , <b>2012</b> , 41, 4324-7	4.3	5
76	N-(diisopropylthiophosphoryl)-N'-(R)-thioureas: synthesis, characterization, crystal structures and competitive bulk liquid membrane transport of some metal ions. <i>Dalton Transactions</i> , <b>2012</b> , 41, 3223-32	4.3	20
75	Influence of CH <sub>2</sub> Cl <sub>2</sub> for the structure stabilization of the Nill complex [Ni{6-MeO(O)CC <sub>6</sub> H <sub>4</sub> NHC(S)NP(S)(OiPr) <sub>2</sub> -1,5-S,S'} <sub>2</sub> ][CH <sub>2</sub> Cl <sub>2</sub> . <i>CrystEngComm</i> , <b>2012</b> , 14, 370-373	3.3	10
74	Spin-canted ferromagnetic behaviour and unusual thermal decomposition of [Co{MeO(O)CC <sub>6</sub> H <sub>4</sub> NHC(S)NP(S)(OiPr) <sub>2</sub> }] <sub>2</sub> . <i>New Journal of Chemistry</i> , <b>2012</b> , 36, 2642	3.6	3
73	Influence of the coordination mode in [Ni{RC(S)NP(S)(OiPr) <sub>2</sub> }] <sub>2</sub> for the formation of nickel-containing nanoparticles. <i>Dalton Transactions</i> , <b>2012</b> , 41, 2234-6	4.3	18
72	Thiophosphorylated bis-thioureas for competitive bulk liquid membrane transport of some metal ions. <i>CrystEngComm</i> , <b>2012</b> , 14, 1324-1329	3.3	15
71	A metal-organic framework made of an asymmetric 1,2,4-triazole and tetrazole ligand. <i>CrystEngComm</i> , <b>2012</b> , 14, 8153	3.3	13
70	Crown ether-containing N-salicylidene aniline derivatives: synthesis, characterization and optical properties. <i>CrystEngComm</i> , <b>2012</b> , 14, 5523	3.3	24
69	Crystal Structure and Interaction of 6-Amino Coumarin with Nitrite Ion for Its Selective Fluorescence Detection. <i>Spectroscopy Letters</i> , <b>2012</b> , 45, 225-235	1.1	11

68	Homoleptic polynuclear CuI and AgI complexes of N-thiophosphorylated thioureas o-RO(O)CC6H4NHC(S)NHP(S)(OiPr)2 (R = Me, Et). <i>CrystEngComm</i> , <b>2012</b> , 14, 774-778	3-3	5
67	Reaction of aminobenzoate esters with N,N'-dimethylformamide azine dihydrochloride: crucial influence of intramolecular hydrogen bonding for the formation of 1,2,4-triazoles. <i>CrystEngComm</i> , <b>2012</b> , 14, 4812	3-3	4
66	Complexation properties of the crown ether-containing N-thiophosphorylated thiourea towards Ni(II). <i>Dalton Transactions</i> , <b>2012</b> , 41, 1451-3	4-3	13
65	Heteroligand copper(I) complexes of N-thiophosphorylated thioureas and phosphanes: Versatile structures and luminescence. <i>Inorganica Chimica Acta</i> , <b>2011</b> , 370, 59-64	2-7	8
64	Highly efficient [Ni{iPrHNC(S)NP(S)(OiPr)2-1,3-N,S'}2]/PR3 (R3 = Me3, Me2Ph) complexes for the generation of Ni(0) for catalysis. <i>Dalton Transactions</i> , <b>2011</b> , 40, 4806-9	4-3	14
63	Solvent-induced 1,3-N,S- vs. 1,5-S,S'-coordination in the NiII complex [Ni{p-Me2NC6H4NHC(S)NP(S)(OiPr)2}2]. <i>CrystEngComm</i> , <b>2011</b> , 13, 5321	3-3	22
62	Intramolecular hydrogen bonding controls 1,3-N,S- vs. 1,5-S,S'-coordination in Ni(II) complexes of N-thiophosphorylated thioureas RNHC(S)NHP(S)(OiPr)2. <i>Dalton Transactions</i> , <b>2011</b> , 40, 3142-53	4-3	30
61	Versatile structures and photophysical properties of poly- and mononuclear CuI complexes with N-thiophosphorylated thioureas RNHC(S)NHP(S)(OiPr)2 and phosphanes. <i>CrystEngComm</i> , <b>2011</b> , 13, 568-576	3-3	13
60	Synthesis and characterization of [H2NC(S)NP(S)(OiPr)2]n complexes of Co(II), Ni(II), Zn(II) and Cd(II). <i>Inorganica Chimica Acta</i> , <b>2011</b> , 365, 32-37	2-7	15
59	Studies on Zn(II) and Cd(II) heteroligand complexes with RC(S)NHP(O)(OiPr)2 (R=Ph, NH2, PhNH) and diimine (bpy and phen) ligands. C-C bond cleavage of CH2Cl2 by [Zn(phen){PhC(S)NP(O)(OiPr)2}2]. <i>Inorganica Chimica Acta</i> , <b>2011</b> , 366, 19-26	2-7	9
58	Zn(II), Co(II) and Ni(II) complexes of a phosphorylthiourea derivative of 4-[(EtO)2P(O)CH2]-C6H4-NHC(S)NHP(O)(OiPr)2. <i>Collection of Czechoslovak Chemical Communications</i> , <b>2010</b> , 75, 507-515		1
57	Copper(I) complexes with N-thiophosphorylated thioureas and phosphines with versatile structures and luminescence. <i>CrystEngComm</i> , <b>2010</b> , 12, 134-143	3-3	13
56	Nitrocellulose Membrane, Modified by RC(S)NHP(X)(OiPr)2 (X = S, R = PhNH; X = O, R = PhNH, Ph), for Sorption Extraction of Cobalt Cations. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , <b>2010</b> , 185, 1815-1822	1	1
55	Complexes of [(iPrO)2P(O)NHC(S)NHCH2]2CH2 with Co(II), Ni(II), Zn(II), and Pd(II): Reaction of [(iPrO)2P(O)NHC(S)NHCH2]2 with KOH Leading to the Imidazolidine-2-Imine Derivative. <i>Phosphorus, Sulfur and Silicon and the Related Elements</i> , <b>2010</b> , 185, 1437-1444	1	7
54	Aerial oxidation of tetrahydrofuran to 2-hydroxotetrahydrofuran in the presence of a trimeric CuI complex [Cu3L3] (HL = tBuNHC(S)NHP(S)(OiPr)2) and trapping of the unstable product at recrystallization. <i>New Journal of Chemistry</i> , <b>2010</b> , 34, 2835	3-6	9
53	Tetraphenylphosphonium salts of N-thiophosphorylated thiobenzamide, mono-thioureas or bis-thioureas. <i>Heteroatom Chemistry</i> , <b>2010</b> , 21, 343-350	1-2	5
52	Synthesis and characterization of silver(I) complexes with N-thiophosphorylated thiobenzamide or thiourea and phosphines. <i>Heteroatom Chemistry</i> , <b>2010</b> , 21, 386-391	1-2	7
51	Synthesis and characterization of crown ether-containing N-phosphorylated thioureas RC(S)NHP(O)(OiPr)2 (R = 1-aza-15-crown-5, 1-aza-12-crown-4) and their Co(II) and Ni(II) complexes. <i>Heteroatom Chemistry</i> , <b>2010</b> , 21, 486-491	1-2	3



50	Synthesis, Characterisation and Luminescent Properties of Mixed-Ligand Copper(I) Complexes Incorporating N-Thiophosphorylated Thioureas and Phosphane Ligands. <i>European Journal of Inorganic Chemistry</i> , <b>2010</b> , 2010, 4018-4026	2.3	29
49	Synthesis and characterization of [Me <sub>2</sub> NC(S)NP(S)(OiPr) <sub>2</sub> ] <sub>n</sub> complexes of Co(II), Ni(II), Zn(II) and Cd(II). <i>Polyhedron</i> , <b>2010</b> , 29, 488-492	2.7	17
48	Synthesis and characterization of N-thiophosphorylated thioureas RC(S)NHP(S)(OiPr) <sub>2</sub> (p-(4-imidazol-1-yl)C <sub>6</sub> H <sub>4</sub> NH, p-NC <sub>6</sub> H <sub>4</sub> NH, (CH <sub>2</sub> CH=C <sub>2</sub> H <sub>2</sub> ) <sub>2</sub> N): Unexpected formation of o-HO-C <sub>6</sub> H <sub>4</sub> NHC(S)NHP(S)(OiPr) <sub>2</sub> from p-NC <sub>6</sub> H <sub>4</sub> NHC(S)NHP(S)(OiPr) <sub>2</sub> . <i>Polyhedron</i> , <b>2010</b> , 29, 715-718	2.7	2
47	Synthesis and characterization of Zn(II), Cd(II) and Ni(II) complexes with N-thiophosphorylated thioureas RC(S)NP(S)(OiPr) <sub>2</sub> (R=MeNH, 2-MeC <sub>6</sub> H <sub>4</sub> NH, 2,6-Me <sub>2</sub> C <sub>6</sub> H <sub>3</sub> NH, 2,4,6-Me <sub>3</sub> C <sub>6</sub> H <sub>2</sub> NH, 2-Py(6-Me)NH). <i>Polyhedron</i> , <b>2010</b> , 29, 1515-1519	2.7	11
46	Solvent-dependent formation of two different Cd(II) complexes with p-BrC <sub>6</sub> H <sub>4</sub> C(S)NHP(O)(OiPr) <sub>2</sub> (HL). Crystallographic modification of Cd(HL) <sub>2</sub> L <sub>2</sub> . <i>Polyhedron</i> , <b>2010</b> , 29, 1837-1841	2.7	6
45	Synthesis, characterization and luminescent properties of heteroligand copper(I) complexes with N-thiophosphorylated thioureas RNHC(S)NHP(S)(OiPr) <sub>2</sub> (R=iPr, tBu, Ph, 2,6-Me <sub>2</sub> C <sub>6</sub> H <sub>3</sub> , 2,4,6-Me <sub>3</sub> C <sub>6</sub> H <sub>2</sub> ) and phosphines (PPh <sub>3</sub> , Ph <sub>2</sub> P(C <sub>5</sub> H <sub>4</sub> FeC <sub>5</sub> H <sub>4</sub> )PPh <sub>2</sub> ). <i>Inorganica Chimica Acta</i> , <b>2010</b> , 343, 1887-1894	2.7	14
44	Synthesis, characterization and complexation properties of N-phosphorylated thioureas RNHC(S)NHP(O)(OiPr) <sub>2</sub> (R=2-Py, 3-Py) towards Ni(II). <i>Inorganica Chimica Acta</i> , <b>2010</b> , 363, 1791-1795	2.7	9
43	Synthesis and Characterization of Co(II) Complexes of N-Thiophosphorylated Thioureas RC(S)NHP(S)(OiPr) <sub>2</sub> (R = Me <sub>2</sub> N, 2-MeC <sub>6</sub> H <sub>4</sub> NH, 2,6-Me <sub>2</sub> C <sub>6</sub> H <sub>3</sub> NH, 2,4,6-Me <sub>3</sub> C <sub>6</sub> H <sub>2</sub> NH). <i>Acta Chimica Slovenica</i> , <b>2010</b> , 57, 475-9	1.9	3
42	4-Bromo-N-(diisopropoxyphosphor-yl)benzamide. <i>Acta Crystallographica Section E: Structure Reports Online</i> , <b>2009</b> , 65, o2926		2
41	Suzuki Cross-Coupling Reaction Catalyzed by the Palladium Complex Pd[N-MorphC(S)NP(O)(OiPr) <sub>2</sub> -O,S] <sub>2</sub> . <i>Catalysis Letters</i> , <b>2009</b> , 129, 363-366	2.8	9
40	Pd[tBuNH(S)NHP(O)(OiPr) <sub>2</sub> -S] <sub>2</sub> Cl <sub>2</sub> Complex as Air- and Moisture-Stable Catalyst for Palladium Catalyzed Suzuki Cross-Coupling Reaction. <i>Catalysis Letters</i> , <b>2009</b> , 130, 679-682	2.8	10
39	Complexes of N-(thio)phosphorylthioureas AdNHC(S)NHP(X)(OiPr) <sub>2</sub> (X = O, S) with cobalt(II), zinc(II), and cadmium(II). Crystal structure of Co[AdNHC(S)NP(S)(OiPr) <sub>2</sub> ] <sub>2</sub> . <i>Transition Metal Chemistry</i> , <b>2009</b> , 34, 43-49	2.1	8
38	Complexes of N-Thiophosphorylthiourea NaphthylNHC(S)NHP(S)(OiPr) <sub>2</sub> (HL) with Copper(I). Crystal Structures of HL and Cu(PPh <sub>3</sub> ) <sub>2</sub> L. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , <b>2009</b> , 635, 554-557	1.3	17
37	Complexes of N-thiophosphorylthioureas RNHC(S)NHP(S)(OiPr) <sub>2</sub> (HL) (R=pyridin-2-yl, pyridin-3-yl, 6-amino-pyridin-2-yl) with copper(I): Crystal structures of Cu(PPh <sub>3</sub> ) <sub>n</sub> L (n=1, 2). <i>Inorganica Chimica Acta</i> , <b>2009</b> , 362, 1895-1900	2.7	11
36	The influence of the substituent [PhNHNH and EtN(NH <sub>2</sub> )] on the N-thiophosphorylated thiosemicarbazides RC(S)NHP(S)(OiPr) <sub>2</sub> crystal design. <i>Polyhedron</i> , <b>2009</b> , 28, 501-504	2.7	8
35	The influence of the spacer Z on N-phosphorylated bis-thioureas and 2,5-dithiobiurea Z[C(S)NHP(O)(OiPr) <sub>2</sub> ] <sub>2</sub> (Z = NHCH <sub>2</sub> CH <sub>2</sub> NH, NHC <sub>6</sub> H <sub>4</sub> -2-NH, NHNH) crystal design. <i>Polyhedron</i> , <b>2009</b> , 28, 1403-1408	2.7	13
34	Versatile coordination of N-(diisopropoxyphosphoryl)-p-bromothiobenzamide towards Zn(II) and Cd(II). <i>Polyhedron</i> , <b>2009</b> , 28, 1504-1510	2.7	22
33	Synthesis of N-(thio)phosphorylated thiosemicarbazides RC(S)NHP(X)(OiPr) <sub>2</sub> (X = S, R = NH <sub>2</sub> N(Me)) [X = O, R = NH <sub>2</sub> N(Me)] PhNHNH Reaction of NH <sub>2</sub> N(Me)C(S)NHP(S)(OiPr) <sub>2</sub> with acetone. <i>Polyhedron</i> , <b>2009</b> , 28, 2693-2697	2.7	9

- 32 Novel bicyclic hexanuclear copper(I) aggregate: Structure and solid state  $^{31}\text{P}$  CPMAS NMR spectra of  $[(\text{Cu}_3\text{L}_3)_2]$  and  $[\text{Cu}(\text{PPh}_3)_2\text{L}]$  complexes of N-(diisopropoxythiophosphinyl)-N'-phenylthiourea (HL). *Journal of Organometallic Chemistry*, **2009**, 694, 167-172 2.3 29
- 31 Intramolecular hydrogen bond controlled coordination of N-thiophosphorylated thiourea 2,6-Me<sub>2</sub>C<sub>6</sub>H<sub>3</sub>NHC(S)NHP(S)(OiPr)<sub>2</sub> with Ni(II). *Inorganic Chemistry Communication*, **2009**, 12, 678-681 3.1 25
- 30 C-Cl bond cleavage of CH<sub>2</sub>Cl<sub>2</sub> by the zinc(II) complex  $[\text{Zn}(\text{phen})\text{L}_2]$  (L=PhC(S)NP(O)(OiPr)<sub>2</sub>-) with the formation of  $[\text{Zn}(\text{phen})\text{LCl}]$  and CH<sub>2</sub>[PhC(S)NP(O)(OiPr)<sub>2</sub>]<sub>2</sub>. *Inorganic Chemistry Communication*, **2009**, 12, 913-915 3.1 11
- 29 Nanoparticles and Thin Films of Silver from Complexes of Derivatives of N-(Diisopropylthiophosphoryl)thioureas. *Chemistry of Materials*, **2009**, 21, 4233-4240 9.6 18
- 28 The influence of an intramolecular hydrogen bond on the 1,3-N,S-coordination of crown ether-containing N-phosphorylthiourea with Ni(II). *New Journal of Chemistry*, **2009**, 33, 2443 3.6 18
- 27 Competitive bulk liquid membrane transport and solvent extraction of some metal ions using RC(S)NHP(X)(OiPr)<sub>2</sub> (X = O, S) as ionophores. Formation of the polynuclear complex of  $[\text{Ag}(\text{N}[\text{triple bond}] \text{C}-\text{NP}(\text{S})(\text{OiPr})_2)]_n$ . *Dalton Transactions*, **2009**, 8227-36 4.3 26
- 26 Complexes of podand-containing bis(dithiophosphonate) ligands with cobalt(II), nickel(II) and cadmium(II): recognition of CH<sub>2</sub>Cl<sub>2</sub>. *Transition Metal Chemistry*, **2008**, 33, 921-924 2.1 13
- 25 Coordination Mode of the Nickel(II) Cation with N-Diisopropoxyphosphinyl-p-bromothiobenzamide. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **2008**, 634, 835-838 1.3 10
- 24 Complexes of N-Thiophosphorylthiourea tBuNHC(S)NHP(S)(OiPr)<sub>2</sub> with Zinc(II), Cadmium(II), Nickel(II), and Cobalt(II) Cations. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **2008**, 634, 967-971 1.3 26
- 23 1,10-Diaza-18-crown-ether, Modified by Phosphonate Pendant Arms: Synthesis, Structure, and Picrate Extraction Properties. *Zeitschrift Fur Anorganische Und Allgemeine Chemie*, **2008**, 634, 1836-1838 1.3 6
- 22 Studies on zinc(II) complexes with N-thioacylamidophosphates: X-ray crystal structure of the  $[\text{Zn}(\text{RC}(\text{S})\text{NP}(\text{O})(\text{OiPr})_2)_2]$  (R=NH<sub>2</sub>, tBuNH, c-C<sub>6</sub>H<sub>11</sub>NH). *Polyhedron*, **2008**, 27, 2022-2028 2.7 21
- 21 Monodentate S-vs. bidentate 1,5-O,S-coordination of N-phosphoryl-N'-(R)-thioureas with Pd(II). *Polyhedron*, **2008**, 27, 1995-1998 2.7 6
- 20 Nickel(II) complexes with N-(thio)phosphorylthioureas AdNHC(S)NHP(X)(OiPr)<sub>2</sub>: Versatile coordination of phosphoryl (X=O) and thiophosphoryl (X=S) derivatives. *Polyhedron*, **2008**, 27, 2271-2276 2.7 26
- 19 Complexes of N-thiophosphorylthiourea (EtO)P(O)CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>-4-[NHC(S)NHP(S)(OiPr)<sub>2</sub>] with Zn(II), Cd(II), Co(II) and Cu(PPh<sub>3</sub>)(I). *Polyhedron*, **2008**, 27, 2978-2982 2.7 13
- 18 New bifunctional N-thiophosphorylated thiourea and 2,5-dithiobiurea derivatives. Crystal structures of  $\text{R}[\text{C}(\text{S})\text{NHP}(\text{S})(\text{OiPr})_2]_2$  (R =  $\text{N}(\text{Ph})\text{CH}_2\text{CH}_2\text{N}(\text{Ph})$  and  $\text{N}(\text{H})\text{NH}$ ). *Polyhedron*, **2008**, 27, 3141-3145 2.7 12
- 17 Intramolecular hydrogen bond controlled monodentate S-coordination of N-phosphoryl-N'-(R)-thioureas with Pd(II). *Inorganic Chemistry Communication*, **2008**, 11, 330-333 3.1 11
- 16 The influence of the intramolecular hydrogen bond on the 1,3-N,S- and 1,5-O,S-coordination of N-phosphoryl-N'-(R)-thioureas with Ni(II) and Pd(II). *New Journal of Chemistry*, **2007**, 31, 1661 3.6 36
- 15 Complexes of a novel N-(diisopropoxythiophosphoryl)thiourea derivative of 1,4,8,11-tetraazacyclotetradecane with Na<sup>+</sup>, K<sup>+</sup> and cations. *Polyhedron*, **2007**, 26, 1113-1116 2.7 29

14	A Sodium trap based on benzo-15-crown-5 with an exocyclic N-(thiophosphoryl)thiourea moiety. <i>Polyhedron</i> , <b>2007</b> , 26, 1550-1560	2.7	23
13	Cobalt(II) Complexes with N-Thioacylamidophosphates and 2,2'-Bipyridyl and 1,10-Phenanthroline. Crystal Structures, Solution <sup>1</sup> H NMR Spectral and Magnetic Properties. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , <b>2007</b> , 633, 1472-1479	1.3	19
12	Rich Magnetic Chemistry of Cobalt(II) Complexes with N-Phosphorylthioureas. Crystal Structure and Solution <sup>1</sup> H NMR Spectral Properties. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , <b>2007</b> , 633, 2089-2096	1.3	9
11	Coordination Mode of the Zinc(II) and Cadmium(II) Cations with N-(Diisopropoxyphosphoryl)thiobenzamide. <i>European Journal of Inorganic Chemistry</i> , <b>2006</b> , 2006, 2027-2034	2.3	26
10	Complexes of crown-containing N-thioacylamido(thio)phosphates with Zn(II) and Co(II) cations. <i>Inorganic Chemistry Communication</i> , <b>2006</b> , 9, 1133-1135	3.1	11
9	Combined EPR and DFT study of the copper(II) complexes with N-phosphoryl thioureas. <i>Chemical Physics</i> , <b>2006</b> , 320, 59-74	2.3	17
8	Studies on cobalt(II) complexes with N-thioacylamido(thio)phosphates: X-ray crystal structure of the Co[PhC(S)NP(S)(OPri) <sub>2</sub> ] <sub>2</sub> . <i>Polyhedron</i> , <b>2006</b> , 25, 3330-3336	2.7	18
7	Synthesis, structure and properties of novel N-thiophosphorylated derivatives of 1,3-dihydro-2H-benzimidazol-2-imine and imidazolidine-2-imine. <i>Polyhedron</i> , <b>2006</b> , 25, 3611-3616	2.7	7
6	Reaction of N-diisopropoxyphosphorylthiobenzamide (HL) with the Co(II) cation: the crystal structure of CoL <sub>2</sub> and CoL <sub>2</sub> ·2HL complexes. <i>Mendeleev Communications</i> , <b>2004</b> , 14, 51-52	1.9	20
5	Naphthalene-based bis-N-salicylidene aniline dyes: Crystal structures, Hirshfeld surface analysis, computational study and molecular docking with the SARS-CoV-2 proteins. <i>Journal of the Iranian Chemical Society</i> , 1	2	3
4	Lead(II) supramolecular structures formed through a cooperative influence of the hydrazinecarbothioamide derived and ancillary ligands. <i>CrystEngComm</i> ,	3.3	1
3	Favipiravir: insight into the crystal structure, Hirshfeld surface analysis and computational study. <i>Journal of the Iranian Chemical Society</i> , 1	2	7
2	Ambroxol: Insight into the Crystal Structure, Hirshfeld Surface Analysis and Computational Study. <i>Polycyclic Aromatic Compounds</i> , 1-19	1.3	1
1	6-Amino-2-(4-fluorophenyl)-4-(trifluoromethyl)quinoline: Insight into the Crystal Structure, Hirshfeld Surface Analysis and Computational Study. <i>Polycyclic Aromatic Compounds</i> , 1-18	1.3	1