

Songee Jung

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6743693/songee-jung-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

30
papers

246
citations

10
h-index

15
g-index

30
ext. papers

314
ext. citations

3.2
avg, IF

2.76
L-index

#	Paper	IF	Citations
30	The influence of gut microbiome on progression of overactive bladder symptoms: a community-based 3-year longitudinal study in Aomori, Japan. <i>International Urology and Nephrology</i> , 2021 , 1	2.3	3
29	Application of state-space model with skew-t measurement noise to blood test value prediction. <i>Applied Mathematical Modelling</i> , 2021 , 100, 365-378	4.5	
28	Older Korean adults have lower physical function despite longer exercise times compared to their Japanese counterparts: A Japan-Korea comparative study. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2020 , 69, 39-39	0.1	
27	Pain characteristics and incidence of functional disability among community-dwelling older adults. <i>PLoS ONE</i> , 2019 , 14, e0215467	3.7	14
26	Impact of Sedentary Time on Chronic Kidney Disease and Disability Incidence in Community-Dwelling Japanese Older Adults: A 4-Year Prospective Cohort Study. <i>Journal of Aging and Physical Activity</i> , 2019 , 27, 184-190	1.6	1
25	Daily Physical Activity and Functional Disability Incidence in Community-Dwelling Older Adults with Chronic Pain: A Prospective Cohort Study. <i>Pain Medicine</i> , 2019 , 20, 1702-1710	2.8	8
24	The effect of a multicomponent intervention to promote community activity on cognitive function in older adults with mild cognitive impairment: A randomized controlled trial. <i>Complementary Therapies in Medicine</i> , 2019 , 42, 164-169	3.5	36
23	Relationship between chronic kidney disease without diabetes mellitus and components of frailty in community-dwelling Japanese older adults. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 286-292	2.9	6
22	Relationship between physical activity levels and depressive symptoms in community-dwelling older Japanese adults. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 421-427	2.9	20
21	Effectiveness of the KENKOJISEICHI local revitalization system on cognitive function change in older adults with mild cognitive impairment: study protocol for a randomized controlled trial. <i>Trials</i> , 2018 , 19, 276	2.8	8
20	Older Korean adults have lower physical function despite longer exercise times compared to their Japanese counterparts: A Japan-Korea comparative study. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 576-583	2.9	
19	The role of social frailty in explaining the association between hearing problems and mild cognitive impairment in older adults. <i>Archives of Gerontology and Geriatrics</i> , 2018 , 78, 45-50	4	10
18	Does attending an exercise class with a spouse improve long-term exercise adherence among people aged 65 years and older: a 6-month prospective follow-up study. <i>BMC Geriatrics</i> , 2017 , 17, 170	4.1	7
17	Influence of sharing exercise classes with spouses on marital relationship satisfaction in older adults. <i>Taiikugaku Kenkyu (Japan Journal of Physical Education Health and Sport Sciences)</i> , 2017 , 62, 71-81	0.1	1
16	Obesity and Muscle Weakness as Risk Factors for Mobility Limitation in Community-Dwelling Older Japanese Women: A Two-Year Follow-Up Investigation. <i>Journal of Nutrition, Health and Aging</i> , 2016 , 20, 28-34	5.2	13
15	Walking can be more effective than balance training in fall prevention among community-dwelling older adults. <i>Geriatrics and Gerontology International</i> , 2016 , 16, 118-25	2.9	30
14	Association between habitual light-intensity physical activity and lower-extremity performance: a cross-sectional study of community-dwelling older Japanese adults. <i>Geriatrics and Gerontology International</i> , 2015 , 15, 268-75	2.9	16

13	Longitudinal association between habitual walking and fall occurrences among community-dwelling older adults: analyzing the different risks of falling. <i>Archives of Gerontology and Geriatrics</i> , 2015 , 60, 45-51	4	16
12	Effects of exercise for older married couples on exercise adherence and physical fitness. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2015 , 64, 407-418	0.1	1
11	Translating physical activity reference value for older adults to the number of steps per day. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2015 , 64, 243-250	0.1	1
10	Is habitual sedentary behaviour time associated with lower extremity performance independent of moderate- to vigorous-intensity physical activity in older adults? [Cross-sectional analysis using uniaxial accelerometer] <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2014 , 63, 169-176	0.1	
9	Physical performance measures as a useful indicator of multiple geriatric syndromes in women aged 75 years and older. <i>Geriatrics and Gerontology International</i> , 2013 , 13, 901-10	2.9	12
8	Cross-sectional association between body mass index and muscle strength, and mobility limitation in community-dwelling older women. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2013 , 62, 323-330	0.1	1
7	Comparison of a combination of upper extremity performance measures and usual gait speed alone for discriminating upper extremity functional limitation and disability in older women. <i>Archives of Gerontology and Geriatrics</i> , 2012 , 55, 486-91	4	9
6	Is a composite score of physical performance measures more useful than usual gait speed alone in assessing functional status?. <i>Archives of Gerontology and Geriatrics</i> , 2012 , 55, 392-8	4	14
5	Cross-sectional analysis of hierarchy of higher-level functional capacity and quantity/intensity of physical activity in older women. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2012 , 61, 327-334	0.1	2
4	DISCRIMINATION OF MOBILITY LIMITATION BY HAND-GRIP STRENGTH AMONG COMMUNITY-DWELLING OLDER ADULTS. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2011 , 60, 259-268	0.1	6
3	ASSOCIATION BETWEEN HABITUAL WALKING AND MULTIPLE OR INJURIOUS FALLS AMONG COMMUNITY-DWELLING OLDER ADULTS ^ ^sim; DIFFERENCE BY RISK OF FALLING ^ ^sim;. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2011 , 60, 239-248	0.1	4
2	IDENTIFICATION OF THE PHYSICAL FUNCTION OF FRAIL OLDER ADULTS AND EFFECTIVITY OF THE HEALTH CHECK-UP QUESTIONNAIRE (KIHON CHECK-LIST). <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2011 , 60, 413-422	0.1	4
1	FALL-RELATED FACTORS TO TARGET IN COMMUNITY-BASED INTERVENTIONS FOR PREVENTION OF FALLS. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2010 , 59, 415-426	0.1	3