
## Shun'ichi Hisamatsu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/674784/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Estimation of dietary 14C dose coefficient using 13C-labelled compound administration analysis. Scientific Reports, 2020, 10, 8156.	3.3	3

 $_{2}$  Short-term metabolism of biologically incorporated 1251 ingested by olive flounder (Paralichthys) Tj ETQq0 0 0 rgBT  $_{1.7}^{1/2}$  Verlock 10 Tf 50 7

3	Effects of organic amendments on the natural attenuation of radiocesium transferability in grassland soils with high potassium fertility. Journal of Environmental Radioactivity, 2020, 217, 106207.	1.7	7
4	240Pu/239Pu and 242Pu/239Pu atom ratios of Japanese monthly atmospheric deposition samples during 1963–1966. Scientific Reports, 2019, 9, 8105.	3.3	6
5	A SIMULATION STUDY OF DEPOSITION PARAMETERS FOR 129I DISCHARGED FROM THE ROKKASHO REPROCESSING PLANT. Radiation Protection Dosimetry, 2019, 184, 376-379.	0.8	3
6	Concentrations of iodine-129 in livestock, agricultural, and fishery products around spent nuclear fuel reprocessing plant in Rokkasho, Japan, during and after its test operation. Environmental Monitoring and Assessment, 2019, 191, 61.	2.7	11
7	Soil-soil solution distribution coefficient of soil organic matter is a key factor for that of radioiodide in surface and subsurface soils. Journal of Environmental Radioactivity, 2017, 169-170, 131-136.	1.7	20
8	Regional and global contributions of anthropogenic iodine-129 in monthly deposition samples collected in North East Japan between 2006 and 2015. Journal of Environmental Radioactivity, 2017, 171, 65-73.	1.7	24
9	Biokinetics of13C in the human body after oral administration of13C-labeled glucose as an index for the biokinetics of14C. Journal of Radiological Protection, 2016, 36, 532-546.	1.1	2
10	Temporal variation of post-accident atmospheric 137 Cs in an evacuated area of Fukushima Prefecture: Size-dependent behaviors of 137 Cs-bearing particles. Journal of Environmental Radioactivity, 2016, 165, 131-139.	1.7	19
11	Determination of Iodide, Iodate and Total Iodine in Natural Water Samples by HPLC with Amperometric and Spectrophotometric Detection, and Off-line UV Irradiation. Analytical Sciences, 2016, 32, 839-845.	1.6	23
12	Spatial and temporal changes of 137Cs concentrations derived from nuclear power plant accident in river waters in eastern Fukushima, Japan during 2012–2014. Journal of Radioanalytical and Nuclear Chemistry, 2016, 307, 2167-2172.	1.5	12
13	Long-term variations in water quality of lakes in Rokkasho, Aomori, Japan, from 2004 to 2015. Japanese Journal of Limnology, 2016, 78, 75-85.	0.1	4
14	Tritium activity concentrations and residence times of groundwater collected in Rokkasho, Japan. Radiation Protection Dosimetry, 2015, 167, 201-205.	0.8	4
15	Concentration of 129I in aquatic biota collected from a lake adjacent to the spent nuclear fuel reprocessing plant in Rokkasho, Japan. Radiation Protection Dosimetry, 2015, 167, 176-180.	0.8	12
16	Vertical distribution of radiation dose rates in the water of a brackish lake in Aomori Prefecture, Japan. Radiation Protection Dosimetry, 2015, 167, 235-238.	0.8	0
17	Atmospheric deposition of radionuclides (7Be, 210Pb, 134Cs, 137Cs and 40K) during 2000–2012 at Rokkasho, Japan, and impact of the Fukushima Dai-ichi Nuclear Power Plant accident. Journal of Radioanalytical and Nuclear Chemistry, 2015, 303, 1217-1222.	1.5	4
18	Iodine-129 in water samples collected adjacent to a spent nuclear fuel reprocessing plant in Rokkasho, Japan. Journal of Radioanalytical and Nuclear Chemistry, 2015, 303, 1211-1215.	1.5	17

#	Article	IF	CITATIONS
19	Nuclear accident-derived 3H in river water of Fukushima Prefecture during 2011–2014. Journal of Environmental Radioactivity, 2015, 146, 102-109.	1.7	26
20	Effects of radiocesium inventory on 137Cs concentrations in river waters of Fukushima, Japan, under base-flow conditions. Journal of Environmental Radioactivity, 2015, 144, 86-95.	1.7	49
21	Relationship between the radiocesium interception potential and the transfer of radiocesium from soil to soybean cultivated in 2011 in Fukushima Prefecture, Japan. Journal of Environmental Radioactivity, 2014, 137, 119-124.	1.7	35
22	Characteristics of hydrogen and oxygen stable isotope ratios in precipitation collected in a snowfall region, Aomori Prefecture, Japan. Geochemical Journal, 2014, 48, 9-18.	1.0	16
23	Fluvial discharges of radiocaesium from watersheds contaminated by the Fukushima Dai-ichi Nuclear Power Plant accident, Japan. Journal of Environmental Radioactivity, 2013, 118, 96-104.	1.7	170
24	Time-dependent changes of phytoavailability of Cs added to allophanic Andosols in laboratory cultivations and extraction tests. Journal of Environmental Radioactivity, 2013, 122, 29-36.	1.7	56
25	Daily Radionuclide Ingestion and Internal Radiation Doses in Aomori Prefecture, Japan. Health Physics, 2013, 105, 340-350.	0.5	4
26	Radiocarbon Concentrations in Environmental Samples Collected Near the Spent Nuclear Fuel Reprocessing Plant at Rokkasho, Aomori, Japan, During Test Operation Using Spent Nuclear Fuel. Health Physics, 2013, 105, 236-244.	0.5	10
27	Concentration of 3H in plants around Fukushima Dai-ichi Nuclear Power Station. Scientific Reports, 2012, 2, 947.	3.3	28
28	Preface to first special issue on Fukushima. Journal of Environmental Radioactivity, 2012, 111, 1.	1.7	12
29	Tritium concentrations in the atmospheric environment at Rokkasho, Japan before the final testing of the spent nuclear fuel reprocessing plant. Journal of Environmental Radioactivity, 2011, 102, 837-842.	1.7	49
30	AtKUP/HAK/KT9, a K <sup>+</sup> Transporter from <i>Arabidopsis thaliana</i> , Mediates Cs <sup>+</sup> Uptake in <i>Escherichia coli</i> . Bioscience, Biotechnology and Biochemistry, 2010, 74, 203-205.	1.3	42
31	Effect of the counter anion of cesium on foliar uptake and translocation. Journal of Environmental Radioactivity, 2009, 100, 54-57.	1.7	21
32	Air mass origins by back trajectory analysis for evaluating atmospheric 210Pb concentrations at Rokkasho, Aomori, Japan. Journal of Radioanalytical and Nuclear Chemistry, 2009, 279, 493-498.	1.5	6
33	Inventories of 239+240Pu, 137Cs, and excess 210Pb in sediments from freshwater and brackish lakes in Rokkasho, Japan, adjacent to a spent nuclear fuel reprocessing plant. Journal of Environmental Radioactivity, 2009, 100, 835-840.	1.7	17
34	Concentration and specific activity of fallout 137Cs in extracted and particle-size fractions of cultivated soils. Journal of Environmental Radioactivity, 2008, 99, 875-881.	1.7	129
35	Total deposition velocities and scavenging ratios of 7Be and 210Pb at Rokkasho, Japan. Journal of Radioanalytical and Nuclear Chemistry, 2008, 277, 347-355.	1.5	33
36	Plant induced changes in concentrations of caesium, strontium and uranium in soil solution with reference to major ions and dissolved organic matter. Journal of Environmental Radioactivity, 2008, 99, 900-911.	1.7	25

Shun'ichi Hisamatsu

#	Article	IF	CITATIONS
37	Deposition of 137Cs in Rokkasho, Japan and its relation to Asian dust. Journal of Environmental Radioactivity, 2007, 95, 1-9.	1.7	34
38	Rapid Method for the Analysis of Plutonium Isotopes in a Soil Sample within 60 min. Analytical Sciences, 2006, 22, 309-311.	1.6	22
39	Accumulation of uranium derived from long-term fertilizer applications in a cultivated Andisol. Science of the Total Environment, 2006, 367, 924-931.	8.0	66
40	Development of Rapid Plutonium Analysis for Environmental Samples by Isotope Dilution/Inductively Coupled Plasma Mass Spectrometry with On-line Column. Analytical Sciences, 2005, 21, 205-208.	1.6	28
41	Uptake and distribution of 90Sr and stable Sr in rice plants. Journal of Environmental Radioactivity, 2005, 81, 221-231.	1.7	60
42	Effect of Long-term Fertilizer Application on the Concentration and Solubility of Major and Trace Elements in a Cultivated Andisol. Soil Science and Plant Nutrition, 2005, 51, 251-260.	1.9	26
43	Free water 3H concentrations in serum samples collected during 1969–1992 in Akita, Japan. Health Physics, 2003, 85, 204-209.	0.5	2
44	VALIDATION OF ICRP METABOLIC MODELS FOR THE TRANSURANICS IN A JAPANESE POPULATION. Health Physics, 2003, 85, 701-708.	0.5	3
45	Transfer of 137Cs and stable Cs from paddy soil to polished rice in Aomori, Japan. Journal of Environmental Radioactivity, 2002, 59, 351-363.	1.7	136
46	Effective Dose Due to Environmental .GAMMARay for the People of Akita Prefecture Radioisotopes, 2001, 50, 435-441.	0.2	0
47	210Pb Ingestion in Akita City, Japan Radioisotopes, 1992, 41, 574-576.	0.2	3
48	Interlaboratory comparison of low-level organic tritium measurement in environmental samples Radioisotopes, 1990, 39, 457-463.	0.2	9
49	Tritium concentrations in some European foods. Journal of Environmental Radioactivity, 1989, 10, 251-255.	1.7	9
50	Concentration of Fallout Plutonium in Tissues of Japanese Who Died during 1980-1984. Radiation Research, 1987, 109, 245.	1.5	7
51	Effect of Gavaged Chemical Form of 241 Am on Its Retention in Mice. Radiation Research, 1987, 111, 334.	1.5	2
52	Geographical distribution of cerebrovascular disease mortality and food intakes in Japan. Social Science and Medicine, 1987, 24, 401-407.	3.8	28
53	New spectrometric technique for the determination of the isotopic ratio of. Nuclear Instruments and Methods in Physics Research, Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, 1984, 226, 482-486.	1.6	8
54	Placental Transfer and Distribution of 241 Am in the Rat. Radiation Research, 1983, 94, 81.	1.5	6