Cecily J Forsyth

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6753972/publications.pdf

Version: 2024-02-01

1040056 839539 31 345 9 18 citations g-index h-index papers 32 32 32 626 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	RUNX1-mutated families show phenotype heterogeneity and a somatic mutation profile unique to germline predisposed AML. Blood Advances, 2020, 4, 1131-1144.	5.2	102
2	TIDEL-II: first-line use of imatinib in CML with early switch to nilotinib for failure to achieve time-dependent molecular targets. Blood, 2015, 125, 915-923.	1.4	77
3	The incidence and natural history of dasatinib complications in the treatment of chronic myeloid leukemia. Blood Advances, 2017, 1, 802-811.	5.2	31
4	Upfront lower dose lenalidomide is less toxic and does not compromise efficacy for vulnerable patients with relapsed refractory multiple myeloma: final analysis of the phase II RevLite study. British Journal of Haematology, 2017, 177, 441-448.	2.5	21
5	The natural history of vascular and other complications in patients treated with nilotinib for chronic myeloid leukemia. Blood Advances, 2019, 3, 1084-1091.	5.2	17
6	Changing incidence of myeloproliferative neoplasms in Australia, 2003â€2014. American Journal of Hematology, 2019, 94, E107-E109.	4.1	16
7	Non-anaemic iron deficiency. Australian Prescriber, 2021, 44, 193-196.	1.0	15
8	A Randomised Dose De-Escalation Study of Oral Fludarabine, ±Oral Cyclophosphamide and Intravenous Rituximab As First-Line Therapy of Fit Patients with Chronic Lymphocytic Leukaemia (CLL) Aged ≥65 Years: Final Analysis of Response and Toxicity. Blood, 2014, 124, 3325-3325.	1.4	13
9	Efficacy and Safety of ABP 798: Results from the JASMINE Trial in Patients with Follicular Lymphoma in Comparison with Rituximab Reference Product. Targeted Oncology, 2020, 15, 599-611.	3.6	12
10	Recommendations for the use of pegylated interferonâ€Î± in the treatment of classical myeloproliferative neoplasms. Internal Medicine Journal, 2019, 49, 948-954.	0.8	7
11	The delayed diagnosis of myeloproliferative neoplasms is common and results in a high incidence of potentially preventable thrombotic complications. Pathology, 2018, 50, 775-776.	0.6	6
12	Outpatient rituximab, ifosfamide, etoposide (R-IE) in patients older than 60 years with relapsed or refractory diffuse large B-cell lymphoma who are not candidates for stem cell transplantation. Leukemia and Lymphoma, 2020, 61, 91-97.	1.3	6
13	Toxicity Is Not Associated with Age or Comorbidity Score in a Randomised Study of Oral Fludarabine and Cyclophosphamide and IV Rituximab (FCR) As First-Line Therapy of Fit, Elderly Patients with Chronic Lymphocytic Leukemia (CLL). Blood, 2014, 124, 4695-4695.	1.4	5
14	Early Dose-Escalation in Chronic Myeloid Leukaemia Patients with Low Plasma Imatinib Levels Leads to Equivalent BCR-ABL Values and Drug Levels at 6 Months to Those with Optimal Drug Levels: First Analysis From the TIDEL II Trial of De-Novo Patients Treated with 600mg Imatinib Blood, 2009, 114, 1131-1131.	1.4	4
15	Delayedâ€onset heparinâ€induced thrombocytopenia complicated by arterial and venous thromboses. Internal Medicine Journal, 2018, 48, 98-100.	0.8	3
16	Proteasome inhibitor-induced coronary vasospasm in multiple myeloma: a case report. European Heart Journal - Case Reports, 2021, 5, ytab076.	0.6	3
17	Enzastaurin in Patients with Non-Hodgkin Lymphomas: A Multicenter, Open-Label, Screening Study Blood, 2009, 114, 3719-3719.	1.4	2
18	Raccoon eyes in systemic light chain amyloidosis. Medical Journal of Australia, 2017, 206, 384-384.	1.7	1

#	Article	IF	CITATIONS
19	Cutaneous multiple myeloma. British Journal of Haematology, 2020, 190, 134-134.	2.5	1
20	Subacute combined degeneration of the spinal cord due to pernicious anaemia with an erroneously normal active <scp>B12</scp> level. Internal Medicine Journal, 2021, 51, 1184-1185.	0.8	1
21	Long-Term Outcomes and Health-Related Quality of Life (HRQoL) By Response Status for Bortezomib, Melphalan, and Prednisone (VMP) ± Daratumumab (DARA) in Alcyone. Blood, 2020, 136, 43-44.	1.4	1
22	115â€Erroneously normal active B12 level in a case of subacute combined degeneration of the cord. , 2021, , .		0
23	FDG PET-CT in Primary Staging and Management of Hodgkin Lymphoma (HL) and Non-Hodgkin Lymphoma (NHL): Experience in 465 Consecutive Patients Blood, 2006, 108, 2398-2398.	1.4	O
24	Immunoglobulin G (IgG) Subclasses in Chronic Lymphocytic Leukaemia. Blood, 2008, 112, 4180-4180.	1.4	0
25	Quality of Life in Fit Elderly Patients with Chronic Lymphocytic Leukemia (CLL) Receiving Oral Fludarabine-Based Regimens As First Line Therapy: Australasian Leukaemia and Lymphoma Group (ALLG) CLL5 Trial. Blood, 2015, 126, 5295-5295.	1.4	O
26	Upfront Imatinib with Selective Early Switching to Nilotinib Leads to Excellent Achievement of Deep Molecular Response in Chronic Phase CML: 5 Year (Final) Analysis of the TIDEL-II Study. Blood, 2016, 128, 939-939.	1.4	0
27	Efficacy and safety of ABP 798 compared with rituximab: Results from the comparative clinical study in patients with non-Hodgkin's Journal of Clinical Oncology, 2020, 38, 8044-8044.	1.6	O
28	Variation in Use of Immunoglobulin and Impact on Survival in Multiple Myeloma: A Report from the Australian and New Zealand Myeloma and Related Diseases Registry (MRDR). Blood, 2021, 138, 4757-4757.	1.4	0
29	Variable incidence of myeloproliferative neoplasms in Australia. Internal Medicine Journal, 2021, 51, 1979-1980.	0.8	O
30	Totality of Scientific Evidence in the Development of ABP 798, a Biosimilar to Rituximab. Blood, 2020, 136, 35-36.	1.4	0
31	Paraprotein associated heparin resistance during cardiopulmonary bypass. Perfusion (United) Tj ETQq1 1 0.7843	314 ₁ rgBT /0	Overlock 10 T