Akira Osawa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6761730/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Seasonal patterns of fine root dynamics and their contribution to net primary production in hinoki cypress (Chamaecyparis obtusa) and konara oak (Quercus serrata) forests. Trees - Structure and Function, 2021, 35, 255-271.	1.9	7
2	Application of a u-w method for the detection of boreal forest response to environmental changes in Canada. Journal of Forest Research, 2021, 26, 303-313.	1.4	1
3	Quantification and Understanding of Above and Belowground Biomass in Medium Saline Zone of the Sundarbans, Bangladesh: The Relationships with Forest Attributes. Journal of Sustainable Forestry, 2020, 39, 331-345.	1.4	12
4	Soil conditions required for reaction wood formation of drunken trees in a continuous permafrost region. Arctic, Antarctic, and Alpine Research, 2020, 52, 47-59.	1.1	5
5	Two-decadal trends in aboveground litterfall and net primary production in self-thinning <i>Pinus banksiana</i> stands in Wood Buffalo National Park, NWT, Canada. Scandinavian Journal of Forest Research, 2019, 34, 102-114.	1.4	1
6	Decomposition rate of extraradical hyphae of arbuscular mycorrhizal fungi decreases rapidly over time and varies by hyphal diameter and season. Soil Biology and Biochemistry, 2019, 136, 107533.	8.8	10
7	Phenology and litterfall production of <i>Bruguiera sexangula</i> (Lour.) Poir. in the Sundarbans mangrove forests, Bangladesh. Forest Science and Technology, 2019, 15, 165-172.	0.8	5
8	Comparison of biomass and net primary productivity among three species in a subtropical mangrove forest at Manko Wetland, Okinawa, Japan. Regional Studies in Marine Science, 2019, 25, 100475.	0.7	6
9	Evidence for the coupling of extraradical mycorrhizal hyphae production to plant C assimilation in Japanese warm-temperate forest of arbuscular mycorrhizal and ectomycorrhizal tree species. European Journal of Soil Biology, 2018, 88, 73-79.	3.2	2
10	Species composition, biomass, and net primary productivity of mangrove forest in Okukubi River, Okinawa Island, Japan. Regional Studies in Marine Science, 2017, 12, 19-27.	0.7	23
11	Fine root dynamics after soil disturbance evaluated with a root scanner method. Plant and Soil, 2017, 419, 467-487.	3.7	16
12	Fine roots: when anisotropy matters. Tree Physiology, 2017, 37, 693-696.	3.1	3
13	Estimation of fine-root production using rates of diameter-dependent root mortality, decomposition and thickening in forests. Tree Physiology, 2016, 36, 513-523.	3.1	15
14	Size-dependent morphological and chemical property of fine root litter decomposition. Plant and Soil, 2015, 393, 283-295.	3.7	12
15	A new approach to estimate fine root production, mortality, and decomposition using litter bag experiments and soil core techniques. Plant and Soil, 2012, 355, 167-181.	3.7	50
16	Population changes of early successional forest species after shifting cultivation in Northwestern Vietnam. New Forests, 2011, 41, 247-262.	1.7	25
17	Recovery of Vegetation Structure and Species Diversity after Shifting Cultivation in Northwestern Vietnam, with Special Reference to Commercially Valuable Tree Species. ISRN Ecology, 2011, 2011, 1-12.	1.0	4
18	Individual-based measurement and analysis of root system development: case studies for Larix gmelinii trees growing on the permafrost region in Siberia. Journal of Forest Research, 2007, 12, 103-112.	1.4	31