

# Iserhard, Ca

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/676886/publications.pdf>

Version: 2024-02-01

30  
papers

414  
citations

759233

12  
h-index

839539

18  
g-index

31  
all docs

31  
docs citations

31  
times ranked

322  
citing authors

#	ARTICLE	IF	CITATIONS
1	Morphological matching and phenological overlap promote niche partitioning and shape a mutualistic plant-hawkmoth network. <i>Ecological Entomology</i> , 2021, 46, 292-300.	2.2	6
2	The hidden side of diversity: Effects of imperfect detection on multiple dimensions of biodiversity. <i>Ecology and Evolution</i> , 2021, 11, 12508-12519.	1.9	8
3	Sampling performance of bait traps in high Andean fruit-feeding butterflies. <i>Neotropical Biodiversity</i> , 2021, 7, 507-513.	0.5	2
4	Discolouring the Amazon Rainforest: how deforestation is affecting butterfly coloration. <i>Biodiversity and Conservation</i> , 2020, 29, 2821-2838.	2.6	14
5	"INSETOS, E DA RESSIGNIFICANDO AS DIMENSÕES DA EXTENSÃO UNIVERSITÁRIA COM A PANDEMIA DA COVID-19. <i>Expressa Extensão</i> , 2020, 26, 285-299.	0.1	0
6	Combining functional traits and phylogeny to disentangling Amazonian butterfly assemblages on anthropogenic gradients. <i>Ecosphere</i> , 2019, 10, e02837.	2.2	27
7	Effect of habitat loss and fragmentation on fruit-feeding butterflies in the Brazilian Atlantic Forest. <i>Canadian Journal of Zoology</i> , 2019, 97, 588-596.	1.0	17
8	How urbanization affects multiple dimensions of biodiversity in tropical butterfly assemblages. <i>Biodiversity and Conservation</i> , 2019, 28, 621-638.	2.6	33
9	Atlantic butterflies: a data set of fruit-feeding butterfly communities from the Atlantic forests. <i>Ecology</i> , 2018, 99, 2875-2875.	3.2	24
10	Monitoring fruit-feeding butterfly assemblages in two vertical strata in seasonal Atlantic Forest: temporal species turnover is lower in the canopy. <i>Journal of Tropical Ecology</i> , 2017, 33, 345-355.	1.1	29
11	Monitoring Temporal Variation to Assess Changes in the Structure of Subtropical Atlantic Forest Butterfly Communities. <i>Environmental Entomology</i> , 2017, 46, 804-813.	1.4	16
12	<i>Thysbe silvestre</i> sp. nov. (Lepidoptera: Riodinidae): a new myrmecophilous butterfly from the Brazilian Atlantic Forest. <i>Austral Entomology</i> , 2016, 55, 138-146.	1.4	3
13	<i>Stichelia pelotensis</i> (Lepidoptera, Riodinidae): conservation, notes, and rediscovery of an endangered butterfly from southern Brazil. <i>Revista Brasileira De Entomologia</i> , 2016, 60, 105-108.	0.4	1
14	First Record of <i>Aricoris cinerica</i> (Stichel, 1910) (Riodinidae) from Brazil and Update on Its Geographical Distribution. <i>Journal of the Lepidopterists' Society</i> , 2014, 68, 69-71.	0.2	3
15	Distribution patterns of riodinid butterflies (Lepidoptera: Riodinidae) from southern Brazil. <i>Zoological Studies</i> , 2014, 53, .	0.3	12
16	<i>Paulogramma hydarnis</i> (n. comb.) (Nymphalidae: Biblidinae): Distribution, Systematic Position, and Conservation Status of a Rare and Endangered Butterfly. <i>Neotropical Entomology</i> , 2014, 43, 218-226.	1.2	14
17	New Geographical Records for the Threatened Butterfly <i>Actinote quadra</i> (Lepidoptera: Nymphalidae: Tj ETQq1 1 0.784314 rgBT /Overloc	0.2	9
18	Maximized sampling of butterflies to detect temporal changes in tropical communities. <i>Journal of Insect Conservation</i> , 2013, 17, 615-622.	1.4	17

#	ARTICLE	IF	CITATIONS
19	Borboletas (Lepidoptera: Papilionoidea e Hesperioidea) de Matas Paludosas e Matas de Restinga da Planície Costeira da região Sul do Brasil. <i>Biota Neotropica</i> , 2012, 12, 181-190.	1.0	12
20	The endangered butterfly <i>Charonias theano</i> (Boisduval) (Lepidoptera: Pieridae): current status, threats and its rediscovery in the state of São Paulo, southeastern Brazil. <i>Neotropical Entomology</i> , 2011, 40, 669-676.	1.2	7
21	Borboletas (Lepidoptera: Papilionoidea e Hesperioidea) do Sudoeste do Pampa Brasileiro, Uruguaiana, Rio Grande do Sul, Brasil. <i>Biota Neotropica</i> , 2011, 11, 355-360.	1.0	8
22	A new genus and species of Euptychiina (Lepidoptera: Nymphalidae: Satyrinae) from southern Brazil. <i>Neotropical Entomology</i> , 2011, 40, 231-237.	1.2	21
23	Borboletas frugívoras (Lepidoptera: Nymphalidae) ocorrentes em um fragmento de Floresta Ombrófila Mista no Rio Grande do Sul, Brasil. <i>Biota Neotropica</i> , 2011, 11, 385-390.	1.0	11
24	Fruit-feeding butterflies guide of subtropical Atlantic Forest and Araucaria Moist Forest in State of Rio Grande do Sul, Brazil. <i>Biota Neotropica</i> , 2011, 11, 253-274.	1.0	23
25	Borboletas (Lepidoptera: Papilionoidea e Hesperioidea) ocorrentes em diferentes ambientes na Floresta Ombrófila Mista e nos Campos de Cima da Serra do Rio Grande do Sul, Brasil. <i>Biota Neotropica</i> , 2010, 10, 309-320.	1.0	23
26	Natural History, New Records, and Notes on the Conservation Status of <i>Cyanophrys bertha</i> (Jones) (Lepidoptera: Lycaenidae). <i>Proceedings of the Entomological Society of Washington</i> , 2010, 112, 54-60.	0.2	9
27	Lepidoptera, Nymphalidae, Heliconiinae, <i>Heliconius sara</i> apseudes (Hübner, 1813): distribution extension. <i>Check List</i> , 2010, 6, 316.	0.4	2
28	Occurrence of Lepidopterism caused by the moth <i>Hylesia nigricans</i> (Berg) (Lepidoptera: Saturniidae) in Rio Grande do Sul State, Brazil. <i>Neotropical Entomology</i> , 2007, 36, 612-615.	1.2	17
29	Lista de espécies de borboletas (Lepidoptera, Papilionoidea e Hesperioidea) da região do vale do rio Maquinã, Rio Grande do Sul, Brasil. <i>Revista Brasileira De Zoologia</i> , 2004, 21, 649-662.	0.5	41
30	Are there differences in the diversity of bees between organic and conventional agroecosystems in the Pampa biome?. <i>Journal of Apicultural Research</i> , 0, , 1-13.	1.5	4