

Cristina Petrisor

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6769689/cristina-petrisor-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

16
papers

110
citations

7
h-index

10
g-index

19
ext. papers

144
ext. citations

2.5
avg, IF

2.61
L-index

#	Paper	IF	Citations
16	Association of Moral Distress with Anxiety, Depression, and an Intention to Leave among Nurses Working in Intensive Care Units during the COVID-19 Pandemic. <i>Healthcare (Switzerland)</i> , 2021 , 9,	3.4	7
15	A.B.C. approach proposal for POCUS in COVID-19 critically ill patients. <i>Medical Ultrasonography</i> , 2021 , 23, 94-102	1.4	1
14	Clinical Ultrasound Measurements of Hyomental Distance Ratio for the Prediction of Difficult Airway in Patients with and without Morbid Obesity. <i>Diagnostics</i> , 2020 , 10,	3.8	4
13	Which is the Optimal Method of Reconstruction After Laparoscopic Right Hemicolectomy, the Intracorporeal or Extracorporeal Anastomosis Technique?. <i>Chirurgia (Romania)</i> , 2020 , 115, 493-504	1.8	2
12	Diagnostic testing for penicillin allergy: A survey of practices and cost perceptions. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2020 , 75, 436-441	9.3	12
11	Preoperative difficult airway prediction using suprahyoid and infrahyoid ultrasonography derived measurements in anesthesiology. <i>Medical Ultrasonography</i> , 2019 , 21, 83-88	1.4	9
10	The performance of ultrasound-based quantification of the hyomental distance ratio in predicting difficult airway in anaesthesia: A STARD-compliant prospective diagnostic study. <i>European Journal of Anaesthesiology</i> , 2018 , 35, 627-628	2.3	3
9	Ultrasound-based assessment of hyomental distances in neutral, ramped, and maximum hyperextended positions, and derived ratios, for the prediction of difficult airway in the obese population: a pilot diagnostic accuracy study. <i>Anaesthesiology Intensive Therapy</i> , 2018 , 50, 110-116	1.7	7
8	sTREM-1, sIL-2R α and IL-6, but not sCD163, might predict sepsis in polytrauma patients: a prospective cohort study. <i>European Journal of Trauma and Emergency Surgery</i> , 2017 , 43, 363-370	2.3	9
7	Multiple Anesthetic Drug Sensitization in a Hairdresser With Previous Immediate-Type Hypersensitivity to Penicillin: A Case Report. <i>A & A Case Reports</i> , 2017 , 9, 151-153		
6	Flow-assisted basophil activation test for sevoflurane hypersensitivity: A dose-finding in-vitro experimental study. <i>European Journal of Anaesthesiology</i> , 2017 , 34, 851-852	2.3	
5	Can APACHE II, SOFA, ISS, and RTS Severity Scores be used to Predict Septic Complications in Multiple Trauma Patients?. <i>The Journal of Critical Care Medicine</i> , 2016 , 2, 124-130	1.2	7
4	Antibiotic-induced immediate type hypersensitivity is a risk factor for positive allergy skin tests for neuromuscular blocking agents. <i>Allergology International</i> , 2016 , 65, 52-5	4.4	12
3	Letter to the Editor: Early serum procalcitonin, but not C reactive protein, might improve the prediction of sepsis in multiple trauma patients. A prospective observational study. <i>Romanian Journal of Laboratory Medicine</i> , 2015 , 23, 347-350	0.3	1
2	Threshold for basophil activation test positivity in neuromuscular blocking agents hypersensitivity reactions. <i>Allergy, Asthma and Clinical Immunology</i> , 2013 , 9, 42	3.2	21
1	Threshold for positivity and optimal dipyrone concentration in flow cytometry-assisted basophil activation test. <i>Allergy, Asthma and Immunology Research</i> , 2013 , 5, 383-8	5.3	15