
Alf Inge Larsen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6769732/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Targeted Temperature Management for 48 vs 24 Hours and Neurologic Outcome After Out-of-Hospital Cardiac Arrest. JAMA - Journal of the American Medical Association, 2017, 318, 341.	7.4	260
2	Identification of vulnerable plaques and patients by intracoronary near-infrared spectroscopy and ultrasound (PROSPECT II): a prospective natural history study. Lancet, The, 2021, 397, 985-995.	13.7	208
3	Characteristics and outcomes of patients with acute myocardial infarction and angiographically normal coronary arteries. American Journal of Cardiology, 2005, 95, 261-263.	1.6	152
4	Cardiac arrest with continuous mechanical chest compression during percutaneous coronary intervention. Resuscitation, 2007, 75, 454-459.	3.0	94
5	Intracoronary autologous bone marrow cell transfer after myocardial infarction: the BOOST-2 randomised placebo-controlled clinical trial. European Heart Journal, 2017, 38, 2936-2943.	2.2	91
6	Long-Term Prognosis of Patients Presenting With ST-Segment Elevation Myocardial Infarction With No Significant Coronary Artery Disease (from The HORIZONS-AMI Trial). American Journal of Cardiology, 2013, 111, 643-648.	1.6	71
7	Effect of exercise training in patients with heart failure: a pilot study on autonomic balance assessed by heart rate variability. European Journal of Cardiovascular Prevention and Rehabilitation, 2004, 11, 162-167.	2.8	54
8	Coronary blood flow and perfusion pressure during coronary angiography in patients with ongoing mechanical chest compression: A report on 6 cases. Resuscitation, 2010, 81, 493-497.	3.0	42
9	Time-differentiated target temperature management after out-of-hospital cardiac arrest: a multicentre, randomised, parallel-group, assessor-blinded clinical trial (the TTH48 trial): study protocol for a randomised controlled trial. Trials, 2016, 17, 228.	1.6	32
10	BEtablocker Treatment After acute Myocardial Infarction in revascularized patients without reduced left ventricular ejection fraction (BETAMI): Rationale and design of a prospective, randomized, open, blinded end point study. American Heart Journal, 2019, 208, 37-46.	2.7	20
11	Persistent left superior vena cava. Use of an innominate vein between left and right superior caval veins for the placement of a right ventricular lead during ICD/CRT implantation. European Heart Journal, 2005, 26, 2178-2178.	2.2	17
12	Cardiac arrest as a reportable condition: a cohort study of the first 6 years of the Norwegian out-of-hospital cardiac arrest registry. BMJ Open, 2020, 10, e038133.	1.9	17
13	Burden of treatment in patients with chronic heart failure – A cross-sectional study. Heart and Lung: Journal of Acute and Critical Care, 2021, 50, 369-374.	1.6	15
14	B-type natriuretic peptide is related to histological skeletal muscle abnormalities in patients with chronic heart failure. International Journal of Cardiology, 2009, 136, 358-362.	1.7	11
15	BNP in acute coronary syndromes: the heart expresses its suffering. European Heart Journal, 2004, 25, 1284-1286.	2.2	9
16	Adding stress biomarkers to high-sensitivity cardiac troponin for rapid non-ST-elevation myocardial infarction rule-out protocols. European Heart Journal: Acute Cardiovascular Care, 2022, 11, 201-212.	1.0	9
17	Cardiac resynchronization therapy improves minute ventilation/carbon dioxide production slope and skeletal muscle capillary density without reversal of skeletal muscle pathology or inflammation. Europace, 2013, 15, 857-864.	1.7	8
18	Effects of interval training on inflammatory biomarkers in patients with ischemic heart failure. Scandinavian Cardiovascular Journal, 2019, 53, 213-219.	1.2	8

Alf Inge Larsen

#	Article	IF	CITATIONS
19	Confronting one's vulnerability – patients with chest pain participating in a highâ€intensity exercise programme. Journal of Clinical Nursing, 2017, 26, 2006-2015.	3.0	7
20	Exercise training and highâ€sensitivity cardiac troponin T in patients with heart failure with reduced ejection fraction. ESC Heart Failure, 2021, 8, 2183-2192.	3.1	7
21	Change to a primary PCI program increases number of patients offered reperfusion therapy and significantly reduces mortality. International Journal of Cardiology, 2008, 127, 208-213.	1.7	5
22	A statistical analysis protocol for the time-differentiated target temperature management after out-of-hospital cardiac arrest (TTH48) clinical trial. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2016, 24, 138.	2.6	5
23	Impact of an exercise training program on cardiac neuronal function in heart failure patients on optimal medical therapy. Journal of Nuclear Cardiology, 2018, 25, 1164-1171.	2.1	5
24	Large and small vessel vasoconstriction following coronary artery stenting. International Journal of Cardiology, 2006, 113, 61-65.	1.7	4
25	Post resuscitation care $\hat{a} \in$ some words of caution and a call for action. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2015, 23, 89.	2.6	4
26	<scp>QRS</scp> fragmentation is associated with increased risk of ventricular arrhythmias in highâ€risk patients; Data from the <scp>SMASH</scp> 1 Study. Annals of Noninvasive Electrocardiology, 0, , .	1.1	2
27	Aortic calcification; from innocent bystander to independent predictor; the delicate balance in biology; da aaaCapo: Editorial accompanying †Abdominal aortic calcification – from ancient friend to modern foe'. European Journal of Preventive Cardiology, 2022, 28, e20-e24.	1.8	1
28	Increased functional capacity after 12 weeks of exercise training does not transform into improved skeletal muscle metabolism or ultrastructure in heart failure patients on modern optimal medical therapy. European Journal of Preventive Cardiology, 2020, , 2047487320919863.	1.8	1
29	The Localization and Characterization of Ischemic Scars in relation to the Infarct Related Coronary Artery Assessed by Cardiac Magnetic Resonance and a Novel Automatic Postprocessing Method. Cardiology Research and Practice, 2015, 2015, 1-9.	1.1	0
30	Psycho-educational programmes led by experienced nurses may improve the coping abilities of implantable cardioverter defibrillator recipients. Evidence-based Nursing, 2016, 19, 79-80.	0.2	0
31	Blood Lactate AUC Is a Sensitive Test for Evaluating the Effect of Exercise Training on Functional Work Capacity in Patients with Chronic Heart Failure. Rehabilitation Research and Practice, 2021, 2021, 1-6.	0.6	0