## Katarzyna SykÅ,owska-Baranek

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6774450/publications.pdf

Version: 2024-02-01

471371 526166 45 820 17 27 citations h-index g-index papers 46 46 46 734 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	HPLC-PDA-ESI-HRMS-Based Profiling of Secondary Metabolites of Rindera graeca Anatomical and Hairy Roots Treated with Drought and Cold Stress. Cells, 2022, 11, 931.	1.8	7
2	Polyurethane Foam Rafts Supported In Vitro Cultures of Rindera graeca Roots for Enhanced Production of Rinderol, Potent Proapoptotic Naphthoquinone Compound. International Journal of Molecular Sciences, 2022, 23, 56.	1.8	5
3	Polyscias filicifolia (Araliaceae) Hairy Roots with Antigenotoxic and Anti-Photogenotoxic Activity. Molecules, 2022, 27, 186.	1.7	3
4	Metabolic Modifications in Terpenoid and Steroid Pathways Triggered by Methyl Jasmonate in Taxus $ ilde{A}-$ media Hairy Roots. Plants, 2022, $11,1120$ .	1.6	8
5	Application of Priming Strategy for Enhanced Paclitaxel Biosynthesis in Taxus × Media Hairy Root Cultures. Cells, 2022, 11, 2062.	1.8	6
6	In Vitro Response of Polyscias filicifolia (Araliaceae) Shoots to Elicitation with Alarmone–Diadenosine Triphosphate, Methyl Jasmonate, and Salicylic Acid. Cells, 2021, 10, 419.	1.8	8
7	MTMS-Based Aerogel Constructs for Immobilization of Plant Hairy Roots: Effects on Proliferation of Rindera graeca Biomass and Extracellular Secretion of Naphthoquinones. Journal of Functional Biomaterials, 2021, 12, 19.	1.8	9
8	Chemical Profile and Screening of Bioactive Metabolites of Rindera graeca (A. DC.) Bois. & DC. (Boraginaceae) In Vitro Cultures. Plants, 2021, 10, 834.	1.6	7
9	Comparative Study of the Genetic and Biochemical Variability of Polyscias filicifolia (Araliaceae) Regenerants Obtained by Indirect and Direct Somatic Embryogenesis as a Source of Triterpenes. International Journal of Molecular Sciences, 2021, 22, 5752.	1.8	8
10	Development of Taxus spp. Hairy Root Cultures for Enhanced Taxane Production. Reference Series in Phytochemistry, 2021, , 541-559.	0.2	0
11	Antigenotoxic, Anti-photogenotoxic, and Antioxidant Properties of Polyscias filicifolia Shoots Cultivated In Vitro. Molecules, 2020, 25, 1090.	1.7	11
12	Bioactive rinderol and cynoglosol isolated from Cynoglossum columnae Ten. in vitro root culture. Industrial Crops and Products, 2019, 137, 446-452.	2.5	11
13	Comparison of elicitor-based effects on metabolic responses of Taxus × media hairy roots in perfluorodecalin-supported two-phase culture system. Plant Cell Reports, 2019, 38, 85-99.	2.8	25
14	Development of Taxus spp. Hairy Root Cultures for Enhanced Taxane Production. Reference Series in Phytochemistry, 2019, , 1-19.	0.2	1
15	Lignan accumulation in two-phase cultures of Taxus x media hairy roots. Plant Cell, Tissue and Organ Culture, 2018, 133, 371-384.	1.2	8
16	Hairy Root Cultures for the Production of Anti-cancer Naphthoquinone Compounds. Current Medicinal Chemistry, 2018, 25, 4718-4739.	1.2	14
17	A cellulase-supported two-phase in situ system for enhanced biosynthesis of paclitaxel in Taxus × media hairy roots. Acta Physiologiae Plantarum, 2018, 40, 1.	1.0	11
18	Stimulation of phenolic compounds production in the in vitro cultivated Polyscias filicifolia Bailey shoots and evaluation of the antioxidant and cytotoxic potential of plant extracts. Acta Societatis Botanicorum Poloniae, 2018, 87, .	0.8	7

#	Article	IF	CITATIONS
19	Pyrrolizidine alkaloids from Cynoglossum columnae Ten. (Boraginaceae). Phytochemistry Letters, 2016, 15, 234-237.	0.6	16
20	Antigenotoxic, anti-photogenotoxic and antioxidant activities of natural naphthoquinone shikonin and arnebia euchroma callus extracts evaluated by the umu-test and EPR method. Toxicology in Vitro, 2015, 30, 364-372.	1.1	33
21	Perfluorodecalin-supported system enhances taxane production in hairy root cultures of Taxus xÂmedia var. Hicksii carrying a taxadiene synthase transgene. Plant Cell, Tissue and Organ Culture, 2015, 120, 1051-1059.	1.2	35
22	Paclitaxel production and PAL activity in hairy root cultures of Taxus x media var. Hicksii carrying a taxadiene synthase transgene elicited with nitric oxide and methyl jasmonate. Acta Physiologiae Plantarum, 2015, 37, 1.	1.0	24
23	Biotechnological approaches to enhance salidroside, rosin and its derivatives production in selected Rhodiola spp. in vitro cultures. Phytochemistry Reviews, 2015, 14, 657-674.	3.1	63
24	Approaches of <i>Rhodiola kirilowii</i> and <i>Rhodiola rosea</i> field cultivation in Poland and their potential health benefits. Annals of Agricultural and Environmental Medicine, 2015, 22, 281-285.	0.5	17
25	Biotransformation of cinnamyl alcohol to rosavins by non-transformed wild type and hairy root cultures of Rhodiola kirilowii. Biotechnology Letters, 2014, 36, 649-656.	1.1	22
26	Liquid Perfluorodecalin Application for In Situ Extraction and Enhanced Naphthoquinones Production in Arnebia euchroma Cell Suspension Cultures. Applied Biochemistry and Biotechnology, 2014, 172, 2618-2627.	1.4	18
27	TYROSOL GLUCOSYLTRANSFERASE ACTIVITY AND SALIDROSIDE PRODUCTION IN NATURAL AND TRANSFORMED ROOT CULTURES OF RHODIOLA KIRILOWII (REGEL) REGEL ET MAXIMOWICZ. Acta Biologica Cracoviensia Series Botanica, 2013, 55, .	0.5	0
28	Phenolic compounds from in vitro cultures of Rindera graeca Boiss. & Feldr Planta Medica, 2013, 79, .	0.7	5
29	lsolation of pyrrolizidine alkaloids from Cynoglossum columnae Ten. (Boraginaceae). Planta Medica, 2013, 79, .	0.7	2
30	Effect of l-phenylalanine on PAL activity and production of naphthoquinone pigments in suspension cultures of Arnebia euchroma (Royle) Johnst. In Vitro Cellular and Developmental Biology - Plant, 2012, 48, 555-564.	0.9	63
31	Antimicrobial and Cytotoxic Isohexenylnaphthazarins from Arnebia euchroma (Royle) Jonst. (Boraginaceae) Callus and Cell Suspension Culture. Molecules, 2012, 17, 14310-14322.	1.7	64
32	Enhanced production of antitumour naphthoquinones in transgenic hairy root lines of Lithospermum canescens. Plant Cell, Tissue and Organ Culture, 2012, 108, 213-219.	1.2	28
33	Activity of tyrosol glucosyltransferase in Rhodiola kirilowii transgenic root cultures. Planta Medica, 2012, 78, .	0.7	1
34	Phenolic compounds from in vitro cultures of Rindera gareca Boiss. & Heldr Planta Medica, 2012, 78, .	0.7	3
35	Cytotoxic and antimicrobial activity of Cynoglossum columnae Ten. in vitro roots. Planta Medica, 2012, 78, .	0.7	2
36	Lignans from in vitro cultures of transgenic roots of Taxus x media var. Hicksii. Planta Medica, 2012, 78, .	0.7	0

#	Article	IF	CITATIONS
37	Metabolic responses of <i>Taxus media</i> transformed cell cultures to the addition of methyl jasmonate. Biotechnology Progress, 2010, 26, 1145-1153.	1.3	70
38	Enhancement of taxane production in hairy root culture of Taxus x media var. Hicksii. Journal of Plant Physiology, 2009, 166, 1950-1954.	1.6	50
39	Establishment of Rindera graeca transgenic root culture as a source of shikonin derivatives. Planta Medica, 2008, 74, .	0.7	5
40	Production of shikonin derivatives in transgenic roots of Lithospermum canescens (Michx.) Lehm. cultivated in mist bioreactor. Planta Medica, 2008, 74, .	0.7	2
41	The shikonin derivatives and pyrrolizidine alkaloids in hairy root cultures of Lithospermum canescens (Michx.) Lehm Plant Cell Reports, 2006, 25, 1052-1058.	2.8	27
42	Taxane Production in Suspension Culture of Taxus $\tilde{A}-$ Media var. Hicksii Carried Out in Flasks and Bioreactor. Biotechnology Letters, 2005, 27, 1301-1304.	1.1	8
43	Title is missing!. Biotechnology Letters, 2000, 22, 1449-1452.	1.1	30
44	Title is missing!. Biotechnology Letters, 2000, 22, 683-686.	1.1	50
45	Title is missing!. Plant Cell, Tissue and Organ Culture, 1997, 49, 75-79.	1.2	33