
Carina A Thorstensson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6778136/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Physiotherapist or physician as primary assessor for patients with suspected knee osteoarthritis in primary care – a cost-effectiveness analysis of a pragmatic trial. BMC Musculoskeletal Disorders, 2022, 23, 260. | 0.8 | 4 |
| 2 | Change in self-efficacy after participation in a supported self-management program for osteoarthritis – an observational study of 11Â906 patients. Disability and Rehabilitation, 2020, 42, 2133-2140. | 0.9 | 12 |
| 3 | High self-efficacy – a predictor of reduced pain and higher levels of physical activity among patients with osteoarthritis: an observational study. BMC Musculoskeletal Disorders, 2020, 21, 380. | 0.8 | 27 |
| 4 | Physiotherapist as primary assessor for patients with suspected knee osteoarthritis in primary care—a randomised controlled pragmatic study. BMC Musculoskeletal Disorders, 2019, 20, 329. | 0.8 | 4 |
| 5 | Chronic widespread pain and its associations with quality of life and function at a 20- year follow-up of individuals with chronic knee pain at inclusion. BMC Musculoskeletal Disorders, 2019, 20, 592. | 0.8 | 13 |
| 6 | The effect of education and supervised exercise on physical activity, pain, quality of life and self-efficacy - an intervention study with a reference group. BMC Musculoskeletal Disorders, 2018, 19, 198. | 0.8 | 57 |
| 7 | Health enhancing physical activity in patients with hip or knee osteoarthritis - an observational intervention study. BMC Musculoskeletal Disorders, 2017, 18, 42. | 0.8 | 16 |
| 8 | Osteoarthritis: Models for appropriate care across the disease continuum. Best Practice and Research in Clinical Rheumatology, 2016, 30, 503-535. | 1.4 | 123 |
| 9 | A Web-Based Platform for Patients With Osteoarthritis of the Hip and Knee: A Pilot Study. JMIR Research Protocols, 2016, 5, e115. | 0.5 | 49 |
| 10 | Better Management of Patients with Osteoarthritis: Development and Nationwide Implementation of an Evidenceâ€Based Supported Osteoarthritis Selfâ€Management Programme. Musculoskeletal Care, 2015, 13, 67-75. | 0.6 | 121 |
| 11 | Knee joint stabilization therapy in patients with osteoarthritis of the knee and knee instability: Subgroup analyses in a randomized, controlled trial. Journal of Rehabilitation Medicine, 2014, 46, 703-707. | 0.8 | 18 |
| 12 | Dissatisfied patients after total knee arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 2014, 85, 229-233. | 1.2 | 60 |
| 13 | Similar Problem in the Activities of Daily Living but Different Experience: A Qualitative Analysis in Six Rheumatic Conditions and Eight European Countries. Musculoskeletal Care, 2014, 12, 22-33. | 0.6 | 23 |
| 14 | Differences in muscle activity during hand-dexterity tasks between women with arthritis and a healthy reference group. BMC Musculoskeletal Disorders, 2014, 15, 154. | 0.8 | 12 |
| 15 | Risk of sick leave and disability pension in working-age women and men with knee osteoarthritis. Annals of the Rheumatic Diseases, 2013, 72, 401-405. | 0.5 | 85 |
| 16 | ls knee osteoarthritis a symmetrical disease? Analysis of a 12 year prospective cohort study. BMC Musculoskeletal Disorders, 2012, 13, 153. | 0.8 | 85 |
| 17 | Relationship between finger flexion and extension force in healthy women and women with rheumatoid arthritis. Journal of Rehabilitation Medicine, 2012, 44, 605-608. | 0.8 | 12 |
| 18 | Association of lower muscle strength with selfâ€reported knee instability in osteoarthritis of the knee: Results from the Amsterdam Osteoarthritis Cohort. Arthritis Care and Research, 2012, 64, 38-45. | 1.5 | 92 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Sickness absence in musculoskeletal disorders - patients' experiences of interactions with the Social Insurance Agency and health care. A qualitative study. BMC Public Health, 2011, 11, 107. | 1.2 | 34 |
| 20 | Identification of phenotypes with different clinical outcomes in knee osteoarthritis: Data from the osteoarthritis initiative. Arthritis Care and Research, 2011, 63, 1535-1542. | 1.5 | 115 |
| 21 | Choosing surgery: patients' preferences within a trial of treatments for anterior cruciate ligament injury. A qualitative study. BMC Musculoskeletal Disorders, 2009, 10, 100. | 0.8 | 45 |
| 22 | Help-seeking behaviour among people living with chronic hip or knee pain in the community. BMC Musculoskeletal Disorders, 2009, 10, 153. | 0.8 | 52 |
| 23 | Cooperation between gatekeepers in sickness insurance – the perspective of social insurance officers. A qualitative study. BMC Health Services Research, 2008, 8, 231. | 0.9 | 16 |
| 24 | Invited Commentary. Physical Therapy, 2008, 88, 1541-1543. | 1.1 | 1 |
| 25 | How do middle-aged patients conceive exercise as a form of treatment for knee osteoarthritis?. Disability and Rehabilitation, 2006, 28, 51-59. | 0.9 | 53 |
| 26 | Serum levels of Cartilage Oligomeric Matrix Protein (COMP) increase temporarily after physical exercise in patients with knee osteoarthritis. BMC Musculoskeletal Disorders, 2006, 7, 98. | 0.8 | 94 |
| 27 | Six-week high-intensity exercise program for middle-aged patients with knee osteoarthritis: a randomized controlled trial [ISRCTN20244858]. BMC Musculoskeletal Disorders, 2005, 6, 27. | 0.8 | 82 |