

Bo Engdahl

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6781984/bo-engdahl-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

79
papers

1,345
citations

18
h-index

33
g-index

83
ext. papers

1,614
ext. citations

3.4
avg. IF

4.64
L-index

#	Paper	IF	Citations
79	Childhood hearing impairment and fertility in Norway.. <i>Scientific Reports</i> , 2022 , 12, 402	4.9	
78	Weaker association between hearing loss and non-employment in recent generations: the HUNT cohort study.. <i>International Journal of Audiology</i> , 2022 , 1-8	2.6	0
77	Cohort difference in the association between use of recreational firearms and hearing loss: findings from the HUNT study.. <i>International Journal of Audiology</i> , 2022 , 1-7	2.6	1
76	Suicide trends in Norway during the first year of the Covid-19 pandemic. A register-based cohort study.. <i>European Psychiatry</i> , 2022 , 1-24	6	2
75	A possible effect of montelukast on neurological aging examined by the use of register data. <i>International Journal of Clinical Pharmacy</i> , 2021 , 43, 541-548	2.3	7
74	Tinnitus and associations with chronic pain: The population-based Tromsø Study (2015-2016). <i>PLoS ONE</i> , 2021 , 16, e0247880	3.7	1
73	Genome-wide association identifies the first risk loci for psychosis in Alzheimer disease. <i>Molecular Psychiatry</i> , 2021 ,	15.1	5
72	Personal Music Players and Hearing Loss: The HUNT Cohort Study. <i>Trends in Hearing</i> , 2021 , 25, 23312165211015881	3.2	1
71	Explaining better hearing in Norway: a comparison of two cohorts 20 years apart - the HUNT study. <i>BMC Public Health</i> , 2021 , 21, 242	4.1	0
70	Occupational noise exposure and asymmetric hearing loss: Results from the HUNT population study in Norway. <i>American Journal of Industrial Medicine</i> , 2020 , 63, 535-542	2.7	2
69	Physical exercise and chronic pain in university students. <i>PLoS ONE</i> , 2020 , 15, e0235419	3.7	11
68	Better Hearing in Norway: A Comparison of Two HUNT Cohorts 20 Years Apart. <i>Ear and Hearing</i> , 2020 , 42, 42-52	3.4	8
67	Otitis Media in Childhood and Disease in Adulthood: A 40-Year Follow-Up Study. <i>Ear and Hearing</i> , 2020 , 41, 67-71	3.4	5
66	A population-based study of inflammatory mechanisms and pain sensitivity. <i>Pain</i> , 2020 , 161, 338-350	8	7
65	The Graphical Index of Pain: a new web-based method for high-throughput screening of pain. <i>Pain</i> , 2020 , 161, 2255-2262	8	4
64	Physical exercise and chronic pain in university students 2020 , 15, e0235419		
63	Physical exercise and chronic pain in university students 2020 , 15, e0235419		

62 Physical exercise and chronic pain in university students **2020**, 15, e0235419

61 Physical exercise and chronic pain in university students **2020**, 15, e0235419

60 Childhood Sensorineural Hearing Loss and Educational Attainment in Adulthood: Results From the HUNT Study. *Ear and Hearing*, **2019**, 40, 1359-1367 3.4 9

59 Hearing loss, family status and mortality - Findings from the HUNT study, Norway. *Social Science and Medicine*, **2019**, 220, 219-225 5.1 6

58 Prescription database analyses indicates that the asthma medicine montelukast might protect against dementia: a hypothesis to be verified. *Immunity and Ageing*, **2017**, 14, 20 9.7 11

57 Occupational noise exposure, hearing loss, and notched audiograms in the HUNT Nord-Trøndelag hearing loss study, 1996-1998. *Laryngoscope*, **2017**, 127, 1442-1450 3.6 14

56 Childhood otitis media is associated with dizziness in adulthood: the HUNT cohort study. *European Archives of Oto-Rhino-Laryngology*, **2016**, 273, 2047-54 3.5 16

55 Childhood sensorineural hearing loss: effects of combined exposure with aging or noise exposure later in life. *European Archives of Oto-Rhino-Laryngology*, **2016**, 273, 1099-105 3.5 8

54 Occupational noise exposure and hearing: a systematic review. *International Archives of Occupational and Environmental Health*, **2016**, 89, 351-72 3.2 157

53 Simplified risk assessment of noise induced hearing loss by means of 2 spreadsheet models. *International Journal of Occupational Medicine and Environmental Health*, **2016**, 29, 991-999 1.5

52 Noise-induced hearing loss in a longitudinal study of Norwegian railway workers. *BMJ Open*, **2016**, 6, e011923 3 8

51 Noise-induced hearing loss: the diagnosis depends on the doctor's belief. *Occupational and Environmental Medicine*, **2015**, 72, 234 2.1

50 Associations between parental hearing impairment and children's mental health: Results from the Nord-Trøndelag Health Study. *Social Science and Medicine*, **2015**, 147, 252-60 5.1 2

49 The prevalence of notched audiograms in a cross-sectional study of 12,055 railway workers. *Ear and Hearing*, **2015**, 36, e86-92 3.4 17

48 No Association Between Time of Onset of Hearing Loss (Childhood Versus Adulthood) and Self-Reported Hearing Handicap in Adults. *American Journal of Audiology*, **2015**, 24, 549-56 1.8

47 Childhood Otitis Media: A Cohort Study With 30-Year Follow-Up of Hearing (The HUNT Study). *Ear and Hearing*, **2015**, 36, 302-8 3.4 51

46 Association Between Childhood Hearing Disorders and Tinnitus in Adulthood. *JAMA Otolaryngology - Head and Neck Surgery*, **2015**, 141, 983-9 3.9 8

45 Cardiovascular risk factors and hearing loss: The HUNT study. *International Journal of Audiology*, **2015**, 54, 958-66 2.6 32

44	Short- and long-term mortality risk associated with the use of antipsychotics among 26,940 dementia outpatients: a population-based study. <i>American Journal of Geriatric Psychiatry</i> , 2014 , 22, 321-31	6.5	50
43	Sensorineural hearing loss in children: the association with Apgar score. A registry-based study of 392,371 children in Norway. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2014 , 78, 1940-4	1.7	5
42	A cross-sectional study of hearing thresholds among 4627 Norwegian train and track maintenance workers. <i>BMJ Open</i> , 2014 , 4, e005529	3	15
41	Otoacoustic emissions, pure-tone audiometry, and self-reported hearing. <i>International Journal of Audiology</i> , 2013 , 52, 74-82	2.6	18
40	The effect of emotional distress on persistent pelvic girdle pain after delivery: a longitudinal population study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2013 , 120, 32-40	3.7	55
39	Non-random mating and convergence over time for mental health, life satisfaction, and personality: the Nord-Trøndelag Health Study. <i>Behavior Genetics</i> , 2013 , 43, 108-19	3.2	17
38	Hearing status among Norwegian train drivers and train conductors. <i>Occupational Medicine</i> , 2013 , 63, 544-8	2.1	10
37	Occupation and the risk of bothersome tinnitus: results from a prospective cohort study (HUNT). <i>BMJ Open</i> , 2012 , 2, e000512	3	26
36	Heritability of hearing loss. <i>Epidemiology</i> , 2012 , 23, 328-31	3.1	26
35	Longitudinal analysis of emotional problems in children with congenital heart defects: a follow-up from age 6 to 36 months. <i>Journal of Developmental and Behavioral Pediatrics</i> , 2011 , 32, 461-4	2.4	10
34	Longitudinal findings from a Norwegian case-cohort study on internalizing problems in children with congenital heart defects. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2011 , 100, 236-41	3.1	3
33	Concomitant use of anti-dementia drugs with psychotropic drugs in Norway--a population-based study. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 1319-26	2.6	6
32	Effects of changed aircraft noise exposure on the use of outdoor recreational areas. <i>International Journal of Environmental Research and Public Health</i> , 2010 , 7, 3890-915	4.6	4
31	Low heritability of tinnitus: results from the second Nord-Trøndelag health study. <i>JAMA Otolaryngology</i> , 2010 , 136, 178-82		37
30	Effects of changed aircraft noise exposure on experiential qualities of outdoor recreational areas. <i>International Journal of Environmental Research and Public Health</i> , 2010 , 7, 3739-59	4.6	5
29	Maternal infection with toxoplasma gondii in pregnancy and the risk of hearing loss in the offspring. <i>International Journal of Audiology</i> , 2010 , 49, 65-8	2.6	14
28	The association between tinnitus and mental health in a general population sample: results from the HUNT Study. <i>Journal of Psychosomatic Research</i> , 2010 , 69, 289-98	4.1	123
27	Cardiovascular co-medication among users of antiobesity drugs: a population-based study. <i>International Journal of Clinical Pharmacy</i> , 2010 , 32, 752-8		5

26	Use of psychotropic drugs and analgesics among users of antiobesity drugs--a population based study. <i>Pharmacoepidemiology and Drug Safety</i> , 2010 , 19, 273-9	2.6	2
25	Occupation and the risk of hearing impairment--results from the Nord-Trøndelag study on hearing loss. <i>Scandinavian Journal of Work, Environment and Health</i> , 2010 , 36, 250-7	4.3	37
24	Self-reported sleep disturbances due to railway noise: exposure-response relationships for nighttime equivalent and maximum noise levels. <i>Journal of the Acoustical Society of America</i> , 2008 , 124, 257-68	2.2	34
23	Annoyance and self-reported sleep disturbances due to structurally radiated noise from railway tunnels. <i>Applied Acoustics</i> , 2007 , 68, 970-981	3.1	16
22	Birthweight and the risk of childhood sensorineural hearing loss. <i>Paediatric and Perinatal Epidemiology</i> , 2007 , 21, 495-500	2.7	12
21	Hearing loss induced by occupational and impulse noise: results on threshold shifts by frequencies, age and gender from the Nord-Trøndelag Hearing Loss Study. <i>International Journal of Audiology</i> , 2006 , 45, 309-17	2.6	45
20	Screened and unscreened hearing threshold levels for the adult population: results from the Nord-Trøndelag Hearing Loss Study. <i>International Journal of Audiology</i> , 2005 , 44, 213-30	2.6	77
19	Otoacoustic emissions in the general adult population of Nord-Trøndelag, Norway: III. Relationships with pure-tone hearing thresholds. <i>International Journal of Audiology</i> , 2005 , 44, 15-23	2.6	6
18	Annoyance with aircraft noise in local recreational areas and the recreationists' noise situation at home. <i>Journal of the Acoustical Society of America</i> , 2005 , 117, 221-31	2.2	7
17	Annoyance with aircraft noise in local recreational areas, contingent on changes in exposure and other context variables. <i>Journal of the Acoustical Society of America</i> , 2004 , 116, 323-33	2.2	12
16	Subjective responses to aircraft noise in an outdoor recreational setting: a combined field and laboratory study. <i>Journal of Sound and Vibration</i> , 2004 , 276, 981-996	3.9	21
15	Hearing loss associated with ear infections in Nord-Trøndelag, Norway. <i>Ear and Hearing</i> , 2004 , 25, 388-96	3.4	14
14	Detection of transient-evoked otoacoustic emissions and the design of time windows. <i>IEEE Transactions on Biomedical Engineering</i> , 2002 , 49, 132-9	5	8
13	Otoacoustic emissions in the general adult population of Nord-Trøndelag, Norway: I. Distributions by age, gender, and ear side. <i>International Journal of Audiology</i> , 2002 , 41, 64-77	2.6	19
12	Otoacoustic emissions in the general adult population of Nord-Trøndelag, Norway: II. Effects of noise, head injuries, and ear infections. <i>International Journal of Audiology</i> , 2002 , 41, 78-87	2.6	13
11	Otoacoustic emissions and improved pass/fail separation using wavelet analysis and time windowing. <i>Medical and Biological Engineering and Computing</i> , 2001 , 39, 134-9	3.1	12
10	Aircraft noise in recreational areas: A quasi-experimental field study on individual annoyance responses and dose-response relationships. <i>Noise Control Engineering Journal</i> , 1999 , 47, 158-162	0.6	5
9	Aircraft noise in recreational areas: Effects on visitors' experience and well-being. <i>Noise Control Engineering Journal</i> , 1999 , 47, 147-149	0.6	2

8	Effects of noise and exercise on distortion product otoacoustic emissions. <i>Hearing Research</i> , 1996 , 93, 72-82	3.9	23
7	Transient evoked otoacoustic emissions as screening for hearing losses at the school for military training. <i>Scandinavian Audiology</i> , 1996 , 25, 71-8		15
6	Transient-evoked otoacoustic emissions. Helpful tool in the detection of pseudohypacusis. <i>Scandinavian Audiology</i> , 1996 , 25, 173-7		7
5	The effect of noise exposure on the details of distortion product otoacoustic emissions in humans. <i>Journal of the Acoustical Society of America</i> , 1996 , 99, 1573-87	2.2	76
4	Temporary threshold shift and otoacoustic emissions after industrial noise exposure. <i>Scandinavian Audiology</i> , 1995 , 24, 137-41		17
3	Otoacoustic emissions in the first year of life. <i>Scandinavian Audiology</i> , 1994 , 23, 195-200		15
2	Reproducibility and short-term variability of transient evoked otoacoustic emissions. <i>Scandinavian Audiology</i> , 1994 , 23, 99-104		25
1	Genome-Wide Association Identifies the First Risk Loci for Psychosis in Alzheimer Disease		1