

Sudipta Das

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6787077/sudipta-das-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19

papers

730

citations

12

h-index

19

g-index

19

ext. papers

776

ext. citations

4.2

avg, IF

3.53

L-index

#	Paper	IF	Citations
19	Exploring aggregation-induced emission through tuning of ligand structure for picomolar detection of pyrene. <i>Journal of Molecular Recognition</i> , 2019 , 32, e2771	2.6	2
18	A unique benzimidazole-naphthalene hybrid molecule for independent detection of Zn and N ions: Experimental and theoretical investigations. <i>Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy</i> , 2019 , 209, 170-185	4.4	13
17	Visible Light Excitable SCN ⁻ Selective Fluorescence Probe Derived from Thiophene. <i>Chinese Journal of Chemistry</i> , 2015 , 33, 1173-1177	4.9	6
16	2-(2-Pyridyl) benzimidazole-based ternary Mn(II) complex as an arsenate selective turn-on fluorescence probe: ppb level determination and cell imaging studies. <i>New Journal of Chemistry</i> , 2014 , 38, 2744	3.6	9
15	A visible light excitable pyrene-Naphthalene conjugate for ON fluorescence sensing of histidine in living cells. <i>RSC Advances</i> , 2014 , 4, 7495	3.7	7
14	pH-controlled solid-phase enrichment of Mn(II): confirmation of the structure of the extracted ternary Mn(II) complex by single crystal X-ray structure analysis. <i>Desalination and Water Treatment</i> , 2014 , 52, 6069-6078		1
13	Fluorescent probes for selective determination of trace level Al ³⁺ : recent developments and future prospects. <i>Analytical Methods</i> , 2013 , 5, 6262	3.2	141
12	An INHIBIT logic gate from a thiophene derivative using iron and zinc ions as the input: tuning the efficiency on moving from naphthalene to anthracene to pyrene for the green luminescent detection of the intracellular iron. <i>Dalton Transactions</i> , 2013 , 42, 16387-95	4.3	12
11	Naphthalene based highly selective OFF-ON-OFF type fluorescent probe for Al ³⁺ and NO ₂ ⁻ ions for living cell imaging at physiological pH. <i>Inorganica Chimica Acta</i> , 2013 , 398, 64-71	2.7	30
10	Ratiometric fluorescence sensing and intracellular imaging of Al ³⁺ ions driven by an intramolecular excimer formation of a pyrimidine-pyrene scaffold. <i>Dalton Transactions</i> , 2013 , 42, 4757-63	4.3	84
9	A naphthalene-thiophene hybrid molecule as a fluorescent AND logic gate with Zn ²⁺ and OAc ⁻ ions as inputs: cell imaging and computational studies. <i>Dalton Transactions</i> , 2013 , 42, 6708-15	4.3	29
8	THIOCYANATE-ASSISTED SOLID PHASE ENRICHMENT OF Cu(II) USING PYRIDINE DERIVATIVE WITH X-RAY CRYSTALLOGRAPHIC EVIDENCE. <i>Chemical Engineering Communications</i> , 2013 , 200, 638-654	2.2	1
7	Interaction of a naphthalene based fluorescent probe with Al ³⁺ : experimental and computational studies. <i>Analytical Methods</i> , 2012 , 4, 3620	3.2	21
6	A naphthalene exciplex based Al ³⁺ selective on-type fluorescent probe for living cells at the physiological pH range: experimental and computational studies. <i>Analyst, The</i> , 2012 , 137, 2166-75	5	97
5	Cd(II)-triggered excimer-monomer conversion of a pyrene derivative: time dependent red-shift of monomer emission with cell staining application. <i>Analyst, The</i> , 2012 , 137, 3910-3	5	29
4	Thiophene anchored naphthalene derivative: Cr ³⁺ selective turn-on fluorescent probe for living cell imaging. <i>Analytical Methods</i> , 2012 , 4, 2254	3.2	21
3	A naphthalene-based Al ³⁺ selective fluorescent sensor for living cell imaging. <i>Organic and Biomolecular Chemistry</i> , 2011 , 9, 5523-9	3.9	176

LIST OF PUBLICATIONS

- 2 2-(2-Pyridyl) benzimidazole based Co(II) complex as an efficient fluorescent probe for trace level determination of aspartic and glutamic acid in aqueous solution: a displacement approach. *Organic and Biomolecular Chemistry*, **2011**, 9, 7097-104 3.9 43
- 1 pH dependent separation of uranium by chelation chromatography using pyridine 2,6-dimethanol as a chelator: single crystal X-ray structural confirmation of the chelated uranium complex. *Journal of Hazardous Materials*, **2010**, 181, 154-60 12.8 8