

# Tae-Hyun Yoo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6789904/publications.pdf>

Version: 2024-02-01

281  
papers

6,832  
citations

76196

40  
h-index

114278

63  
g-index

284  
all docs

284  
docs citations

284  
times ranked

8763  
citing authors

#	ARTICLE	IF	CITATIONS
1	An increase in red blood cell distribution width from baseline predicts mortality in patients with severe sepsis or septic shock. <i>Critical Care</i> , 2013, 17, R282.	2.5	162
2	Changing prevalence of glomerular diseases in Korean adults: a review of 20 years of experience. <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 2406-2410.	0.4	160
3	Circulating $\beta$ -Klotho Levels in CKD and Relationship to Progression. <i>American Journal of Kidney Diseases</i> , 2013, 61, 899-909.	2.1	151
4	Changes in Causative Organisms and Their Antimicrobial Susceptibilities in Capd Peritonitis: A Single Center's Experience over one Decade. <i>Peritoneal Dialysis International</i> , 2004, 24, 424-432.	1.1	138
5	Red blood cell distribution width is an independent predictor of mortality in acute kidney injury patients treated with continuous renal replacement therapy. <i>Nephrology Dialysis Transplantation</i> , 2012, 27, 589-594.	0.4	137
6	Characterization of ferroptosis in kidney tubular cell death under diabetic conditions. <i>Cell Death and Disease</i> , 2021, 12, 160.	2.7	133
7	Exosome-based delivery of super-repressor $\beta$ -catenin relieves sepsis-associated organ damage and mortality. <i>Science Advances</i> , 2020, 6, eaaz6980.	4.7	132
8	PGC-1 $\alpha$ Protects from Notch-Induced Kidney Fibrosis Development. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 3312-3322.	3.0	127
9	Sphingomyelinase-Like Phosphodiesterase 3b Expression Levels Determine Podocyte Injury Phenotypes in Glomerular Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 133-147.	3.0	119
10	Decreased Circulating C3 Levels and Mesangial C3 Deposition Predict Renal Outcome in Patients with IgA Nephropathy. <i>PLoS ONE</i> , 2012, 7, e40495.	1.1	112
11	Obesity, Metabolic Abnormality, and Progression of CKD. <i>American Journal of Kidney Diseases</i> , 2018, 72, 400-410.	2.1	105
12	Thyroid Hormone Replacement Therapy Attenuates the Decline of Renal Function in Chronic Kidney Disease Patients with Subclinical Hypothyroidism. <i>Thyroid</i> , 2013, 23, 654-661.	2.4	102
13	Preservation of Renal Function by Thyroid Hormone Replacement Therapy in Chronic Kidney Disease Patients with Subclinical Hypothyroidism. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, 2732-2740.	1.8	101
14	An Increase in Mean Platelet Volume from Baseline Is Associated with Mortality in Patients with Severe Sepsis or Septic Shock. <i>PLoS ONE</i> , 2015, 10, e0119437.	1.1	101
15	Renal outcomes in patients with type 2 diabetes with or without coexisting non-diabetic renal disease. <i>Diabetes Research and Clinical Practice</i> , 2011, 92, 198-204.	1.1	95
16	Clinical Features and Outcomes of IgA Nephropathy with Nephrotic Syndrome. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012, 7, 427-436.	2.2	88
17	Indoxyl sulfate (IS)-mediated immune dysfunction provokes endothelial damage in patients with end-stage renal disease (ESRD). <i>Scientific Reports</i> , 2017, 7, 3057.	1.6	87
18	High-Dose Versus Conventional-Dose Continuous Venovenous Hemodiafiltration and Patient and Kidney Survival and Cytokine Removal in Sepsis-Associated Acute Kidney Injury: A Randomized Controlled Trial. <i>American Journal of Kidney Diseases</i> , 2016, 68, 599-608.	2.1	84

#	ARTICLE	IF	CITATIONS
19	Irisin, a novel myokine is an independent predictor for sarcopenia and carotid atherosclerosis in dialysis patients. <i>Atherosclerosis</i> , 2015, 242, 476-482.	0.4	75
20	A population-based approach indicates an overall higher patient mortality with peritoneal dialysis compared to hemodialysis in Korea. <i>Kidney International</i> , 2014, 86, 991-1000.	2.6	74
21	Induction of heme oxygenase-1 protects against podocyte apoptosis under diabetic conditions. <i>Kidney International</i> , 2009, 76, 838-848.	2.6	67
22	Activation of local aldosterone system within podocytes is involved in apoptosis under diabetic conditions. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 297, F1381-F1390.	1.3	66
23	Uric acid is associated with the rate of residual renal function decline in peritoneal dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 3520-3525.	0.4	64
24	Angiotensin II receptor blocker inhibits p27Kip1 expression in glucose-stimulated podocytes and in diabetic glomeruli. <i>Kidney International</i> , 2005, 67, 944-952.	2.6	63
25	Plasma levels of soluble receptor for advanced glycation end products (sRAGE) and proinflammatory ligand for RAGE (EN-RAGE) are associated with carotid atherosclerosis in patients with peritoneal dialysis. <i>Atherosclerosis</i> , 2012, 220, 208-214.	0.4	61
26	Using the Oxford classification of IgA nephropathy to predict long-term outcomes of Henoch-Schönlein purpura nephritis in adults. <i>Modern Pathology</i> , 2014, 27, 972-982.	2.9	59
27	Decreased Circulating Klotho Levels in Patients Undergoing Dialysis and Relationship to Oxidative Stress and Inflammation. <i>Peritoneal Dialysis International</i> , 2015, 35, 43-51.	1.1	59
28	Left atrial volume is an independent predictor of mortality in CAPD patients. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 3732-3739.	0.4	54
29	Interdialytic Weight Gain and Cardiovascular Outcome in Incident Hemodialysis Patients. <i>American Journal of Nephrology</i> , 2014, 39, 427-435.	1.4	54
30	The Prevalence and Management of Anemia in Chronic Kidney Disease Patients: Result from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>Journal of Korean Medical Science</i> , 2017, 32, 249.	1.1	53
31	Colchicine attenuates inflammatory cell infiltration and extracellular matrix accumulation in diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 297, F200-F209.	1.3	52
32	Clinical Implications of Subclinical Hypothyroidism in Continuous Ambulatory Peritoneal Dialysis Patients. <i>American Journal of Nephrology</i> , 2008, 28, 908-913.	1.4	51
33	Urinary Potassium Excretion and Progression of CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019, 14, 330-340.	2.2	50
34	Exosome-based delivery of super-repressor $\beta$ -catenin ameliorates kidney ischemia-reperfusion injury. <i>Kidney International</i> , 2021, 100, 570-584.	2.6	50
35	MCP-1/CCR2 system is involved in high glucose-induced fibronectin and type IV collagen expression in cultured mesangial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F749-F757.	1.3	49
36	A Diet Rich in Vegetables and Fruit and Incident CKD: A Community-Based Prospective Cohort Study. <i>American Journal of Kidney Diseases</i> , 2019, 74, 491-500.	2.1	48

#	ARTICLE	IF	CITATIONS
37	Pathophysiologic Mechanisms and Potential Biomarkers in Diabetic Kidney Disease. <i>Diabetes and Metabolism Journal</i> , 2022, 46, 181-197.	1.8	48
38	High and low sodium intakes are associated with incident chronic kidney disease in patients with normal renal function and hypertension. <i>Kidney International</i> , 2018, 93, 921-931.	2.6	47
39	Clinical outcomes, when matched at presentation, do not vary between adult-onset Henoch-Schönlein purpura nephritis and IgA nephropathy. <i>Kidney International</i> , 2012, 82, 1304-1312.	2.6	46
40	Early Catheter Removal Improves Patient Survival in Peritoneal Dialysis Patients with Fungal Peritonitis: Results of Ninety-Four Episodes of Fungal Peritonitis at a Single Center. <i>Peritoneal Dialysis International</i> , 2011, 31, 60-66.	1.1	44
41	Differential Expression of Nephrin According to Glomerular Size in Early Diabetic Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 2303-2310.	3.0	42
42	Phosphate is a potential biomarker of disease severity and predicts adverse outcomes in acute kidney injury patients undergoing continuous renal replacement therapy. <i>PLoS ONE</i> , 2018, 13, e0191290.	1.1	40
43	Risk of major cardiovascular events among incident dialysis patients: A Korean national population-based study. <i>International Journal of Cardiology</i> , 2015, 198, 95-101.	0.8	39
44	Prognostic Value of Residual Urine Volume, GFR by 24-hour Urine Collection, and eGFR in Patients Receiving Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2017, 12, 426-434.	2.2	39
45	Indoxyl sulfate-induced TNF $\alpha$ is regulated by crosstalk between the aryl hydrocarbon receptor, NF $\kappa$ B, and SOCS2 in human macrophages. <i>FASEB Journal</i> , 2019, 33, 10844-10858.	0.2	39
46	Association of Blood Pressure With the Progression of CKD: Findings From KNOW-CKD Study. <i>American Journal of Kidney Diseases</i> , 2021, 78, 236-245.	2.1	39
47	The monocyte chemoattractant protein-1 (MCP-1)/CCR2 system is involved in peritoneal dialysis-related epithelial-mesenchymal transition of peritoneal mesothelial cells. <i>Laboratory Investigation</i> , 2012, 92, 1698-1711.	1.7	38
48	Smoking, Smoking Cessation, and Progression of Chronic Kidney Disease: Results From KNOW-CKD Study. <i>Nicotine and Tobacco Research</i> , 2021, 23, 92-98.	1.4	38
49	FR167653 inhibits fibronectin expression and apoptosis in diabetic glomeruli and in high-glucose-stimulated mesangial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F595-F604.	1.3	37
50	Vitamin D deficiency is significantly associated with depression in patients with chronic kidney disease. <i>PLoS ONE</i> , 2017, 12, e0171009.	1.1	37
51	Early initiation of continuous renal replacement therapy improves patient survival in severe progressive septic acute kidney injury. <i>Journal of Critical Care</i> , 2012, 27, 743.e9-743.e18.	1.0	36
52	Hyponatremia as a Predictor of Mortality in Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2014, 9, e111373.	1.1	36
53	Circulating Fibroblast Growth Factor-23 Levels are Associated with an Increased Risk of Anemia Development in Patients with Nondialysis Chronic Kidney Disease. <i>Scientific Reports</i> , 2018, 8, 7294.	1.6	36
54	Diastolic Dysfunction Is an Independent Predictor of Cardiovascular Events in Incident Dialysis Patients with Preserved Systolic Function. <i>PLoS ONE</i> , 2015, 10, e0118694.	1.1	35

#	ARTICLE	IF	CITATIONS
55	Change of Nutritional Status Assessed Using Subjective Global Assessment Is Associated With All-Cause Mortality in Incident Dialysis Patients. <i>Medicine (United States)</i> , 2016, 95, e2714.	0.4	35
56	High-protein diet with renal hyperfiltration is associated with rapid decline rate of renal function: a community-based prospective cohort study. <i>Nephrology Dialysis Transplantation</i> , 2019, 35, 98-106.	0.4	34
57	Secondhand Smoke and CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019, 14, 515-522.	2.2	34
58	Alcohol Consumption and Progression of Chronic Kidney Disease: Results From the Korean Cohort Study for Outcome in Patients with Chronic Kidney Disease. <i>Mayo Clinic Proceedings</i> , 2020, 95, 293-305.	1.4	34
59	Good Glycemic Control Is Associated with Better Survival in Diabetic Patients on Peritoneal Dialysis: A Prospective Observational Study. <i>PLoS ONE</i> , 2012, 7, e30072.	1.1	34
60	Warfarin Use in Patients With Atrial Fibrillation Undergoing Hemodialysis. <i>Stroke</i> , 2017, 48, 2472-2479.	1.0	33
61	The MCP-1/CCR2 axis in podocytes is involved in apoptosis induced by diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2012, 17, 1-13.	2.2	32
62	Association Between Serum High-Density Lipoprotein Cholesterol Levels and Progression of Chronic Kidney Disease: Results From the KNOW-CKD. <i>Journal of the American Heart Association</i> , 2019, 8, e011162.	1.6	32
63	Apoptosis occurs differentially according to glomerular size in diabetic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2012, 27, 259-266.	0.4	31
64	Glomerular IgG deposition predicts renal outcome in patients with IgA nephropathy. <i>Modern Pathology</i> , 2016, 29, 743-752.	2.9	31
65	Effects of Coffee Intake on Incident Chronic Kidney Disease: A Community-Based Prospective Cohort Study. <i>American Journal of Medicine</i> , 2018, 131, 1482-1490.e3.	0.6	31
66	Predictive value of mesangial C3 and C4d deposition in IgA nephropathy. <i>Clinical Immunology</i> , 2020, 211, 108331.	1.4	31
67	Progression of Aortic Arch Calcification Over 1 Year Is an Independent Predictor of Mortality in Incident Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2012, 7, e48793.	1.1	31
68	Comparison of the Haas and the Oxford classifications for prediction of renal outcome in patients with IgA nephropathy. <i>Human Pathology</i> , 2014, 45, 236-243.	1.1	30
69	Selective tubular activation of hypoxia-inducible factor-2 $\alpha$ has dual effects on renal fibrosis. <i>Scientific Reports</i> , 2017, 7, 11351.	1.6	30
70	Gamma Linolenic Acid Exerts Anti-Inflammatory and Anti-Fibrotic Effects in Diabetic Nephropathy. <i>Yonsei Medical Journal</i> , 2012, 53, 1165.	0.9	29
71	Leptin/Adiponectin Ratio is an Independent Predictor of Mortality in Nondiabetic Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2013, 33, 67-74.	1.1	29
72	The KNOW-CKD Study: What we have learned about chronic kidney diseases. <i>Kidney Research and Clinical Practice</i> , 2020, 39, 121-135.	0.9	29

#	ARTICLE	IF	CITATIONS
73	Metabolic syndrome predicts mortality in non-diabetic patients on continuous ambulatory peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 599-604.	0.4	28
74	Vitamin D Deficiency Is an Independent Risk Factor for Urinary Tract Infections After Renal Transplants. <i>Medicine (United States)</i> , 2015, 94, e594.	0.4	28
75	Changes in obese metabolic phenotypes over time and risk of incident chronic kidney disease. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 2778-2791.	2.2	28
76	Dietary zinc intake and incident chronic kidney disease. <i>Clinical Nutrition</i> , 2021, 40, 1039-1045.	2.3	28
77	Enhanced glycogen synthase kinase-3 $\beta$ activity mediates podocyte apoptosis under diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2014, 19, 1678-1690.	2.2	27
78	The benefit of specialized team approaches in patients with acute kidney injury undergoing continuous renal replacement therapy: propensity score matched analysis. <i>Critical Care</i> , 2014, 18, 454.	2.5	27
79	Effect of Icodextrin Solution on the Preservation of Residual Renal Function in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2016, 95, e2991.	0.4	27
80	Klotho plays a protective role against glomerular hypertrophy in a cell cycle-dependent manner in diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 315, F791-F805.	1.3	27
81	Measured sodium excretion is associated with CKD progression: results from the KNOW-CKD study. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, 512-519.	0.4	27
82	P-Cadherin is decreased in diabetic glomeruli and in glucose-stimulated podocytes in vivo and in vitro studies. <i>Nephrology Dialysis Transplantation</i> , 2005, 20, 524-531.	0.4	26
83	Creatinine-Cystatin C Ratio and Mortality in Patients Receiving Intensive Care and Continuous Kidney Replacement Therapy: A Retrospective Cohort Study. <i>American Journal of Kidney Diseases</i> , 2021, 77, 509-516.e1.	2.1	26
84	Usefulness of 23S rRNA Amplification by PCR in the Detection of Bacteria in CAPD Peritonitis. <i>American Journal of Nephrology</i> , 2006, 26, 115-120.	1.4	25
85	Clinical implication of crescentic lesions in immunoglobulin A nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2014, 29, 356-364.	0.4	25
86	Association of serum lipid levels over time with survival in incident peritoneal dialysis patients. <i>Journal of Clinical Lipidology</i> , 2017, 11, 945-954.e3.	0.6	25
87	Prevalence of depression and suicidal ideation increases proportionally with renal function decline, beginning from early stages of chronic kidney disease. <i>Medicine (United States)</i> , 2017, 96, e8476.	0.4	25
88	Permissive fluid volume in adult patients undergoing extracorporeal membrane oxygenation treatment. <i>Critical Care</i> , 2018, 22, 270.	2.5	25
89	Diastolic dysfunction is associated with an increased risk of contrast-induced nephropathy: a retrospective cohort study. <i>BMC Nephrology</i> , 2013, 14, 146.	0.8	24
90	Urine output is associated with prognosis in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , 2013, 28, 379-388.	1.0	24

#	ARTICLE	IF	CITATIONS
91	Glycemic Control Modifies Difference in Mortality Risk Between Hemodialysis and Peritoneal Dialysis in Incident Dialysis Patients With Diabetes. <i>Medicine (United States)</i> , 2016, 95, e3118.	0.4	24
92	Electrolyte and mineral disturbances in septic acute kidney injury patients undergoing continuous renal replacement therapy. <i>Medicine (United States)</i> , 2016, 95, e4542.	0.4	24
93	Body mass index is inversely associated with mortality in patients with acute kidney injury undergoing continuous renal replacement therapy. <i>Kidney Research and Clinical Practice</i> , 2017, 36, 39-47.	0.9	24
94	Visceral Fat Thickness Is Associated With Carotid Atherosclerosis in Peritoneal Dialysis Patients. <i>Obesity</i> , 2012, 20, 1301-1307.	1.5	23
95	The Effect of Statin on Epithelial-Mesenchymal Transition in Peritoneal Mesothelial Cells. <i>PLoS ONE</i> , 2014, 9, e109628.	1.1	23
96	Podocyte hypertrophy precedes apoptosis under experimental diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2015, 20, 1056-1071.	2.2	23
97	High dietary phosphorus density is a risk factor for incident chronic kidney disease development in diabetic subjects: a community-based prospective cohort study. <i>American Journal of Clinical Nutrition</i> , 2017, 106, 311-321.	2.2	23
98	The Optimal Blood Pressure Target in Different Dialysis Populations. <i>Scientific Reports</i> , 2018, 8, 14123.	1.6	23
99	Extracellular Fluid Excess Is Significantly Associated With Coronary Artery Calcification in Patients With Chronic Kidney Disease. <i>Journal of the American Heart Association</i> , 2018, 7, .	1.6	23
100	PGC-1 $\alpha$ inhibits the NLRP3 inflammasome via preserving mitochondrial viability to protect kidney fibrosis. <i>Cell Death and Disease</i> , 2022, 13, 31.	2.7	23
101	The impact of dialysis modality on skin hyperpigmentation in haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2009, 24, 2803-2809.	0.4	22
102	Elevated osteoprotegerin is associated with inflammation, malnutrition and new onset cardiovascular events in peritoneal dialysis patients. <i>Atherosclerosis</i> , 2011, 219, 925-930.	0.4	22
103	Non-Dipper Status and Left Ventricular Hypertrophy as Predictors of Incident Chronic Kidney Disease. <i>Journal of Korean Medical Science</i> , 2011, 26, 1185.	1.1	22
104	The impact of pretransplant 25-hydroxy vitamin D deficiency on subsequent graft function: An observational study. <i>BMC Nephrology</i> , 2012, 13, 22.	0.8	22
105	Clinical features and outcomes of focal segmental glomerulosclerosis pathologic variants in Korean adult patients. <i>BMC Nephrology</i> , 2014, 15, 52.	0.8	22
106	The Effect of Specialized Continuous Renal Replacement Therapy Team in Acute Kidney Injury Patients Treatment. <i>Yonsei Medical Journal</i> , 2015, 56, 658.	0.9	22
107	Nonimmunologic targets of immunosuppressive agents in podocytes. <i>Kidney Research and Clinical Practice</i> , 2015, 34, 69-75.	0.9	22
108	Low serum intact parathyroid hormone level is an independent risk factor for overall mortality and major adverse cardiac and cerebrovascular events in incident dialysis patients. <i>Osteoporosis International</i> , 2016, 27, 2717-2726.	1.3	22



#	ARTICLE	IF	CITATIONS
109	Baseline Cardiovascular Characteristics of Adult Patients with Chronic Kidney Disease from the KoreanN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). Journal of Korean Medical Science, 2017, 32, 231.	1.1	22
110	Framingham risk score and risk of incident chronic kidney disease: A community-based prospective cohort study. Kidney Research and Clinical Practice, 2019, 38, 49-59.	0.9	22
111	Electrocardiographic Left Ventricular Hypertrophy and Outcome in Hemodialysis Patients. PLoS ONE, 2012, 7, e35534.	1.1	22
112	The atherogenic index of plasma and the risk of mortality in incident dialysis patients: Results from a nationwide prospective cohort in Korea. PLoS ONE, 2017, 12, e0177499.	1.1	22
113	Optimal Proteinuria Target for Renoprotection in Patients with IgA Nephropathy. PLoS ONE, 2014, 9, e101935.	1.1	21
114	Mean platelet volume is a prognostic factor in patients with acute kidney injury requiring continuous renal replacement therapy. Journal of Critical Care, 2014, 29, 1016-1021.	1.0	21
115	Endothelial Dysfunction Is Associated With Major Adverse Cardiovascular Events in Peritoneal Dialysis Patients. Medicine (United States), 2014, 93, e73.	0.4	21
116	Creatinineâ€“cystatin C ratio and mortality in cancer patients: a retrospective cohort study. Journal of Cachexia, Sarcopenia and Muscle, 2022, 13, 2064-2072.	2.9	21
117	Congenital Nephrogenic Diabetes Insipidus Presented with Bilateral Hydronephrosis: Genetic Analysis of V2R Gene Mutations. Yonsei Medical Journal, 2006, 47, 126.	0.9	20
118	Serum klotho is inversely associated with metabolic syndrome in chronic kidney disease: results from the KNOW-CKD study. BMC Nephrology, 2019, 20, 119.	0.8	20
119	Reduced Residual Renal Function is Associated with Endothelial Dysfunction in Patients Receiving Peritoneal Dialysis. Peritoneal Dialysis International, 2012, 32, 149-158.	1.1	19
120	Can early initiation of continuous renal replacement therapy improve patient survival with septic acute kidney injury when enrolled in early goal-directed therapy?. Journal of Critical Care, 2016, 35, 51-56.	1.0	19
121	Serum adiponectin and proteinâ€“energy wasting in predialysis chronic kidney disease. Nutrition, 2017, 33, 254-260.	1.1	18
122	Urine Osmolality and Renal Outcome in Patients with Chronic Kidney Disease: Results from the KNOW-CKD. Kidney and Blood Pressure Research, 2019, 44, 1089-1100.	0.9	18
123	High muscleâ€“toâ€“fat ratio is associated with lower risk of chronic kidney disease development. Journal of Cachexia, Sarcopenia and Muscle, 2020, 11, 726-734.	2.9	18
124	The Relationship of Initial Transferrin Saturation to Cardiovascular Parameters and Outcomes in Patients Initiating Dialysis. PLoS ONE, 2014, 9, e87231.	1.1	18
125	Canagliflozin protects against cisplatin-induced acute kidney injury by AMPK-mediated autophagy in renal proximal tubular cells. Cell Death Discovery, 2022, 8, 12.	2.0	18
126	Insulin resistance and lower plasma adiponectin increase malignancy risk in nondiabetic continuous ambulatory peritoneal dialysis patients. Metabolism: Clinical and Experimental, 2011, 60, 121-126.	1.5	17



#	ARTICLE	IF	CITATIONS
127	Thyroid Hormone Replacement Reduces The Risk of Cardiovascular Diseases in Diabetic Nephropathy Patients With Subclinical Hypothyroidism. <i>Endocrine Practice</i> , 2018, 24, 265-272.	1.1	17
128	Ambulatory blood pressure variability and risk of cardiovascular events, all-cause mortality, and progression of kidney disease. <i>Journal of Hypertension</i> , 2020, 38, 1712-1721.	0.3	17
129	Identification of novel mutations in Na-Cl cotransporter gene in a Korean patient with atypical Gitelman's syndrome. <i>American Journal of Kidney Diseases</i> , 2003, 42, e26.1-e26.6.	2.1	16
130	Effects of an oral adsorbent on oxidative stress and fibronectin expression in experimental diabetic nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 2134-2141.	0.4	16
131	Local kallikrein-kinin system is involved in podocyte apoptosis under diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2011, 16, 478-490.	2.2	16
132	The impact of low triiodothyronine levels on mortality is mediated by malnutrition and cardiac dysfunction in incident hemodialysis patients. <i>European Journal of Endocrinology</i> , 2013, 169, 409-419.	1.9	16
133	Normal body mass index with central obesity has increased risk of coronary artery calcification in Korean patients with chronic kidney disease. <i>Kidney International</i> , 2016, 90, 1368-1376.	2.6	16
134	Severe vitamin D deficiency is a risk factor for renal hyperfiltration. <i>American Journal of Clinical Nutrition</i> , 2018, 108, 1342-1351.	2.2	16
135	The Effect of Mycophenolate Mofetil versus Cyclosporine as Combination Therapy with Low Dose Corticosteroids in High-risk Patients with Idiopathic Membranous Nephropathy: a Multicenter Randomized Trial. <i>Journal of Korean Medical Science</i> , 2018, 33, e74.	1.1	16
136	Association of smoking with incident CKD risk in the general population: A community-based cohort study. <i>PLoS ONE</i> , 2020, 15, e0238111.	1.1	16
137	The difference between cystatin C- and creatinine-based eGFR is associated with adverse cardiovascular outcome in patients with chronic kidney disease. <i>Atherosclerosis</i> , 2021, 335, 53-61.	0.4	16
138	Stepwise Treatment Using Corticosteroids Alone and in Combination with Cyclosporine in Korean Patients with Idiopathic Membranous Nephropathy. <i>Yonsei Medical Journal</i> , 2013, 54, 973.	0.9	15
139	Periostin-binding DNA aptamer treatment attenuates renal fibrosis under diabetic conditions. <i>Scientific Reports</i> , 2017, 7, 8490.	1.6	15
140	Hyponatremia Predicts New-Onset Cardiovascular Events in Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2015, 10, e0129480.	1.1	15
141	The pattern of choosing dialysis modality and related mortality outcomes in Korea: a national population-based study. <i>Korean Journal of Internal Medicine</i> , 2017, 32, 699-710.	0.7	15
142	High glucose decreases collagenase expression and increases TIMP expression in cultured human peritoneal mesothelial cells. <i>Nephrology Dialysis Transplantation</i> , 2007, 23, 534-541.	0.4	14
143	Which Biomarker is the Best for Predicting Mortality in Incident Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2015, 94, e1636.	0.4	14
144	Low Mitochondrial DNA Copy Number is Associated With Adverse Clinical Outcomes in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2016, 95, e2717.	0.4	14

#	ARTICLE	IF	CITATIONS
145	Extracellular vesicles in kidneys and their clinical potential in renal diseases. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 194-207.	0.9	14
146	Association of white blood cell count with metabolic syndrome in patients undergoing peritoneal dialysis. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 1379-1385.	1.5	13
147	High Peritoneal Transport Status is Not an Independent Risk Factor for High Mortality in Patients Treated with Automated Peritoneal Dialysis. <i>Journal of Korean Medical Science</i> , 2010, 25, 1313.	1.1	13
148	Removal of Kidney Stones by Extracorporeal Shock Wave Lithotripsy Is Associated with Delayed Progression of Chronic Kidney Disease. <i>Yonsei Medical Journal</i> , 2012, 53, 708.	0.9	13
149	A Low Serum Bicarbonate Concentration as a Risk Factor for Mortality in Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2013, 8, e82912.	1.1	13
150	Serum Ferritin Predicts Mortality Regardless of Inflammatory and Nutritional Status in Patients Starting Dialysis: A Prospective Cohort Study. <i>Blood Purification</i> , 2015, 40, 209-217.	0.9	13
151	Changes in geriatric nutritional risk index and risk of major adverse cardiac and cerebrovascular events in incident peritoneal dialysis patients. <i>Kidney Research and Clinical Practice</i> , 2017, 36, 377-386.	0.9	13
152	Micellized Protein Transduction Domain-Bone Morphogenetic Protein-7 Efficiently Blocks Renal Fibrosis Via Inhibition of Transforming Growth Factor- $\beta$ -Mediated Epithelial-Mesenchymal Transition. <i>Frontiers in Pharmacology</i> , 2020, 11, 591275.	1.6	13
153	Sex disparities and adverse cardiovascular and kidney outcomes in patients with chronic kidney disease: results from the KNOW-CKD. <i>Clinical Research in Cardiology</i> , 2021, 110, 1116-1127.	1.5	13
154	Predictors of Mortality in Patients Returning to Dialysis after Allograft Loss. <i>Blood Purification</i> , 2010, 30, 56-63.	0.9	12
155	Low circulating adiponectin levels are associated with insulin resistance in non-obese peritoneal dialysis patients. <i>Endocrine Journal</i> , 2012, 59, 685-695.	0.7	12
156	Immunogenicity of recombinant human erythropoietin in Korea: A two-year cross-sectional study. <i>Biologicals</i> , 2012, 40, 254-261.	0.5	12
157	The effect of renin-angiotensin system blockade on renal protection in chronic kidney disease patients with hyperkalemia. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2014, 15, 491-497.	1.0	12
158	Baseline Chloride Levels are Associated with the Incidence of Contrast-Associated Acute Kidney Injury. <i>Scientific Reports</i> , 2017, 7, 17431.	1.6	12
159	The impact of disease severity on paradoxical association between body mass index and mortality in patients with acute kidney injury undergoing continuous renal replacement therapy. <i>BMC Nephrology</i> , 2018, 19, 32.	0.8	12
160	Relationship between complement deposition and the Oxford classification score and their combined effects on renal outcome in immunoglobulin A nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 2130-2137.	0.4	12
161	Low High-Sensitivity C-Reactive Protein Level in Korean Patients With Chronic Kidney Disease and Its Predictive Significance for Cardiovascular Events, Mortality, and Adverse Kidney Outcomes: Results From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2020, 9, e017980.	1.6	12
162	Systolic blood pressure and chronic kidney disease progression in patients with primary glomerular disease. <i>Journal of Nephrology</i> , 2021, 34, 1057-1067.	0.9	12

#	ARTICLE	IF	CITATIONS
163	Coronary Artery Calcification Score and the Progression of Chronic Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2022, 33, 1590-1601.	3.0	12
164	Insulin resistance is associated with new-onset cardiovascular events in nondiabetic patients undergoing peritoneal dialysis. <i>Kidney Research and Clinical Practice</i> , 2014, 33, 192-198.	0.9	11
165	Double transduction of a Cre/LoxP lentiviral vector: a simple method to generate kidney cell-specific knockdown mice. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, F1060-F1069.	1.3	11
166	Clinical usefulness of the Oxford classification in determining immunosuppressive treatment in IgA nephropathy. <i>Annals of Medicine</i> , 2017, 49, 217-229.	1.5	11
167	Urinary angiotensinogen level is associated with potassium homeostasis and clinical outcome in patients with polycystic kidney disease: a prospective cohort study. <i>BMC Nephrology</i> , 2019, 20, 104.	0.8	11
168	Association of Reproductive Lifespan Duration and Chronic Kidney Disease in Postmenopausal Women. <i>Mayo Clinic Proceedings</i> , 2020, 95, 2621-2632.	1.4	11
169	Association of Longitudinal Trajectories of Systolic BP with Risk of Incident CKD: Results from the Korean Genome and Epidemiology Study. <i>Journal of the American Society of Nephrology: JASN</i> , 2020, 31, 2133-2144.	3.0	11
170	Left Atrial Enlargement Is Associated with a Rapid Decline in Residual Renal Function in ESRD Patients on Peritoneal Dialysis. <i>Journal of the American Society of Echocardiography</i> , 2012, 25, 421-427.	1.2	10
171	Immunoglobulin A Nephropathy Associated with <i>Plasmodium falciparum</i> Malaria. <i>Journal of Korean Medical Science</i> , 2012, 27, 446.	1.1	10
172	Estimating Average Glucose Levels from Glycated Albumin in Patients with End-Stage Renal Disease. <i>Yonsei Medical Journal</i> , 2012, 53, 578.	0.9	10
173	Rifampicin-Induced Minimal Change Disease Is Improved after Cessation of Rifampicin without Steroid Therapy. <i>Yonsei Medical Journal</i> , 2015, 56, 582.	0.9	10
174	Association between Vitamin D Deficiency and Anemia in Patients with End-Stage Renal Disease: A Cross-Sectional Study. <i>Yonsei Medical Journal</i> , 2016, 57, 1159.	0.9	10
175	Periostin-Binding DNA Aptamer Treatment Ameliorates Peritoneal Dialysis-Induced Peritoneal Fibrosis. <i>Molecular Therapy - Nucleic Acids</i> , 2017, 7, 396-407.	2.3	10
176	Association Between Nocturnal Blood Pressure Dipping and Chronic Kidney Disease Among Patients With Controlled Office Blood Pressure. <i>American Journal of Hypertension</i> , 2021, 34, 821-830.	1.0	10
177	The Effect of HMG-CoA Reductase Inhibitor on Insulin Resistance in Patients Undergoing Peritoneal Dialysis. <i>Cardiovascular Drugs and Therapy</i> , 2012, 26, 501-509.	1.3	9
178	Effect of Peritoneal Dialysis Modality on the 1-Year Rate of Decline of Residual Renal Function. <i>Yonsei Medical Journal</i> , 2014, 55, 141.	0.9	9
179	Low Serum Bicarbonate Predicts Residual Renal Function Loss in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2015, 94, e1276.	0.4	9
180	The locally activated renin-angiotensin system is involved in albumin permeability in glomerular endothelial cells under high glucose conditions. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, gfw089.	0.4	9

#	ARTICLE	IF	CITATIONS
181	Hypoadiponectinemia and the presence of metabolic syndrome in patients with chronic kidney disease: results from the KNOW-CKD study. <i>Diabetology and Metabolic Syndrome</i> , 2016, 8, 75.	1.2	9
182	Preoperative Low Serum Bicarbonate Levels Predict Acute Kidney Injury After Cardiac Surgery. <i>Medicine (United States)</i> , 2016, 95, e3216.	0.4	9
183	Circulating CD89-IgA complex does not predict deterioration of kidney function in Korean patients with IgA nephropathy. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017, 56, 75-85.	1.4	9
184	Carbohydrate-Rich Diet Is Associated with Increased Risk of Incident Chronic Kidney Disease in Non-Diabetic Subjects. <i>Journal of Clinical Medicine</i> , 2019, 8, 793.	1.0	9
185	Soluble Urokinase-type Plasminogen Activator Receptor, Changes of 24-hour Blood Pressure, and Progression of Chronic Kidney Disease. <i>Journal of the American Heart Association</i> , 2021, 10, e017225.	1.6	9
186	Inflammation Alters Relationship Between High-density Lipoprotein Cholesterol and Cardiovascular Risk in Patients With Chronic Kidney Disease: Results From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2021, 10, e021731.	1.6	9
187	Synergic association of diabetes mellitus and chronic kidney disease with muscle loss and cachexia: results of a 16-year longitudinal follow-up of a community-based prospective cohort study. <i>Aging</i> , 2021, 13, 21941-21961.	1.4	9
188	Application of a bridging ELISA for detection of anti-erythropoietin binding antibodies and a cell-based bioassay for neutralizing antibodies in human sera. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010, 52, 289-293.	1.4	8
189	Relation of Homocysteinemia to Contrast-Induced Nephropathy in Patients Undergoing Percutaneous Coronary Intervention. <i>American Journal of Cardiology</i> , 2011, 108, 1086-1091.	0.7	8
190	Delta neutrophil index is an independent predictor of mortality in septic acute kidney injury patients treated with continuous renal replacement therapy. <i>BMC Nephrology</i> , 2017, 18, 94.	0.8	8
191	The Impact of Diabetes Mellitus on Vascular Biomarkers in Patients with End-Stage Renal Disease. <i>Yonsei Medical Journal</i> , 2017, 58, 75.	0.9	8
192	High serum adiponectin is associated with anemia development in chronic kidney disease: The results from the KNOW-CKD study. <i>Cytokine</i> , 2018, 103, 1-9.	1.4	8
193	The association between socioeconomic disparities and left ventricular hypertrophy in chronic kidney disease: results from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>BMC Nephrology</i> , 2018, 19, 203.	0.8	8
194	Incidence of and risk factors for delayed acute kidney injury in patients undergoing colorectal surgery. <i>American Journal of Surgery</i> , 2019, 218, 907-912.	0.9	8
195	Increased risk of acute kidney injury in coronavirus disease patients with renin-angiotensin-aldosterone-system blockade use: a systematic review and meta-analysis. <i>Scientific Reports</i> , 2021, 11, 13588.	1.6	8
196	Sagittal Abdominal Diameter Is an Independent Predictor of All-Cause and Cardiovascular Mortality in Incident Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2013, 8, e77082.	1.1	8
197	Albuminuria as a Risk Factor for Anemia in Chronic Kidney Disease: Result from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>PLoS ONE</i> , 2015, 10, e0139747.	1.1	8
198	Association of serum mineral parameters with mortality in hemodialysis patients: Data from the Korean end-stage renal disease registry. <i>Kidney Research and Clinical Practice</i> , 2018, 37, 266-276.	0.9	8

#	ARTICLE	IF	CITATIONS
199	Low-density lipoprotein cholesterol levels and adverse clinical outcomes in chronic kidney disease: Results from the KNOW-CKD. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022, 32, 410-419.	1.1	8
200	Renal Manifestations in 2007 Korean Patients with Behçet's Disease. <i>Yonsei Medical Journal</i> , 2013, 54, 189.	0.9	7
201	Changes in Echocardiographic Parameters According to the Rate of Residual Renal Function Decline in Incident Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2015, 94, e427.	0.4	7
202	Neck circumference predicts renal function decline in overweight women. <i>Medicine (United States)</i> , 2016, 95, e4844.	0.4	7
203	Smoking Cessation and Coronary Artery Calcification in CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2021, 16, 870-879.	2.2	7
204	Low bone mineral density is associated with coronary arterial calcification progression and incident cardiovascular events in patients with chronic kidney disease. <i>CKJ: Clinical Kidney Journal</i> , 2022, 15, 119-127.	1.4	7
205	Association Between Longitudinal Blood Pressure Trajectory and the Progression of Chronic Kidney Disease: Results From the KNOW-CKD. <i>Hypertension</i> , 2021, 78, 1355-1364.	1.3	7
206	Association of blood pressure with cardiovascular outcome and mortality: results from the KNOW-CKD study. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, 1722-1730.	0.4	7
207	Association Between Systolic Blood Pressure Variability and Major Adverse Cardiovascular Events in Korean Patients With Chronic Kidney Disease: Findings From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2022, 11, .	1.6	7
208	Association between the ratio of insulin-like growth factor-I to insulin-like growth factor binding protein-3 and inflammation in incident automated peritoneal dialysis patients. <i>Growth Hormone and IGF Research</i> , 2013, 23, 170-174.	0.5	6
209	Increased Dialysate MCP-1 is Associated with Cardiovascular Mortality in Peritoneal Dialysis Patients: A Prospective Observational Study. <i>American Journal of Nephrology</i> , 2014, 40, 291-299.	1.4	6
210	NT-proBNP. <i>Medicine (United States)</i> , 2014, 93, e241.	0.4	6
211	The Incidence of Cardiovascular Events Is Comparable Between Normoalbuminuric and Albuminuric Diabetic Patients With Chronic Kidney Disease. <i>Medicine (United States)</i> , 2016, 95, e3175.	0.4	6
212	The Association of Glomerular Glucocorticoid Receptor Expression with Responsiveness to Corticosteroid Treatment in IgA Nephropathy. <i>American Journal of Nephrology</i> , 2019, 50, 187-195.	1.4	6
213	The effect of interactions between proteinuria, activity of fibroblast growth factor 23 and serum phosphate on renal progression in patients with chronic kidney disease: a result from the KoreaN cohort study for Outcome in patients With Chronic Kidney Disease study. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 438-446.	0.4	6
214	Newly designed Protein Transduction Domain (PTD)-mediated BMP7 is a potential therapeutic for peritoneal fibrosis. <i>Journal of Cellular and Molecular Medicine</i> , 2020, 24, 13507-13522.	1.6	6
215	Urinary chloride concentration and progression of chronic kidney disease: results from the KoreaN cohort study for Outcomes in patients With Chronic Kidney Disease. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, 673-680.	0.4	6
216	Erythropoiesis stimulating agent recommendation model using recurrent neural networks for patient with kidney failure with replacement therapy. <i>Computers in Biology and Medicine</i> , 2021, 137, 104718.	3.9	6

#	ARTICLE	IF	CITATIONS
217	Intensity of statin therapy and renal outcome in chronic kidney disease: Results from the Korean Cohort Study for Outcome in Patients With Chronic Kidney Disease. <i>Kidney Research and Clinical Practice</i> , 2020, 39, 93-102.	0.9	6
218	Ideal cardiovascular health duration and risk of chronic kidney disease and cardiovascular disease. <i>Heart</i> , 2022, 108, 523-528.	1.2	6
219	Membranous glomerulonephritis in a patient with myelodysplastic syndrome-refractory cytopenia with multilineage dysplasia. <i>Kidney Research and Clinical Practice</i> , 2013, 32, 134-137.	0.9	5
220	LDL Cholesterol Affects Clinical Outcomes in Incident Hemodialysis Patients During the Early Stages of Dialysis. <i>Blood Purification</i> , 2014, 38, 131-139.	0.9	5
221	Characteristics and Clinical Outcomes of End-Stage Renal Disease Patients on Peritoneal Dialysis for over 15 Years: A Single-Center Experience. <i>Peritoneal Dialysis International</i> , 2017, 37, 535-541.	1.1	5
222	Determination of the optimal target level of proteinuria in the management of patients with glomerular diseases by using different definitions of proteinuria. <i>Medicine (United States)</i> , 2017, 96, e8154.	0.4	5
223	Snoring and incident chronic kidney disease: a community-based prospective cohort study. <i>BMJ Open</i> , 2019, 9, e030671.	0.8	5
224	Lower serum beta-2 microglobulin levels are associated with worse survival in incident peritoneal dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 138-145.	0.4	5
225	Mediation of the relationship between proteinuria and serum phosphate: Insight from the KNOW-CKD study. <i>PLoS ONE</i> , 2020, 15, e0235077.	1.1	5
226	CD71 mesangial IgA1 receptor and the progression of IgA nephropathy. <i>Translational Research</i> , 2021, 230, 34-43.	2.2	5
227	Mixed- versus predilution hemodiafiltration effects on convection volume and small and middle molecule clearance in hemodialysis patients: a prospective randomized controlled trial. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 445-456.	0.9	5
228	Increased Risk of Chronic Kidney Disease Associated With Weight Gain in Healthy Adults: Insight From Metabolic Profiles and Body Composition. <i>Frontiers in Medicine</i> , 2021, 8, 705881.	1.2	5
229	Fatty liver associated with metabolic derangement in patients with chronic kidney disease: A controlled attenuation parameter study. <i>Kidney Research and Clinical Practice</i> , 2017, 36, 48-57.	0.9	5
230	Cumulative fluid balance and mortality in elderly patients with acute kidney injury requiring continuous renal-replacement therapy: a multicenter prospective cohort study. <i>Kidney Research and Clinical Practice</i> , 2020, 39, 414-425.	0.9	5
231	Preoperative Ionized Magnesium Levels and Risk of Acute Kidney Injury After Cardiac Surgery. <i>American Journal of Kidney Diseases</i> , 2022, 80, 629-637.e1.	2.1	5
232	Trends in the association between body mass index and blood pressure among 19-year-old men in Korea from 2003 to 2017. <i>Scientific Reports</i> , 2022, 12, 6767.	1.6	5
233	Association of coronary artery calcium with adverse cardiovascular outcomes and death in patients with chronic kidney disease: results from the KNOW-CKD. <i>Nephrology Dialysis Transplantation</i> , 2023, 38, 712-721.	0.4	5
234	Is it Beneficial to Convert to a Neutral-pH Bicarbonate/Lactate-Buffered PD Solution in Long-Term CAPD Patients? A Single-Center Prospective Study. <i>Peritoneal Dialysis International</i> , 2015, 35, 366-369.	1.1	4



#	ARTICLE	IF	CITATIONS
235	Low Dentin Matrix Protein 1 Is Associated With Incident Cardiovascular Events in Peritoneal Dialysis Patients. <i>Journal of Bone and Mineral Research</i> , 2016, 31, 2149-2158.	3.1	4
236	Efficacy and safety of a balanced salt solution versus a 0.9% saline infusion for the prevention of contrast-induced acute kidney injury (BASIC trial): a study protocol for a randomized controlled trial. <i>Trials</i> , 2017, 18, 461.	0.7	4
237	Cystatin C is Better than Serum Creatinine for Estimating Glomerular Filtration Rate to Detect Osteopenia in Chronic Kidney Disease Patients. <i>Yonsei Medical Journal</i> , 2017, 58, 380.	0.9	4
238	Effect of Psychosocial Distress on the Rate of Kidney Function Decline. <i>Journal of General Internal Medicine</i> , 2021, 36, 2966-2974.	1.3	4
239	A case of membranous nephropathy as a manifestation of graft-versus-host disease. <i>Kidney Research and Clinical Practice</i> , 2013, 32, 39-42.	0.9	3
240	Differential effects of arterial stiffness and fluid overload on blood pressure according to renal function in patients at risk for cardiovascular disease. <i>Hypertension Research</i> , 2019, 42, 341-353.	1.5	3
241	Prognostic value of soluble ST2 and soluble LR11 on mortality and cardiovascular events in peritoneal dialysis patients. <i>BMC Nephrology</i> , 2020, 21, 228.	0.8	3
242	Physical performance and chronic kidney disease development in elderly adults: results from a nationwide cohort study. <i>Aging</i> , 2020, 12, 17393-17417.	1.4	3
243	The fate of manuscripts rejected from <i>Kidney Research and Clinical Practice</i> . <i>Kidney Research and Clinical Practice</i> , 2020, 39, 230-231.	0.9	3
244	Incidence of Acute Kidney Injury after Adrenalectomy in Patients with Primary Aldosteronism. <i>Electrolyte and Blood Pressure</i> , 2019, 17, 45.	0.6	3
245	Dialysis Adequacy and Risk of Dementia in Elderly Hemodialysis Patients. <i>Frontiers in Medicine</i> , 2021, 8, 769490.	1.2	3
246	Colchicine use and the risk of CKD progression: a multicentre nested case-control study. <i>Rheumatology</i> , 2022, 61, 4314-4323.	0.9	3
247	Body weight fluctuation is associated with rapid kidney function decline. <i>Obesity</i> , 2022, 30, 257-267.	1.5	3
248	Effect of baseline glomerular filtration rate on renal function following stenting for atherosclerotic renal artery stenosis. <i>Scandinavian Journal of Urology and Nephrology</i> , 2010, 44, 169-174.	1.4	2
249	End-Stage Renal Disease Impairs the Multidirectional Movements of the Common Carotid Artery: Assessment Using Dimensional Speckle-Tracking Carotid Strain Ultrasonography. <i>Journal of Cardiovascular Imaging</i> , 2018, 26, 155.	0.2	2
250	Indoxyl Sulfate-Mediated Metabolic Alteration of Transcriptome Signatures in Monocytes of Patients with End-Stage Renal Disease (ESRD). <i>Toxins</i> , 2020, 12, 621.	1.5	2
251	Efficacy and Safety of a Balanced Salt Solution Versus a 0.9% Saline Infusion for the Prevention of Contrast-Induced Acute Kidney Injury After Contrast-Enhanced Computed Tomography. <i>Kidney Medicine</i> , 2020, 2, 189-195.	1.0	2
252	P1374DEVELOPMENT OF HEMOGLOBIN PREDICTION AND ERYTHROCYTE STIMULATING AGENT RECOMMENDATION ALGORITHM (HPERA) USING RECURRENT NEURAL NETWORK IN END-STAGE KIDNEY DISEASE PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.4	2



#	ARTICLE	IF	CITATIONS
253	Effect of Steroid Replacement on Long-Term Kidney Function in Patients With Adrenal Insufficiency. <i>Endocrine Practice</i> , 2022, 28, 384-390.	1.1	2
254	External validation of the international prediction tool in Korean patients with immunoglobulin A nephropathy. <i>Kidney Research and Clinical Practice</i> , 2022, 41, 556-566.	0.9	2
255	Minimal Change Disease in Systemic Lupus: Another Renal Manifestation of Lupus?. <i>The Ewha Medical Journal</i> , 2013, 36, 139.	0.1	1
256	Renal Parenchymal Malakoplakia with Acute Interstitial Nephritis Presented with Acute Kidney Injury. <i>The Ewha Medical Journal</i> , 2015, 38, 36.	0.1	1
257	P0930SERUM LEVELS OF OSTEOPROTEGERIN ARE ASSOCIATED WITH OBESITY IN CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.4	1
258	Septicemia, necrotizing fasciitis, and peritonitis due to <i>Vibrio vulnificus</i> treated with early use of polymyxin B hemoperfusion in a patient undergoing CAPD: a case report. <i>BMC Nephrology</i> , 2020, 21, 127.	0.8	1
259	Glomerular subepithelial microparticles - a footprint for podocyte injury. <i>Ultrastructural Pathology</i> , 2021, 45, 236-242.	0.4	1
260	Urinary angiotensinogen as a marker of elevated blood pressure in patients with chronic kidney disease. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 541-543.	0.7	1
261	KRCP's past and future path. <i>Kidney Research and Clinical Practice</i> , 2020, 39, 233-235.	0.9	1
262	Effect of ambulatory blood pressure monitoring guided antihypertensive treatment on renal progression in patients with chronic kidney disease: a randomized comparative study. <i>Journal of Hypertension</i> , 2021, 39, 325-332.	0.3	1
263	Clinical Remission of Renal Amyloidosis after Autologous Peripheral Blood Stem Cell Transplantation. <i>The Ewha Medical Journal</i> , 2013, 36, S25.	0.1	0
264	Syndrome of Inappropriate Secretion of Antidiuretic Hormone after Lung Transplantation. <i>The Ewha Medical Journal</i> , 2014, 37, S41.	0.1	0
265	SP215LOW PRE-OPERATIVE SERUM BICARBONATE LEVELS PREDICT ACUTE KIDNEY INJURY AFTER CARDIAC SURGERY. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i158-i158.	0.4	0
266	MP288ASSOCIATION OF CENTRAL BLOOD PRESSURE WITH MARKERS FOR TARGET ORGAN DAMAGE IN CHRONIC KIDNEY DISEASE PATIENTS: COMPARISON BETWEEN CENTRAL AND PERIPHERAL BLOOD PRESSURE. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i434-i435.	0.4	0
267	In Reply to "Focus Instead on Determining Lower Limit of Continuous Renal Replacement Therapy". <i>American Journal of Kidney Diseases</i> , 2017, 69, 706-707.	2.1	0
268	SP325SERUM ALBUMIN-TO-GLOBULIN RATIO IS AN INDEPENDENT PREDICTOR FOR INCIDENT CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
269	FP021INSULIN-LIKE GROWTH FACTOR BINDING PROTEIN 3 SPECIFIC DNA APTAMER ATTENUATED RENAL TUBULAR FIBROSIS. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0
270	SP227PREOPERATIVE URINE pH IS ASSOCIATED WITH ACUTE KIDNEY INJURY AFTER CARDIOVASCULAR SURGERY. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.4	0

#	ARTICLE	IF	CITATIONS
271	SP045HIGH RENIN-ALDOSTERONE RATIO AND LATERALIZATION INDEX ARE ASSOCIATED WITH ADVERSE RENAL OUTCOMES IN PRIMARY ALDOSTERONISM PATIENTS WHO UNDERWENT ADRENALECTOMY. Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
272	SP302LOW DIETARY ZINC INTAKE IS ASSOCIATED WITH AN INCREASED RISK OF INCIDENT CHRONIC KIDNEY DISEASE DEVELOPMENT. Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
273	FP381ASSOCIATION BETWEEN UNINTENDED WEIGHT LOSS AND INCIDENT CHRONIC KIDNEY DISEASE IN PEOPLE WITH METABOLIC SYNDROME: A COMMUNITY-BASED COHORT STUDY. Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
274	FP400SMOKING, SMOKING CESSATION, AND PROGRESSION OF CHRONIC KIDNEY DISEASE: From the KoreaN cohort study for Outcome in patients With Chronic Kidney Disease (KNOW-CKD). Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
275	FP430SNORING IS ASSOCIATED WITH INCIDENT CHRONIC KIDNEY DISEASE DEVELOPMENT: A COMMUNITY-BASED PROSPECTIVE COHORT STUDY. Nephrology Dialysis Transplantation, 2019, 34, .	0.4	0
276	P0856LOW BONE MASS IS SIGNIFICANTLY ASSOCIATED WITH AN INCREASED RISK OF ANEMIA IN PATIENTS WITH CKD. Nephrology Dialysis Transplantation, 2020, 35, .	0.4	0
277	P0850SERUM OSTEOPROTEGERIN LEVELS ARE ASSOCIATED WITH AN INCREASED RISK OF DEVELOPING ANEMIA IN PATIENTS WITH NON DIALYSIS CHRONIC KIDNEY DISEASE. Nephrology Dialysis Transplantation, 2020, 35, .	0.4	0
278	P0881ANEMIA SIGNIFICANTLY INCREASES RISK OF OSTEOPOROSIS IN PATIENTS WITH NON-DIALYSIS CHRONIC KIDNEY DISEASE. Nephrology Dialysis Transplantation, 2020, 35, .	0.4	0
279	A new opportunity for Kidney Research and Clinical Practice to make a great leap forward as a Science Citation Index Expanded journal. Kidney Research and Clinical Practice, 2021, 40, 1-5.	0.9	0
280	Association between the transtubular potassium gradient and progression of chronic kidney disease: results from KNOW-CKD. Journal of Nephrology, 2021, 34, 2063-2072.	0.9	0
281	Reduction in proteinuria after immunosuppressive therapy and long-term kidney outcomes in patients with immunoglobulin A nephropathy. Korean Journal of Internal Medicine, 2021, 36, 1169-1180.	0.7	0