

Tae-Hyun Yoo

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6789904/tae-hyun-yoo-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

255
papers

4,544
citations

35
h-index

52
g-index

284
ext. papers

5,712
ext. citations

4.5
avg, IF

5.15
L-index

#	Paper	IF	Citations
255	Changing prevalence of glomerular diseases in Korean adults: a review of 20 years of experience. <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 2406-10	4.3	119
254	An increase in red blood cell distribution width from baseline predicts mortality in patients with severe sepsis or septic shock. <i>Critical Care</i> , 2013 , 17, R282	10.8	117
253	Circulating ßklotho levels in CKD and relationship to progression. <i>American Journal of Kidney Diseases</i> , 2013 , 61, 899-909	7.4	116
252	Changes in Causative Organisms and Their Antimicrobial Susceptibilities in Capd Peritonitis: A Single Center's Experience over one Decade. <i>Peritoneal Dialysis International</i> , 2004 , 24, 424-432	2.8	111
251	Red blood cell distribution width is an independent predictor of mortality in acute kidney injury patients treated with continuous renal replacement therapy. <i>Nephrology Dialysis Transplantation</i> , 2012 , 27, 589-94	4.3	104
250	Sphingomyelinase-like phosphodiesterase 3b expression levels determine podocyte injury phenotypes in glomerular disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2015 , 26, 133-47	12.7	97
249	Thyroid hormone replacement therapy attenuates the decline of renal function in chronic kidney disease patients with subclinical hypothyroidism. <i>Thyroid</i> , 2013 , 23, 654-61	6.2	79
248	Preservation of renal function by thyroid hormone replacement therapy in chronic kidney disease patients with subclinical hypothyroidism. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 2732-40	5.6	79
247	An increase in mean platelet volume from baseline is associated with mortality in patients with severe sepsis or septic shock. <i>PLoS ONE</i> , 2015 , 10, e0119437	3.7	78
246	PGC-1 Protects from Notch-Induced Kidney Fibrosis Development. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 3312-3322	12.7	76
245	Renal outcomes in patients with type 2 diabetes with or without coexisting non-diabetic renal disease. <i>Diabetes Research and Clinical Practice</i> , 2011 , 92, 198-204	7.4	76
244	Decreased circulating C3 levels and mesangial C3 deposition predict renal outcome in patients with IgA nephropathy. <i>PLoS ONE</i> , 2012 , 7, e40495	3.7	73
243	Obesity, Metabolic Abnormality, and Progression of CKD. <i>American Journal of Kidney Diseases</i> , 2018 , 72, 400-410	7.4	67
242	Clinical features and outcomes of IgA nephropathy with nephrotic syndrome. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 427-36	6.9	67
241	High-Dose Versus Conventional-Dose Continuous Venovenous Hemodiafiltration and Patient and Kidney Survival and Cytokine Removal in Sepsis-Associated Acute Kidney Injury: A Randomized Controlled Trial. <i>American Journal of Kidney Diseases</i> , 2016 , 68, 599-608	7.4	64
240	Indoxyl sulfate (IS)-mediated immune dysfunction provokes endothelial damage in patients with end-stage renal disease (ESRD). <i>Scientific Reports</i> , 2017 , 7, 3057	4.9	61
239	Exosome-based delivery of super-repressor IBTrelieves sepsis-associated organ damage and mortality. <i>Science Advances</i> , 2020 , 6, eaaz6980	14.3	58

238	A population-based approach indicates an overall higher patient mortality with peritoneal dialysis compared to hemodialysis in Korea. <i>Kidney International</i> , 2014 , 86, 991-1000	9.9	57
237	Induction of heme oxygenase-1 protects against podocyte apoptosis under diabetic conditions. <i>Kidney International</i> , 2009 , 76, 838-48	9.9	57
236	Irisin, a novel myokine is an independent predictor for sarcopenia and carotid atherosclerosis in dialysis patients. <i>Atherosclerosis</i> , 2015 , 242, 476-82	3.1	55
235	Angiotensin II receptor blocker inhibits p27Kip1 expression in glucose-stimulated podocytes and in diabetic glomeruli. <i>Kidney International</i> , 2005 , 67, 944-52	9.9	55
234	Uric acid is associated with the rate of residual renal function decline in peritoneal dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 3520-5	4.3	54
233	Activation of local aldosterone system within podocytes is involved in apoptosis under diabetic conditions. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 297, F1381-90	4.3	54
232	Plasma levels of soluble receptor for advanced glycation end products (sRAGE) and proinflammatory ligand for RAGE (EN-RAGE) are associated with carotid atherosclerosis in patients with peritoneal dialysis. <i>Atherosclerosis</i> , 2012 , 220, 208-14	3.1	50
231	Colchicine attenuates inflammatory cell infiltration and extracellular matrix accumulation in diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 297, F200-9	4.3	46
230	Using the Oxford classification of IgA nephropathy to predict long-term outcomes of Henoch-Schönlein purpura nephritis in adults. <i>Modern Pathology</i> , 2014 , 27, 972-82	9.8	44
229	Left atrial volume is an independent predictor of mortality in CAPD patients. <i>Nephrology Dialysis Transplantation</i> , 2011 , 26, 3732-9	4.3	41
228	Clinical implications of subclinical hypothyroidism in continuous ambulatory peritoneal dialysis patients. <i>American Journal of Nephrology</i> , 2008 , 28, 908-13	4.6	41
227	Decreased circulating klotho levels in patients undergoing dialysis and relationship to oxidative stress and inflammation. <i>Peritoneal Dialysis International</i> , 2015 , 35, 43-51	2.8	40
226	Interdialytic weight gain and cardiovascular outcome in incident hemodialysis patients. <i>American Journal of Nephrology</i> , 2014 , 39, 427-35	4.6	40
225	MCP-1/CCR2 system is involved in high glucose-induced fibronectin and type IV collagen expression in cultured mesangial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2008 , 295, F749-57	4.3	39
224	Differential expression of nephrin according to glomerular size in early diabetic kidney disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 2303-10	12.7	38
223	The monocyte chemoattractant protein-1 (MCP-1)/CCR2 system is involved in peritoneal dialysis-related epithelial-mesenchymal transition of peritoneal mesothelial cells. <i>Laboratory Investigation</i> , 2012 , 92, 1698-711	5.9	37
222	Clinical outcomes, when matched at presentation, do not vary between adult-onset Henoch-Schönlein purpura nephritis and IgA nephropathy. <i>Kidney International</i> , 2012 , 82, 1304-12	9.9	36
221	Early catheter removal improves patient survival in peritoneal dialysis patients with fungal peritonitis: results of ninety-four episodes of fungal peritonitis at a single center. <i>Peritoneal Dialysis International</i> , 2011 , 31, 60-6	2.8	35

220	Risk of major cardiovascular events among incident dialysis patients: A Korean national population-based study. <i>International Journal of Cardiology</i> , 2015 , 198, 95-101	3.2	34
219	The Prevalence and Management of Anemia in Chronic Kidney Disease Patients: Result from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>Journal of Korean Medical Science</i> , 2017 , 32, 249-256	4.7	34
218	FR167653 inhibits fibronectin expression and apoptosis in diabetic glomeruli and in high-glucose-stimulated mesangial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2008 , 295, F595-604	4.3	33
217	High and low sodium intakes are associated with incident chronic kidney disease in patients with normal renal function and hypertension. <i>Kidney International</i> , 2018 , 93, 921-931	9.9	32
216	Prognostic Value of Residual Urine Volume, GFR by 24-hour Urine Collection, and eGFR in Patients Receiving Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2017 , 12, 426-434	6.9	30
215	The MCP-1/CCR2 axis in podocytes is involved in apoptosis induced by diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2012 , 17, 1-13	5.4	30
214	A Diet Rich in Vegetables and Fruit and Incident CKD: A Community-Based Prospective Cohort Study. <i>American Journal of Kidney Diseases</i> , 2019 , 74, 491-500	7.4	29
213	Diastolic dysfunction is an independent predictor of cardiovascular events in incident dialysis patients with preserved systolic function. <i>PLoS ONE</i> , 2015 , 10, e0118694	3.7	29
212	Early initiation of continuous renal replacement therapy improves patient survival in severe progressive septic acute kidney injury. <i>Journal of Critical Care</i> , 2012 , 27, 743.e9-18	4	27
211	Apoptosis occurs differentially according to glomerular size in diabetic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2012 , 27, 259-66	4.3	27
210	Urinary Potassium Excretion and Progression of CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019 , 14, 330-340	6.9	26
209	Comparison of the Haas and the Oxford classifications for prediction of renal outcome in patients with IgA nephropathy. <i>Human Pathology</i> , 2014 , 45, 236-43	3.7	26
208	P-Cadherin is decreased in diabetic glomeruli and in glucose-stimulated podocytes in vivo and in vitro studies. <i>Nephrology Dialysis Transplantation</i> , 2005 , 20, 524-31	4.3	26
207	Progression of aortic arch calcification over 1 year is an independent predictor of mortality in incident peritoneal dialysis patients. <i>PLoS ONE</i> , 2012 , 7, e48793	3.7	26
206	Warfarin Use in Patients With Atrial Fibrillation Undergoing Hemodialysis: A Nationwide Population-Based Study. <i>Stroke</i> , 2017 , 48, 2472-2479	6.7	25
205	Hyponatremia as a predictor of mortality in peritoneal dialysis patients. <i>PLoS ONE</i> , 2014 , 9, e111373	3.7	25
204	Vitamin D deficiency is significantly associated with depression in patients with chronic kidney disease. <i>PLoS ONE</i> , 2017 , 12, e0171009	3.7	25
203	Glomerular IgG deposition predicts renal outcome in patients with IgA nephropathy. <i>Modern Pathology</i> , 2016 , 29, 743-52	9.8	25

202	Good glycemic control is associated with better survival in diabetic patients on peritoneal dialysis: a prospective observational study. <i>PLoS ONE</i> , 2012 , 7, e30072	3.7	24
201	Change of Nutritional Status Assessed Using Subjective Global Assessment Is Associated With All-Cause Mortality in Incident Dialysis Patients. <i>Medicine (United States)</i> , 2016 , 95, e2714	1.8	24
200	Phosphate is a potential biomarker of disease severity and predicts adverse outcomes in acute kidney injury patients undergoing continuous renal replacement therapy. <i>PLoS ONE</i> , 2018 , 13, e0191290	3.7	23
199	Clinical implication of crescentic lesions in immunoglobulin A nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2014 , 29, 356-64	4.3	23
198	Circulating Fibroblast Growth Factor-23 Levels are Associated with an Increased Risk of Anemia Development in Patients with Nondialysis Chronic Kidney Disease. <i>Scientific Reports</i> , 2018 , 8, 7294	4.9	22
197	Enhanced glycogen synthase kinase-3 β activity mediates podocyte apoptosis under diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2014 , 19, 1678-90	5.4	22
196	Visceral fat thickness is associated with carotid atherosclerosis in peritoneal dialysis patients. <i>Obesity</i> , 2012 , 20, 1301-7	8	22
195	Leptin/adiponectin ratio is an independent predictor of mortality in nondiabetic peritoneal dialysis patients. <i>Peritoneal Dialysis International</i> , 2013 , 33, 67-74	2.8	22
194	Usefulness of 23S rRNA amplification by PCR in the detection of bacteria in CAPD peritonitis. <i>American Journal of Nephrology</i> , 2006 , 26, 115-20	4.6	22
193	Vitamin D deficiency is an independent risk factor for urinary tract infections after renal transplants. <i>Medicine (United States)</i> , 2015 , 94, e594	1.8	21
192	Metabolic syndrome predicts mortality in non-diabetic patients on continuous ambulatory peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , 2010 , 25, 599-604	4.3	21
191	Characterization of ferroptosis in kidney tubular cell death under diabetic conditions. <i>Cell Death and Disease</i> , 2021 , 12, 160	9.8	21
190	High-protein diet with renal hyperfiltration is associated with rapid decline rate of renal function: a community-based prospective cohort study. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35, 98-106	4.3	20
189	Indoxyl sulfate-induced TNF- α s regulated by crosstalk between the aryl hydrocarbon receptor, NF- κ B, and SOCS2 in human macrophages. <i>FASEB Journal</i> , 2019 , 33, 10844-10858	0.9	20
188	Diastolic dysfunction is associated with an increased risk of contrast-induced nephropathy: a retrospective cohort study. <i>BMC Nephrology</i> , 2013 , 14, 146	2.7	20
187	Urine output is associated with prognosis in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , 2013 , 28, 379-88	4	20
186	Changes in obese metabolic phenotypes over time and risk of incident chronic kidney disease. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 2778-2791	6.7	19
185	Effects of Coffee Intake on Incident Chronic Kidney Disease: A Community-Based Prospective Cohort Study. <i>American Journal of Medicine</i> , 2018 , 131, 1482-1490.e3	2.4	19

184	Selective tubular activation of hypoxia-inducible factor-2 α has dual effects on renal fibrosis. <i>Scientific Reports</i> , 2017 , 7, 11351	4.9	19
183	Optimal proteinuria target for renoprotection in patients with IgA nephropathy. <i>PLoS ONE</i> , 2014 , 9, e101935	3.7	19
182	The effect of statin on epithelial-mesenchymal transition in peritoneal mesothelial cells. <i>PLoS ONE</i> , 2014 , 9, e109628	3.7	19
181	Reduced residual renal function is associated with endothelial dysfunction in patients receiving peritoneal dialysis. <i>Peritoneal Dialysis International</i> , 2012 , 32, 149-58	2.8	19
180	Podocyte hypertrophy precedes apoptosis under experimental diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2015 , 20, 1056-71	5.4	18
179	Endothelial dysfunction is associated with major adverse cardiovascular events in peritoneal dialysis patients. <i>Medicine (United States)</i> , 2014 , 93, e73	1.8	18
178	Gamma linolenic acid exerts anti-inflammatory and anti-fibrotic effects in diabetic nephropathy. <i>Yonsei Medical Journal</i> , 2012 , 53, 1165-75	3	18
177	The impact of dialysis modality on skin hyperpigmentation in haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 2803-9	4.3	18
176	Low serum intact parathyroid hormone level is an independent risk factor for overall mortality and major adverse cardiac and cerebrovascular events in incident dialysis patients. <i>Osteoporosis International</i> , 2016 , 27, 2717-2726	5.3	18
175	Association of serum lipid levels over time with survival in incident peritoneal dialysis patients. <i>Journal of Clinical Lipidology</i> , 2017 , 11, 945-954.e3	4.9	17
174	Effect of Icodextrin Solution on the Preservation of Residual Renal Function in Peritoneal Dialysis Patients: A Randomized Controlled Study. <i>Medicine (United States)</i> , 2016 , 95, e2991	1.8	17
173	Electrolyte and mineral disturbances in septic acute kidney injury patients undergoing continuous renal replacement therapy. <i>Medicine (United States)</i> , 2016 , 95, e4542	1.8	17
172	The benefit of specialized team approaches in patients with acute kidney injury undergoing continuous renal replacement therapy: propensity score matched analysis. <i>Critical Care</i> , 2014 , 18, 454	10.8	17
171	Prevalence of depression and suicidal ideation increases proportionally with renal function decline, beginning from early stages of chronic kidney disease. <i>Medicine (United States)</i> , 2017 , 96, e8476	1.8	17
170	The impact of pretransplant 25-hydroxy vitamin D deficiency on subsequent graft function: an observational study. <i>BMC Nephrology</i> , 2012 , 13, 22	2.7	17
169	Elevated osteoprotegerin is associated with inflammation, malnutrition and new onset cardiovascular events in peritoneal dialysis patients. <i>Atherosclerosis</i> , 2011 , 219, 925-30	3.1	17
168	Non-dipper status and left ventricular hypertrophy as predictors of incident chronic kidney disease. <i>Journal of Korean Medical Science</i> , 2011 , 26, 1185-90	4.7	17
167	The atherogenic index of plasma and the risk of mortality in incident dialysis patients: Results from a nationwide prospective cohort in Korea. <i>PLoS ONE</i> , 2017 , 12, e0177499	3.7	17

166	Glycemic Control Modifies Difference in Mortality Risk Between Hemodialysis and Peritoneal Dialysis in Incident Dialysis Patients With Diabetes: Results From a Nationwide Prospective Cohort in Korea. <i>Medicine (United States)</i> , 2016 , 95, e3118	1.8	17
165	Framingham risk score and risk of incident chronic kidney disease: A community-based prospective cohort study. <i>Kidney Research and Clinical Practice</i> , 2019 , 38, 49-59	3.6	16
164	Electrocardiographic left ventricular hypertrophy and outcome in hemodialysis patients. <i>PLoS ONE</i> , 2012 , 7, e35534	3.7	16
163	Smoking, Smoking Cessation, and Progression of Chronic Kidney Disease: Results From KNOW-CKD Study. <i>Nicotine and Tobacco Research</i> , 2021 , 23, 92-98	4.9	16
162	High dietary phosphorus density is a risk factor for incident chronic kidney disease development in diabetic subjects: a community-based prospective cohort study. <i>American Journal of Clinical Nutrition</i> , 2017 , 106, 311-321	7	15
161	Secondhand Smoke and CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019 , 14, 515-522	6.9	15
160	Klotho plays a protective role against glomerular hypertrophy in a cell cycle-dependent manner in diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2018 , 315, F791-F805	4.3	15
159	Mean platelet volume is a prognostic factor in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , 2014 , 29, 1016-21	4	15
158	Insulin resistance and lower plasma adiponectin increase malignancy risk in nondiabetic continuous ambulatory peritoneal dialysis patients. <i>Metabolism: Clinical and Experimental</i> , 2011 , 60, 121-6	12.7	15
157	Effects of an oral adsorbent on oxidative stress and fibronectin expression in experimental diabetic nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2010 , 25, 2134-41	4.3	15
156	Association Between Serum High-Density Lipoprotein Cholesterol Levels and Progression of Chronic Kidney Disease: Results From the KNOW-CKD. <i>Journal of the American Heart Association</i> , 2019 , 8, e011162	6	14
155	The impact of low triiodothyronine levels on mortality is mediated by malnutrition and cardiac dysfunction in incident hemodialysis patients. <i>European Journal of Endocrinology</i> , 2013 , 169, 409-19	6.5	14
154	The effect of specialized continuous renal replacement therapy team in acute kidney injury patients treatment. <i>Yonsei Medical Journal</i> , 2015 , 56, 658-65	3	14
153	Body mass index is inversely associated with mortality in patients with acute kidney injury undergoing continuous renal replacement therapy. <i>Kidney Research and Clinical Practice</i> , 2017 , 36, 39-47 ^{3.6}		14
152	Can early initiation of continuous renal replacement therapy improve patient survival with septic acute kidney injury when enrolled in early goal-directed therapy?. <i>Journal of Critical Care</i> , 2016 , 35, 51-6 ⁴		14
151	Nonimmunologic targets of immunosuppressive agents in podocytes. <i>Kidney Research and Clinical Practice</i> , 2015 , 34, 69-75	3.6	13
150	Serum adiponectin and protein-energy wasting in predialysis chronic kidney disease. <i>Nutrition</i> , 2017 , 33, 254-260	4.8	13
149	Baseline Cardiovascular Characteristics of Adult Patients with Chronic Kidney Disease from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>Journal of Korean Medical Science</i> , 2017 , 32, 231-239	4.7	13

148	Which Biomarker is the Best for Predicting Mortality in Incident Peritoneal Dialysis Patients: NT-ProBNP, Cardiac TnT, or hsCRP?: A Prospective Observational Study. <i>Medicine (United States)</i> , 2015 , 94, e1636	1.8	13
147	A low serum bicarbonate concentration as a risk factor for mortality in peritoneal dialysis patients. <i>PLoS ONE</i> , 2013 , 8, e82912	3.7	13
146	Local kallikrein-kinin system is involved in podocyte apoptosis under diabetic conditions. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2011 , 16, 478-90	5.4	13
145	Congenital nephrogenic diabetes insipidus presented with bilateral hydronephrosis: genetic analysis of V2R gene mutations. <i>Yonsei Medical Journal</i> , 2006 , 47, 126-30	3	13
144	Identification of novel mutations in Na-Cl cotransporter gene in a Korean patient with atypical Gitelman's syndrome. <i>American Journal of Kidney Diseases</i> , 2003 , 42, E11-6	7.4	13
143	The KNOW-CKD Study: What we have learned about chronic kidney diseases. <i>Kidney Research and Clinical Practice</i> , 2020 , 39, 121-135	3.6	13
142	Predictive value of mesangial C3 and C4d deposition in IgA nephropathy. <i>Clinical Immunology</i> , 2020 , 211, 108331	9	13
141	Permissive fluid volume in adult patients undergoing extracorporeal membrane oxygenation treatment. <i>Critical Care</i> , 2018 , 22, 270	10.8	13
140	Normal body mass index with central obesity has increased risk of coronary artery calcification in Korean patients with chronic kidney disease. <i>Kidney International</i> , 2016 , 90, 1368-1376	9.9	12
139	Periostin-binding DNA aptamer treatment attenuates renal fibrosis under diabetic conditions. <i>Scientific Reports</i> , 2017 , 7, 8490	4.9	12
138	Low circulating adiponectin levels are associated with insulin resistance in non-obese peritoneal dialysis patients. <i>Endocrine Journal</i> , 2012 , 59, 685-95	2.9	12
137	The Optimal Blood Pressure Target in Different Dialysis Populations. <i>Scientific Reports</i> , 2018 , 8, 14123	4.9	12
136	THYROID HORMONE REPLACEMENT REDUCES THE RISK OF CARDIOVASCULAR DISEASES IN DIABETIC NEPHROPATHY PATIENTS WITH SUBCLINICAL HYPOTHYROIDISM. <i>Endocrine Practice</i> , 2018 , 24, 265-272	3.2	12
135	Extracellular Fluid Excess Is Significantly Associated With Coronary Artery Calcification in Patients With Chronic Kidney Disease. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	12
134	Exosome-based delivery of super-repressor IBA1 ameliorates kidney ischemia-reperfusion injury. <i>Kidney International</i> , 2021 , 100, 570-584	9.9	12
133	Serum klotho is inversely associated with metabolic syndrome in chronic kidney disease: results from the KNOW-CKD study. <i>BMC Nephrology</i> , 2019 , 20, 119	2.7	11
132	Clinical features and outcomes of focal segmental glomerulosclerosis pathologic variants in Korean adult patients. <i>BMC Nephrology</i> , 2014 , 15, 52	2.7	11
131	Predictors of mortality in patients returning to dialysis after allograft loss. <i>Blood Purification</i> , 2010 , 30, 56-63	3.1	11

130	The relationship of initial transferrin saturation to cardiovascular parameters and outcomes in patients initiating dialysis. <i>PLoS ONE</i> , 2014 , 9, e87231	3.7	11
129	Hyponatremia Predicts New-Onset Cardiovascular Events in Peritoneal Dialysis Patients. <i>PLoS ONE</i> , 2015 , 10, e0129480	3.7	11
128	The Effect of Mycophenolate Mofetil versus Cyclosporine as Combination Therapy with Low Dose Corticosteroids in High-risk Patients with Idiopathic Membranous Nephropathy: a Multicenter Randomized Trial. <i>Journal of Korean Medical Science</i> , 2018 , 33, e74	4.7	10
127	Baseline Chloride Levels are Associated with the Incidence of Contrast-Associated Acute Kidney Injury. <i>Scientific Reports</i> , 2017 , 7, 17431	4.9	10
126	Immunogenicity of recombinant human erythropoietin in Korea: a two-year cross-sectional study. <i>Biologicals</i> , 2012 , 40, 254-61	1.8	10
125	Removal of kidney stones by extracorporeal shock wave lithotripsy is associated with delayed progression of chronic kidney disease. <i>Yonsei Medical Journal</i> , 2012 , 53, 708-14	3	10
124	High glucose decreases collagenase expression and increases TIMP expression in cultured human peritoneal mesothelial cells. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 534-41	4.3	10
123	The pattern of choosing dialysis modality and related mortality outcomes in Korea: a national population-based study. <i>Korean Journal of Internal Medicine</i> , 2017 , 32, 699-710	2.5	10
122	Measured sodium excretion is associated with CKD progression: results from the KNOW-CKD study. <i>Nephrology Dialysis Transplantation</i> , 2021 , 36, 512-519	4.3	10
121	Periostin-Binding DNA Aptamer Treatment Ameliorates Peritoneal Dialysis-Induced Peritoneal Fibrosis. <i>Molecular Therapy - Nucleic Acids</i> , 2017 , 7, 396-407	10.7	9
120	High muscle-to-fat ratio is associated with lower risk of chronic kidney disease development. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2020 , 11, 726-734	10.3	9
119	Double transduction of a Cre/LoxP lentiviral vector: a simple method to generate kidney cell-specific knockdown mice. <i>American Journal of Physiology - Renal Physiology</i> , 2015 , 309, F1060-9	4.3	9
118	The effect of renin-angiotensin system blockade on renal protection in chronic kidney disease patients with hyperkalemia. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2014 , 15, 491-7		9
117	Left atrial enlargement is associated with a rapid decline in residual renal function in ESRD patients on peritoneal dialysis. <i>Journal of the American Society of Echocardiography</i> , 2012 , 25, 421-7	5.8	9
116	Immunoglobulin A nephropathy associated with Plasmodium falciparum malaria. <i>Journal of Korean Medical Science</i> , 2012 , 27, 446-9	4.7	9
115	Estimating average glucose levels from glycated albumin in patients with end-stage renal disease. <i>Yonsei Medical Journal</i> , 2012 , 53, 578-86	3	9
114	Alcohol Consumption and Progression of Chronic Kidney Disease: Results From the Korean Cohort Study for Outcome in Patients with Chronic Kidney Disease. <i>Mayo Clinic Proceedings</i> , 2020 , 95, 293-305	6.4	9
113	Hypoadiponectinemia and the presence of metabolic syndrome in patients with chronic kidney disease: results from the KNOW-CKD study. <i>Diabetology and Metabolic Syndrome</i> , 2016 , 8, 75	5.6	9

112	Preoperative Low Serum Bicarbonate Levels Predict Acute Kidney Injury After Cardiac Surgery. <i>Medicine (United States)</i> , 2016 , 95, e3216	1.8	8
111	Changes in geriatric nutritional risk index and risk of major adverse cardiac and cerebrovascular events in incident peritoneal dialysis patients. <i>Kidney Research and Clinical Practice</i> , 2017 , 36, 377-386	3.6	8
110	Serum Ferritin Predicts Mortality Regardless of Inflammatory and Nutritional Status in Patients Starting Dialysis: A Prospective Cohort Study. <i>Blood Purification</i> , 2015 , 40, 209-17	3.1	8
109	The effect of HMG-CoA reductase inhibitor on insulin resistance in patients undergoing peritoneal dialysis. <i>Cardiovascular Drugs and Therapy</i> , 2012 , 26, 501-9	3.9	8
108	High peritoneal transport status is not an independent risk factor for high mortality in patients treated with automated peritoneal dialysis. <i>Journal of Korean Medical Science</i> , 2010 , 25, 1313-7	4.7	8
107	Application of a bridging ELISA for detection of anti-erythropoietin binding antibodies and a cell-based bioassay for neutralizing antibodies in human sera. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2010 , 52, 289-93	3.5	8
106	Creatinine-Cystatin C Ratio and Mortality in Patients Receiving Intensive Care and Continuous Kidney Replacement Therapy: A Retrospective Cohort Study. <i>American Journal of Kidney Diseases</i> , 2021 , 77, 509-516.e1	7.4	8
105	The Impact of Diabetes Mellitus on Vascular Biomarkers in Patients with End-Stage Renal Disease. <i>Yonsei Medical Journal</i> , 2017 , 58, 75-81	3	7
104	The impact of disease severity on paradoxical association between body mass index and mortality in patients with acute kidney injury undergoing continuous renal replacement therapy. <i>BMC Nephrology</i> , 2018 , 19, 32	2.7	7
103	Circulating CD89-IgA complex does not predict deterioration of kidney function in Korean patients with IgA nephropathy. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017 , 56, 75-85	5.9	7
102	Clinical usefulness of the Oxford classification in determining immunosuppressive treatment in IgA nephropathy. <i>Annals of Medicine</i> , 2017 , 49, 217-229	1.5	7
101	Rifampicin-induced minimal change disease is improved after cessation of rifampicin without steroid therapy. <i>Yonsei Medical Journal</i> , 2015 , 56, 582-5	3	7
100	Low Serum Bicarbonate Predicts Residual Renal Function Loss in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2015 , 94, e1276	1.8	7
99	Insulin resistance is associated with new-onset cardiovascular events in nondiabetic patients undergoing peritoneal dialysis. <i>Kidney Research and Clinical Practice</i> , 2014 , 33, 192-8	3.6	7
98	Renal manifestations in 2007 Korean patients with Behçet's disease. <i>Yonsei Medical Journal</i> , 2013 , 54, 189-96	3	7
97	Stepwise treatment using corticosteroids alone and in combination with cyclosporine in Korean patients with idiopathic membranous nephropathy. <i>Yonsei Medical Journal</i> , 2013 , 54, 973-82	3	7
96	Relation of homocysteinemia to contrast-induced nephropathy in patients undergoing percutaneous coronary intervention. <i>American Journal of Cardiology</i> , 2011 , 108, 1086-91	3	7
95	Association of white blood cell count with metabolic syndrome in patients undergoing peritoneal dialysis. <i>Metabolism: Clinical and Experimental</i> , 2009 , 58, 1379-85	12.7	7

94	Sagittal abdominal diameter is an independent predictor of all-cause and cardiovascular mortality in incident peritoneal dialysis patients. <i>PLoS ONE</i> , 2013 , 8, e77082	3.7	7
93	Albuminuria as a Risk Factor for Anemia in Chronic Kidney Disease: Result from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>PLoS ONE</i> , 2015 , 10, e0139747	3.7	7
92	Low High-Sensitivity C-Reactive Protein Level in Korean Patients With Chronic Kidney Disease and Its Predictive Significance for Cardiovascular Events, Mortality, and Adverse Kidney Outcomes: Results From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2020 , 9, e017980	6	7
91	Association between Vitamin D Deficiency and Anemia in Patients with End-Stage Renal Disease: A Cross-Sectional Study. <i>Yonsei Medical Journal</i> , 2016 , 57, 1159-64	3	7
90	Dietary zinc intake and incident chronic kidney disease. <i>Clinical Nutrition</i> , 2021 , 40, 1039-1045	5.9	7
89	Delta neutrophil index is an independent predictor of mortality in septic acute kidney injury patients treated with continuous renal replacement therapy. <i>BMC Nephrology</i> , 2017 , 18, 94	2.7	6
88	Changes in echocardiographic parameters according to the rate of residual renal function decline in incident peritoneal dialysis patients. <i>Medicine (United States)</i> , 2015 , 94, e427	1.8	6
87	NT-proBNP: is it a more significant risk factor for mortality than troponin T in incident hemodialysis patients?. <i>Medicine (United States)</i> , 2014 , 93, e241	1.8	6
86	Association between the ratio of insulin-like growth factor-I to insulin-like growth factor binding protein-3 and inflammation in incident automated peritoneal dialysis patients. <i>Growth Hormone and IGF Research</i> , 2013 , 23, 170-4	2	6
85	Increased dialysate MCP-1 is associated with cardiovascular mortality in peritoneal dialysis patients: a prospective observational study. <i>American Journal of Nephrology</i> , 2014 , 40, 291-9	4.6	6
84	Association of serum mineral parameters with mortality in hemodialysis patients: Data from the Korean end-stage renal disease registry. <i>Kidney Research and Clinical Practice</i> , 2018 , 37, 266-276	3.6	6
83	Low Mitochondrial DNA Copy Number is Associated With Adverse Clinical Outcomes in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , 2016 , 95, e2717	1.8	6
82	Association of Blood Pressure With the Progression of CKD: Findings From KNOW-CKD Study. <i>American Journal of Kidney Diseases</i> , 2021 , 78, 236-245	7.4	6
81	Urinary angiotensinogen level is associated with potassium homeostasis and clinical outcome in patients with polycystic kidney disease: a prospective cohort study. <i>BMC Nephrology</i> , 2019 , 20, 104	2.7	5
80	High serum adiponectin is associated with anemia development in chronic kidney disease: The results from the KNOW-CKD study. <i>Cytokine</i> , 2018 , 103, 1-9	4	5
79	The Incidence of Cardiovascular Events Is Comparable Between Normoalbuminuric and Albuminuric Diabetic Patients With Chronic Kidney Disease. <i>Medicine (United States)</i> , 2016 , 95, e3175	1.8	5
78	Urine Osmolality and Renal Outcome in Patients with Chronic Kidney Disease: Results from the KNOW-CKD. <i>Kidney and Blood Pressure Research</i> , 2019 , 44, 1089-1100	3.1	5
77	Membranous glomerulonephritis in a patient with myelodysplastic syndrome-refractory cytopenia with multilineage dysplasia. <i>Kidney Research and Clinical Practice</i> , 2013 , 32, 134-7	3.6	5

76	LDL cholesterol affects clinical outcomes in incident hemodialysis patients during the early stages of dialysis. <i>Blood Purification</i> , 2014 , 38, 131-9	3.1	5
75	Fatty liver associated with metabolic derangement in patients with chronic kidney disease: A controlled attenuation parameter study. <i>Kidney Research and Clinical Practice</i> , 2017 , 36, 48-57	3.6	5
74	Micellized Protein Transduction Domain-Bone Morphogenetic Protein-7 Efficiently Blocks Renal Fibrosis Via Inhibition of Transforming Growth Factor-Beta-Mediated Epithelial-Mesenchymal Transition. <i>Frontiers in Pharmacology</i> , 2020 , 11, 591275	5.6	5
73	Ambulatory blood pressure variability and risk of cardiovascular events, all-cause mortality, and progression of kidney disease. <i>Journal of Hypertension</i> , 2020 , 38, 1712-1721	1.9	5
72	Characteristics and Clinical Outcomes of End-Stage Renal Disease Patients on Peritoneal Dialysis for Over 15 Years: A Single-Center Experience. <i>Peritoneal Dialysis International</i> , 2017 , 37, 535-541	2.8	4
71	Carbohydrate-Rich Diet Is Associated with Increased Risk of Incident Chronic Kidney Disease in Non-Diabetic Subjects. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	4
70	Incidence of and risk factors for delayed acute kidney injury in patients undergoing colorectal surgery. <i>American Journal of Surgery</i> , 2019 , 218, 907-912	2.7	4
69	Mediation of the relationship between proteinuria and serum phosphate: Insight from the KNOW-CKD study. <i>PLoS ONE</i> , 2020 , 15, e0235077	3.7	4
68	Neck circumference predicts renal function decline in overweight women: A community-based prospective cohort study. <i>Medicine (United States)</i> , 2016 , 95, e4844	1.8	4
67	The Association of Glomerular Glucocorticoid Receptor Expression with Responsiveness to Corticosteroid Treatment in IgA Nephropathy. <i>American Journal of Nephrology</i> , 2019 , 50, 187-195	4.6	4
66	Determination of the optimal target level of proteinuria in the management of patients with glomerular diseases by using different definitions of proteinuria. <i>Medicine (United States)</i> , 2017 , 96, e8154	1.8	4
65	Is It Beneficial to Convert to a Neutral-pH Bicarbonate/Lactate-Buffered PD Solution in Long-Term CAPD Patients? A Single-Center Prospective Study. <i>Peritoneal Dialysis International</i> , 2015 , 35, 366-9	2.8	4
64	Effect of peritoneal dialysis modality on the 1-year rate of decline of residual renal function. <i>Yonsei Medical Journal</i> , 2014 , 55, 141-8	3	4
63	Association of Longitudinal Trajectories of Systolic BP with Risk of Incident CKD: Results from the Korean Genome and Epidemiology Study. <i>Journal of the American Society of Nephrology: JASN</i> , 2020 , 31, 2133-2144	12.7	4
62	Association of smoking with incident CKD risk in the general population: A community-based cohort study. <i>PLoS ONE</i> , 2020 , 15, e0238111	3.7	4
61	Increased risk of acute kidney injury in coronavirus disease patients with renin-angiotensin-aldosterone-system blockade use: a systematic review and meta-analysis. <i>Scientific Reports</i> , 2021 , 11, 13588	4.9	4
60	The locally activated renin-angiotensin system is involved in albumin permeability in glomerular endothelial cells under high glucose conditions. <i>Nephrology Dialysis Transplantation</i> , 2017 , 32, 61-72	4.3	4
59	Relationship between complement deposition and the Oxford classification score and their combined effects on renal outcome in immunoglobulin A nephropathy. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35, 2103-2137	4.3	4

58	Soluble Urokinase-Type Plasminogen Activator Receptor, Changes of 24-Hour Blood Pressure, and Progression of Chronic Kidney Disease. <i>Journal of the American Heart Association</i> , 2021 , 10, e017225	6	4
57	Low Dentin Matrix Protein 1 Is Associated With Incident Cardiovascular Events in Peritoneal Dialysis Patients. <i>Journal of Bone and Mineral Research</i> , 2016 , 31, 2149-2158	6.3	3
56	A case of membranous nephropathy as a manifestation of graft-versus-host disease. <i>Kidney Research and Clinical Practice</i> , 2013 , 32, 39-42	3.6	3
55	Intensity of statin therapy and renal outcome in chronic kidney disease: Results from the Korean Cohort Study for Outcome in Patients With Chronic Kidney Disease. <i>Kidney Research and Clinical Practice</i> , 2020 , 39, 93-102	3.6	3
54	The fate of manuscripts rejected from. <i>Kidney Research and Clinical Practice</i> , 2020 , 39, 230-231	3.6	3
53	Snoring and incident chronic kidney disease: a community-based prospective cohort study. <i>BMJ Open</i> , 2019 , 9, e030671	3	3
52	Lower serum beta-2 microglobulin levels are associated with worse survival in incident peritoneal dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 138-145	4.3	3
51	Differential effects of arterial stiffness and fluid overload on blood pressure according to renal function in patients at risk for cardiovascular disease. <i>Hypertension Research</i> , 2019 , 42, 341-353	4.7	3
50	The effect of interactions between proteinuria, activity of fibroblast growth factor 23 and serum phosphate on renal progression in patients with chronic kidney disease: a result from the KoreaN cohort study for Outcome in patients With Chronic Kidney Disease study. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35, 438-446	4.3	3
49	Urinary chloride concentration and progression of chronic kidney disease: results from the KoreaN cohort study for Outcomes in patients With Chronic Kidney Disease. <i>Nephrology Dialysis Transplantation</i> , 2021 , 36, 673-680	4.3	3
48	Severe vitamin D deficiency is a risk factor for renal hyperfiltration. <i>American Journal of Clinical Nutrition</i> , 2018 , 108, 1342-1351	7	3
47	Pathophysiologic Mechanisms and Potential Biomarkers in Diabetic Kidney Disease.. <i>Diabetes and Metabolism Journal</i> , 2022 , 46, 181-197	5	3
46	Prognostic value of soluble ST2 and soluble LR11 on mortality and cardiovascular events in peritoneal dialysis patients. <i>BMC Nephrology</i> , 2020 , 21, 228	2.7	2
45	P1374DEVELOPMENT OF HEMOGLOBIN PREDICTION AND ERYTHROCYTE STIMULATING AGENT RECOMMENDATION ALGORITHM (HPERA) USING RECURRENT NEURAL NETWORK IN END-STAGE KIDNEY DISEASE PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35,	4.3	2
44	Efficacy and safety of a balanced salt solution versus a 0.9% saline infusion for the prevention of contrast-induced acute kidney injury (BASIC trial): a study protocol for a randomized controlled trial. <i>Trials</i> , 2017 , 18, 461	2.8	2
43	Cystatin C is Better than Serum Creatinine for Estimating Glomerular Filtration Rate to Detect Osteopenia in Chronic Kidney Disease Patients. <i>Yonsei Medical Journal</i> , 2017 , 58, 380-387	3	2
42	The association between socioeconomic disparities and left ventricular hypertrophy in chronic kidney disease: results from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>BMC Nephrology</i> , 2018 , 19, 203	2.7	2
41	Physical performance and chronic kidney disease development in elderly adults: results from a nationwide cohort study. <i>Aging</i> , 2020 , 12, 17393-17417	5.6	2

40	Incidence of Acute Kidney Injury after Adrenalectomy in Patients with Primary Aldosteronism. <i>Electrolyte and Blood Pressure</i> , 2019 , 17, 45-53	1.1	2
39	Newly designed Protein Transduction Domain (PTD)-mediated BMP-7 is a potential therapeutic for peritoneal fibrosis. <i>Journal of Cellular and Molecular Medicine</i> , 2020 , 24, 13507-13522	5.6	2
38	Association of Reproductive Lifespan Duration and Chronic Kidney Disease in Postmenopausal Women. <i>Mayo Clinic Proceedings</i> , 2020 , 95, 2621-2632	6.4	2
37	Efficacy and Safety of a Balanced Salt Solution Versus a 0.9% Saline Infusion for the Prevention of Contrast-Induced Acute Kidney Injury After Contrast-Enhanced Computed Tomography. <i>Kidney Medicine</i> , 2020 , 2, 189-195	2.8	2
36	Sex disparities and adverse cardiovascular and kidney outcomes in patients with chronic kidney disease: results from the KNOW-CKD. <i>Clinical Research in Cardiology</i> , 2021 , 110, 1116-1127	6.1	2
35	Extracellular vesicles in kidneys and their clinical potential in renal diseases. <i>Kidney Research and Clinical Practice</i> , 2021 , 40, 194-207	3.6	2
34	Systolic blood pressure and chronic kidney disease progression in patients with primary glomerular disease. <i>Journal of Nephrology</i> , 2021 , 34, 1057-1067	4.8	2
33	Association Between Nocturnal Blood Pressure Dipping and Chronic Kidney Disease Among Patients With Controlled Office Blood Pressure. <i>American Journal of Hypertension</i> , 2021 , 34, 821-830	2.3	2
32	Low bone mineral density is associated with coronary arterial calcification progression and incident cardiovascular events in patients with chronic kidney disease.. <i>CKJ: Clinical Kidney Journal</i> , 2022 , 15, 1194-127	4.5	2
31	Association Between Longitudinal Blood Pressure Trajectory and the Progression of Chronic Kidney Disease: Results From the KNOW-CKD. <i>Hypertension</i> , 2021 , 78, 1355-1364	8.5	2
30	Erythropoiesis stimulating agent recommendation model using recurrent neural networks for patient with kidney failure with replacement therapy. <i>Computers in Biology and Medicine</i> , 2021 , 137, 104718	7.18	2
29	Renal Parenchymal Malakoplakia with Acute Interstitial Nephritis Presented with Acute Kidney Injury. <i>The Ewha Medical Journal</i> , 2015 , 38, 36	0.1	1
28	Minimal Change Disease in Systemic Lupus: Another Renal Manifestation of Lupus?. <i>The Ewha Medical Journal</i> , 2013 , 36, 139	0.1	1
27	Effect of baseline glomerular filtration rate on renal function following stenting for atherosclerotic renal artery stenosis. <i>Scandinavian Journal of Urology and Nephrology</i> , 2010 , 44, 169-74		1
26	Canagliflozin protects against cisplatin-induced acute kidney injury by AMPK-mediated autophagy in renal proximal tubular cells.. <i>Cell Death Discovery</i> , 2022 , 8, 12	6.9	1
25	Dialysis Adequacy and Risk of Dementia in Elderly Hemodialysis Patients.. <i>Frontiers in Medicine</i> , 2021 , 8, 769490	4.9	1
24	PGC-1 α inhibits the NLRP3 inflammasome via preserving mitochondrial viability to protect kidney fibrosis.. <i>Cell Death and Disease</i> , 2022 , 13, 31	9.8	1
23	Low-density lipoprotein cholesterol levels and adverse clinical outcomes in chronic kidney disease: Results from the KNOW-CKD. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 ,	4.5	1

22	Cumulative fluid balance and mortality in elderly patients with acute kidney injury requiring continuous renal-replacement therapy: a multicenter prospective cohort study. <i>Kidney Research and Clinical Practice</i> , 2020 , 39, 414-425	3.6	1
21	P0930SERUM LEVELS OF OSTEOPROTEGERIN ARE ASSOCIATED WITH OBESITY IN CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35,	4.3	1
20	Glomerular subepithelial microparticles - a footprint for podocyte injury. <i>Ultrastructural Pathology</i> , 2021 , 45, 236-242	1.3	1
19	CD71 mesangial IgA1 receptor and the progression of IgA nephropathy. <i>Translational Research</i> , 2021 , 230, 34-43	11	1
18	End-Stage Renal Disease Impairs the Multidirectional Movements of the Common Carotid Artery: Assessment Using Dimensional Speckle-Tracking Carotid Strain Ultrasonography. <i>Journal of Cardiovascular Imaging</i> , 2018 , 26, 155-164	1.3	1
17	Inflammation Alters Relationship Between High-Density Lipoprotein Cholesterol and Cardiovascular Risk in Patients With Chronic Kidney Disease: Results From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2021 , 10, e021731	6	1
16	Increased Risk of Chronic Kidney Disease Associated With Weight Gain in Healthy Adults: Insight From Metabolic Profiles and Body Composition. <i>Frontiers in Medicine</i> , 2021 , 8, 705881	4.9	1
15	The difference between cystatin C- and creatinine-based eGFR is associated with adverse cardiovascular outcome in patients with chronic kidney disease. <i>Atherosclerosis</i> , 2021 , 335, 53-61	3.1	1
14	Effect of Psychosocial Distress on the Rate of Kidney Function Decline. <i>Journal of General Internal Medicine</i> , 2021 , 36, 2966-2974	4	1
13	Smoking Cessation and Coronary Artery Calcification in CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2021 , 16, 870-879	6.9	0
12	Mixed- versus predilution hemodiafiltration effects on convection volume and small and middle molecule clearance in hemodialysis patients: a prospective randomized controlled trial. <i>Kidney Research and Clinical Practice</i> , 2021 , 40, 445-456	3.6	0
11	Synergic association of diabetes mellitus and chronic kidney disease with muscle loss and cachexia: results of a 16-year longitudinal follow-up of a community-based prospective cohort study. <i>Aging</i> , 2021 , 13, 21941-21961	5.6	0
10	Trends in the association between body mass index and blood pressure among 19-year-old men in Korea from 2003 to 2017.. <i>Scientific Reports</i> , 2022 , 12, 6767	4.9	0
9	In Reply to Focus Instead on Determining Lower Limit of Continuous Renal Replacement TherapyS <i>American Journal of Kidney Diseases</i> , 2017 , 69, 706-707	7.4	
8	Septicemia, necrotizing fasciitis, and peritonitis due to <i>Vibrio vulnificus</i> treated with early use of polymyxin B hemoperfusion in a patient undergoing CAPD: a case report. <i>BMC Nephrology</i> , 2020 , 21, 127	2.7	
7	Syndrome of Inappropriate Secretion of Antidiuretic Hormone after Lung Transplantation. <i>The Ewha Medical Journal</i> , 2014 , 37, S41	0.1	
6	Clinical Remission of Renal Amyloidosis after Autologous Peripheral Blood Stem Cell Transplantation. <i>The Ewha Medical Journal</i> , 2013 , 36, S25	0.1	
5	Effect of ambulatory blood pressure monitoring guided antihypertensive treatment on renal progression in patients with chronic kidney disease: a randomized comparative study. <i>Journal of Hypertension</i> , 2021 , 39, 325-332	1.9	

4	Association between the transtubular potassium gradient and progression of chronic kidney disease: results from KNOW-CKD. <i>Journal of Nephrology</i> , 2021 , 34, 2063-2072	4.8
3	SP215LOW PRE-OPERATIVE SERUM BICARBONATE LEVELS PREDICT ACUTE KIDNEY INJURY AFTER CARDIAC SURGERY. <i>Nephrology Dialysis Transplantation</i> , 2016 , 31, i158-i158	4.3
2	MP288ASSOCIATION OF CENTRAL BLOOD PRESSURE WITH MARKERS FOR TARGET ORGAN DAMAGE IN CHRONIC KIDNEY DISEASE PATIENTS: COMPARISON BETWEEN CENTRAL AND PERIPHERAL BLOOD PRESSURE. <i>Nephrology Dialysis Transplantation</i> , 2016 , 31, i434-i435	4.3
1	Reduction in proteinuria after immunosuppressive therapy and long-term kidney outcomes in patients with immunoglobulin A nephropathy. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 1169-1180	2.5