

Josef KÄhrle

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6792021/publications.pdf>

Version: 2024-02-01

299
papers

15,325
citations

14653

66
h-index

24254

110
g-index

327
all docs

327
docs citations

327
times ranked

12652
citing authors

#	ARTICLE	IF	CITATIONS
1	Perinatal exposure to the thyroperoxidase inhibitors methimazole and amitrole perturbs thyroid hormone system signaling and alters motor activity in rat offspring. <i>Toxicology Letters</i> , 2022, 354, 44-55.	0.8	12
2	100 YEARS OF VITAMIN D: Light and health: a century after the therapeutic use of UV light and vitamin D, hormones advanced medical care. <i>Endocrine Connections</i> , 2022, 11, .	1.9	2
3	Tentative Application of a Streamlined Protocol to Determine Organ-Specific Regulations of Deiodinase 1 and Dehalogenase Activities as Readouts of the Hypothalamus-Pituitary-Thyroid-Periphery-Axis. <i>Frontiers in Toxicology</i> , 2022, 4, 822993.	3.1	3
4	Changes in Thyroid Metabolites after Liothyronine Administration: A Secondary Analysis of Two Clinical Trials That Incorporated Pharmacokinetic Data. <i>Metabolites</i> , 2022, 12, 476.	2.9	0
5	3,5-T2-an Endogenous Thyroid Hormone Metabolite as Promising Lead Substance in Anti-Steatotic Drug Development?. <i>Metabolites</i> , 2022, 12, 582.	2.9	6
6	Obesity and Pregnancy. Guideline of the German Society of Gynecology and Obstetrics (S3-Level, AWMF) Tj ETQq0,0,0 rgBT /Qverlock 1	1.8	10
7	Comparative Analysis of the Effects of Long-Term 3,5-diiodothyronine Treatment on the Murine Hepatic Proteome and Transcriptome Under Conditions of Normal Diet and High-Fat Diet. <i>Thyroid</i> , 2021, 31, 1135-1146.	4.5	7
8	Thyroid hormone system disrupting chemicals. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2021, 35, 101562.	4.7	20
9	Testing for heterotopia formation in rats after developmental exposure to selected inÂvitro inhibitors of thyroperoxidase. <i>Environmental Pollution</i> , 2021, 283, 117135.	7.5	19
10	Selenium in Endocrinologyâ€”Selenoprotein-Related Diseases, Population Studies, and Epidemiological Evidence. <i>Endocrinology</i> , 2021, 162, .	2.8	27
11	Lack of the Thyroid Hormone Transporter Mct8 in Osteoblast and Osteoclast Progenitors Increases Trabecular Bone in Male Mice. <i>Thyroid</i> , 2020, 30, 329-342.	4.5	9
12	Disruption of <sc>BMP</sc> Signaling Prevents Hyperthyroidismâ€”Induced Bone Loss in Male Mice. <i>Journal of Bone and Mineral Research</i> , 2020, 35, 2058-2069.	2.8	13
13	Mass Spectrometry-Based Determination of Thyroid Hormones and Their Metabolites in Endocrine Diagnostics and Biomedical Research â€” Implications for Human Serum Diagnostics. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2020, 128, 358-374.	1.2	4
14	CD5L Constitutes a Novel Biomarker for Integrated Hepatic Thyroid Hormone Action. <i>Thyroid</i> , 2020, 30, 908-923.	4.5	8
15	Endocrine, Metabolic and Pharmacological Effects of Thyronamines (TAM), Thyroacetic Acids (TA) and Thyroid Hormone Metabolites (THM) â€” Evidence from in vitro, Cellular, Experimental Animal and Human Studies. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2020, 128, 401-413.	1.2	10
16	Removing Critical Gaps in Chemical Test Methods by Developing New Assays for the Identification of Thyroid Hormone System-Disrupting Chemicalsâ€”The ATHENA Project. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3123.	4.1	34
17	Adversity Considerations for Thyroid Follicular Cell Hypertrophy and Hyperplasia in Nonclinical Toxicity Studies: Results From the 6th ESTP International Expert Workshop. <i>Toxicologic Pathology</i> , 2020, 48, 920-938.	1.8	12
18	Endocrine Disruptors and Thyroid Function. , 2019, , 787-792.		0

#	ARTICLE	IF	CITATIONS
19	A Thyroid Hormone-Independent Molecular Fingerprint of 3,5-Diiodothyronine Suggests a Strong Relationship with Coffee Metabolism in Humans. <i>Thyroid</i> , 2019, 29, 1743-1754.	4.5	12
20	The Colorful Diversity of Thyroid Hormone Metabolites. <i>European Thyroid Journal</i> , 2019, 8, 115-129.	2.4	55
21	A combined LC-MS/MS and LC-MS3 multi-method for the quantification of iodothyronines in human blood serum. <i>Analytical and Bioanalytical Chemistry</i> , 2019, 411, 5605-5616.	3.7	23
22	3-Iodothyronamine—A Thyroid Hormone Metabolite With Distinct Target Profiles and Mode of Action. <i>Endocrine Reviews</i> , 2019, 40, 602-630.	20.1	38
23	The isoflavones genistein and daidzein increase hepatic concentration of thyroid hormones and affect cholesterol metabolism in middle-aged male rats. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2019, 190, 1-10.	2.5	40
24	The Role of Dickkopf-1 in Thyroid Hormone–Induced Changes of Bone Remodeling in Male Mice. <i>Endocrinology</i> , 2019, 160, 664-674.	2.8	12
25	Association Between 3-Iodothyronamine (T1) Concentrations and Left Ventricular Function in Chronic Heart Failure. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 1232-1238.	3.6	10
26	Aging Alters Phenotypic Traits of Thyroid Dysfunction in Male Mice With Divergent Effects on Complex Systems but Preserved Thyroid Hormone Action in Target Organs. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2019, 74, 1162-1169.	3.6	9
27	3,5-T2—A Janus-Faced Thyroid Hormone Metabolite Exerts Both Canonical T3-Mimetic Endocrine and Intracrine Hepatic Action. <i>Frontiers in Endocrinology</i> , 2019, 10, 787.	3.5	17
28	A combined LC-MS/MS and LC-MS3 multi-method for the quantification of iodothyronines in human blood serum. , 2019, 411, 5605.		1
29	A combined LC-MS/MS and LC-MS3 multi-method for the quantification of iodothyronines in human blood serum. , 2019, 411, 5605.		1
30	In vivo Effects of Repeated Thyronamine Administration in Male C57BL/6J Mice. <i>European Thyroid Journal</i> , 2018, 7, 3-12.	2.4	15
31	Micronutrient status assessment in humans: Current methods of analysis and future trends. <i>TrAC - Trends in Analytical Chemistry</i> , 2018, 102, 110-122.	11.4	24
32	3-Iodothyronamine reduces insulin secretion in vitro via a mitochondrial mechanism. <i>Molecular and Cellular Endocrinology</i> , 2018, 460, 219-228.	3.2	14
33	Relaxin-2 connecting peptide (pro-RLX2) levels in second trimester serum samples to predict preeclampsia. <i>Pregnancy Hypertension</i> , 2018, 11, 124-128.	1.4	6
34	The Effect of High Dose Isoflavone Supplementation on Serum Reverse T3 in Euthyroid Men With Type 2 Diabetes and Post-menopausal Women. <i>Frontiers in Endocrinology</i> , 2018, 9, 698.	3.5	9
35	Thyroid Hormone Metabolism. , 2018, , 420-428.		0
36	Vascular Endothelial Growth Factor (VEGF) Induced Downstream Responses to Transient Receptor Potential Vanilloid 1 (TRPV1) and 3-Iodothyronamine (3-TIAM) in Human Corneal Keratocytes. <i>Frontiers in Endocrinology</i> , 2018, 9, 670.	3.5	16

#	ARTICLE	IF	CITATIONS
37	TRPM8 Activation via 3-Iodothyronamine Blunts VEGF-Induced Transactivation of TRPV1 in Human Uveal Melanoma Cells. <i>Frontiers in Pharmacology</i> , 2018, 9, 1234.	3.5	18
38	Canonical TSH Regulation of Cathepsin-Mediated Thyroglobulin Processing in the Thyroid Gland of Male Mice Requires Taar1 Expression. <i>Frontiers in Pharmacology</i> , 2018, 9, 221.	3.5	22
39	Effects of isoflavones on breast tissue and the thyroid hormone system in humans: a comprehensive safety evaluation. <i>Archives of Toxicology</i> , 2018, 92, 2703-2748.	4.2	62
40	Thyroid Hormones and Derivatives: Endogenous Thyroid Hormones and Their Targets. <i>Methods in Molecular Biology</i> , 2018, 1801, 85-104.	0.9	41
41	Molecular features of the L-type amino acid transporter 2 determine different import and export profiles for thyroid hormones and amino acids. <i>Molecular and Cellular Endocrinology</i> , 2017, 443, 163-174.	3.2	14
42	Sex-specific and inter-individual differences in biomarkers of selenium status identified by a calibrated ELISA for selenoprotein P. <i>Redox Biology</i> , 2017, 11, 403-414.	9.0	79
43	Avoiding the pitfalls when quantifying thyroid hormones and their metabolites using mass spectrometric methods: The role of quality assurance. <i>Molecular and Cellular Endocrinology</i> , 2017, 458, 44-56.	3.2	26
44	Sclerostin Blockade and Zoledronic Acid Improve Bone Mass and Strength in Male Mice With Exogenous Hyperthyroidism. <i>Endocrinology</i> , 2017, 158, 3765-3777.	2.8	20
45	BMPs as new insulin sensitizers: enhanced glucose uptake in mature 3T3-L1 adipocytes via PPAR α and GLUT4 upregulation. <i>Scientific Reports</i> , 2017, 7, 17192.	3.3	43
46	Noncanonical thyroid hormone signaling mediates cardiometabolic effects in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E11323-E11332.	7.1	93
47	Aminoglycoside-driven biosynthesis of selenium-deficient Selenoprotein P. <i>Scientific Reports</i> , 2017, 7, 4391.	3.3	17
48	3-Iodothyronamine Decreases Expression of Genes Involved in Iodide Metabolism in Mouse Thyroids and Inhibits Iodide Uptake in PCCL3 Thyrocytes. <i>Thyroid</i> , 2017, 27, 11-22.	4.5	26
49	Igniting stage two in <i>Endocrine Connections</i> . <i>Endocrine Connections</i> , 2017, 6, E1-E2.	1.9	0
50	Editorial: Get inspired - Lessons learned from evolution of thyroid hormone signaling in developmental processes. <i>Molecular and Cellular Endocrinology</i> , 2017, 459, 1-4.	3.2	2
51	Quantification of Relaxin-2 Connecting Peptide (Pro-RLX2) in Human Blood Samples. <i>Journal of Applied Laboratory Medicine</i> , 2017, 2, 322-334.	1.3	2
52	A validated LC-MS/MS method for cellular thyroid hormone metabolism: Uptake and turnover of mono-iodinated thyroid hormone metabolites by PCCL3 thyrocytes. <i>PLoS ONE</i> , 2017, 12, e0183482.	2.5	17
53	Sex-specific phenotypes of hyperthyroidism and hypothyroidism in aged mice. <i>Biology of Sex Differences</i> , 2017, 8, 38.	4.1	20
54	Avoiding the pitfalls when quantifying thyroid hormones and their metabolites using mass spectrometric methods: The role of quality assurance. , 2017, 458, 44-44.		1

#	ARTICLE	IF	CITATIONS
55	Restoration of type 1 iodothyronine deiodinase expression in renal cancer cells downregulates oncoproteins and affects key metabolic pathways as well as anti-oxidative system. PLoS ONE, 2017, 12, e0190179.	2.5	17
56	Differential Modulation of Adrenergic Receptor Signaling by Octopamine, Tyramine, Phenylethylamine, and 3-Iodothyronamine. , 2016, , 63-81.		2
57	Few Amino Acid Exchanges Expand the Substrate Spectrum of Monocarboxylate Transporter 10*. Molecular Endocrinology, 2016, 30, 796-808.	3.7	17
58	A Nonradioactive DEHAL Assay for Testing Substrates, Inhibitors, and Monitoring Endogenous Activity. Endocrinology, 2016, 157, 4516-4525.	2.8	16
59	Minireview: Insights Into the Structural and Molecular Consequences of the TSH-Î² Mutation C105Vfs114X. Molecular Endocrinology, 2016, 30, 954-964.	3.7	10
60	Sex-specific phenotypes of hyperthyroidism and hypothyroidism in mice. Biology of Sex Differences, 2016, 7, 36.	4.1	34
61	Thyroid hormone status defines brown adipose tissue activity and browning of white adipose tissues in mice. Scientific Reports, 2016, 6, 38124.	3.3	71
62	Silychristin, a Flavonolignan Derived From the Milk Thistle, Is a Potent Inhibitor of the Thyroid Hormone Transporter MCT8. Endocrinology, 2016, 157, 1694-1701.	2.8	54
63	Factors impacting the aminoglycoside-induced UGA stop codon readthrough in selenoprotein translation. Journal of Trace Elements in Medicine and Biology, 2016, 37, 104-110.	3.0	12
64	Thyronamines and Derivatives: Physiological Relevance, Pharmacological Actions, and Future Research Directions. Thyroid, 2016, 26, 1656-1673.	4.5	70
65	Circulating 3-T1AM and 3,5-T2 in Critically Ill Patients: A Cross-Sectional Observational Study. Thyroid, 2016, 26, 1674-1680.	4.5	27
66	Chemical Hybridization of Glucagon and Thyroid Hormone Optimizes Therapeutic Impact for Metabolic Disease. Cell, 2016, 167, 843-857.e14.	28.9	153
67	Efficacy of protocols for induction of chronic hyperthyroidism in male and female mice. Endocrine, 2016, 54, 47-54.	2.3	18
68	Selenium and Endocrine Tissues. , 2016, , 389-400.		2
69	High Variability of Insulin Sensitivity in Closely Related Obese Mouse Inbred Strains. Experimental and Clinical Endocrinology and Diabetes, 2016, 124, 519-528.	1.2	7
70	Selenoprotein Gene Nomenclature. Journal of Biological Chemistry, 2016, 291, 24036-24040.	3.4	207
71	Thyroid hormone and its metabolites in relation to quality of life in patients treated for differentiated thyroid cancer. Clinical Endocrinology, 2016, 85, 781-788.	2.4	41
72	High levels of thyroid-stimulating hormone are associated with aortic wall thickness in the general population. European Radiology, 2016, 26, 4490-4496.	4.5	8

#	ARTICLE	IF	CITATIONS
73	3,5-T2 alters murine genes relevant for xenobiotic, steroid, and thyroid hormone metabolism. <i>Journal of Molecular Endocrinology</i> , 2016, 56, 311-323.	2.5	28
74	Exposure to Thyroid-Disrupting Chemicals: A Transatlantic Call for Action. <i>Thyroid</i> , 2016, 26, 479-480.	4.5	16
75	3-Iodothyronamine increases transient receptor potential melastatin channel 8 (TRPM8) activity in immortalized human corneal epithelial cells. <i>Cellular Signalling</i> , 2016, 28, 136-147.	3.6	41
76	Selenoprotein P and Selenium Distribution in Mammals. , 2016, , 261-274.		8
77	Establishment of an Effective Radioiodide Thyroid Ablation Protocol in Mice. <i>European Thyroid Journal</i> , 2015, 4, 74-80.	2.4	8
78	Differences in Mouse Hepatic Thyroid Hormone Transporter Expression with Age and Hyperthyroidism. <i>European Thyroid Journal</i> , 2015, 4, 81-86.	2.4	22
79	Serum Thyrotropin Concentrations Are Not Associated with the Ankle-Brachial Index: Results from Three Population-Based Studies. <i>European Thyroid Journal</i> , 2015, 4, 101-107.	2.4	3
80	Involvement of the L-Type Amino Acid Transporter Lat2 in the Transport of 3,3- ¹²⁵ I-Diiodothyronine across the Plasma Membrane. <i>European Thyroid Journal</i> , 2015, 4, 42-50.	2.4	22
81	The Multitarget Ligand 3-Iodothyronamine Modulates β -Adrenergic Receptor 2 Signaling. <i>European Thyroid Journal</i> , 2015, 4, 21-29.	2.4	31
82	Lokalisation und Verteilung von Selenoprotein P im humanen Gehirn. <i>Perspectives in Science</i> , 2015, 3, 9-11.	0.6	0
83	Establishment and characterization of a new ELISA for selenoprotein P. <i>Perspectives in Science</i> , 2015, 3, 23-24.	0.6	4
84	Selenium and the thyroid. <i>Current Opinion in Endocrinology, Diabetes and Obesity</i> , 2015, 22, 392-401.	2.3	134
85	A Nonradioactive Uptake Assay for Rapid Analysis of Thyroid Hormone Transporter Function. <i>Endocrinology</i> , 2015, 156, 2739-2745.	2.8	21
86	3-iodothyronamine differentially modulates β -2-adrenergic receptor-mediated signaling. <i>Journal of Molecular Endocrinology</i> , 2015, 54, 205-216.	2.5	54
87	Quantitative Analysis of Thyroid Hormone Metabolites in Cell Culture Samples Using LC-MS/MS. <i>European Thyroid Journal</i> , 2015, 4, 51-58.	2.4	35
88	Testosterone and estradiol treatments differently affect pituitary-thyroid axis and liver deiodinase 1 activity in orchidectomized middle-aged rats. <i>Experimental Gerontology</i> , 2015, 72, 85-98.	2.8	24
89	Trace Amine-Associated Receptor 1 Localization at the Apical Plasma Membrane Domain of Fisher Rat Thyroid Epithelial Cells Is Confined to Cilia. <i>European Thyroid Journal</i> , 2015, 4, 30-41.	2.4	28
90	Urine Metabolomics by 1H-NMR Spectroscopy Indicates Associations between Serum 3,5-T2 Concentrations and Intermediary Metabolism in Euthyroid Humans. <i>European Thyroid Journal</i> , 2015, 4, 92-100.	2.4	32

#	ARTICLE	IF	CITATIONS
91	Nonthyroidal Illness Syndrome in Cardiac Illness Involves Elevated Concentrations of 3,5-Diiodothyronine and Correlates with Atrial Remodeling. <i>European Thyroid Journal</i> , 2015, 4, 129-137.	2.4	67
92	High T3, Low T4 Serum Levels in Mct8 Deficiency Are Not Caused by Increased Hepatic Conversion through Type I Deiodinase. <i>European Thyroid Journal</i> , 2015, 4, 87-91.	2.4	10
93	3,5-Diiodo-L-Thyronine (3,5-T ₂) Exerts Thyromimetic Effects on Hypothalamus-Pituitary-Thyroid Axis, Body Composition, and Energy Metabolism in Male Diet-Induced Obese Mice. <i>Endocrinology</i> , 2015, 156, 389-399.	2.8	97
94	Translating Pharmacological Findings from Hypothyroid Rodents to Euthyroid Humans: Is There a Functional Role of Endogenous 3,5-T ₂ ? <i>Thyroid</i> , 2015, 25, 188-197.	4.5	35
95	Structural Insights Into Thyroid Hormone Transport Mechanisms of the L-Type Amino Acid Transporter 2. <i>Molecular Endocrinology</i> , 2015, 29, 933-942.	3.7	20
96	The Effects of Thyroid Hormones on Gene Expression of Acyl-Coenzyme A Thioesterases in Adipose Tissue and Liver of Mice. <i>European Thyroid Journal</i> , 2015, 4, 59-66.	2.4	12
97	Hyperthyroidism and Hypothyroidism in Male Mice and Their Effects on Bone Mass, Bone Turnover, and the Wnt Inhibitors Sclerostin and Dickkopf-1. <i>Endocrinology</i> , 2015, 156, 3517-3527.	2.8	53
98	Biosynthesis of 3-Iodothyronamine From T4 in Murine Intestinal Tissue. <i>Endocrinology</i> , 2015, 156, 4356-4364.	2.8	63
99	Chronic Kidney Disease Distinctly Affects Relationship Between Selenoprotein P Status and Serum Thyroid Hormone Parameters. <i>Thyroid</i> , 2015, 25, 1091-1096.	4.5	14
100	An Improved Nonradioactive Screening Method Identifies Genistein and Xanthohumol as Potent Inhibitors of Iodothyronine Deiodinases. <i>Thyroid</i> , 2015, 25, 962-968.	4.5	59
101	Thyronamine induces TRPM8 channel activation in human conjunctival epithelial cells. <i>Cellular Signalling</i> , 2015, 27, 315-325.	3.6	43
102	Inverse Agonistic Action of 3-Iodothyronamine at the Human Trace Amine-Associated Receptor 5. <i>PLoS ONE</i> , 2015, 10, e0117774.	2.5	62
103	Quantitative analysis of thyroid hormone metabolites in cell culture samples using LC-MS/MS. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
104	Effects of the thyroid hormone metabolite 3,5-T ₂ on murine heart tissue and the cardiomyocyte cell line H9C2. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
105	Phenotypic and molecular characterization of age-dependent thyroid hormone action in mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
106	Influence of thyroid hormones on brown adipose tissue activity and browning of white adipose tissues in mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
107	Towards the intestinal biosynthesis of 3-iodothyronamine. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
108	Abnormal circulating thyroid hormone levels in MCT8-deficiency are not caused by increased hepatic conversion through type i-deiodinase. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0

#	ARTICLE	IF	CITATIONS
109	New G protein coupled receptor targets for 3-iodotyronamine. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
110	Effects of repeated 3-TIAM treatment on thyrocytes in mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
111	Establishment of a non-isotopic activity assay for thyroid hormone transporters. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2015, 122, .	1.2	0
112	Analysis of Human TAAR8 and Murine Taar8b Mediated Signaling Pathways and Expression Profile. <i>International Journal of Molecular Sciences</i> , 2014, 15, 20638-20655.	4.1	23
113	Thyroxine: beneficial for mutated TR β receptors thwarting thyroid hormone action?. <i>Lancet Diabetes and Endocrinology</i> , 2014, 2, 602-603.	11.4	1
114	Selenium status in patients with autoimmune and non-autoimmune thyroid diseases from four European countries. <i>Expert Review of Endocrinology and Metabolism</i> , 2014, 9, 685-692.	2.4	12
115	Crystal structure of mammalian selenocysteine-dependent iodothyronine deiodinase suggests a peroxiredoxin-like catalytic mechanism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 10526-10531.	7.1	89
116	High Serum Thyrotropin Levels Are Associated with Retinal Arteriolar Narrowing in the General Population. <i>Thyroid</i> , 2014, 24, 1473-1478.	4.5	21
117	Transport of Thyroid Hormone in Brain. <i>Frontiers in Endocrinology</i> , 2014, 5, 98.	3.5	77
118	Supplementieren oder nicht? Das Spurenelement Selen. <i>Perspectives in Medicine</i> , 2014, 2, 72-78.	0.3	0
119	Selenite supplementation in euthyroid subjects with thyroid peroxidase antibodies. <i>Clinical Endocrinology</i> , 2014, 80, 444-451.	2.4	49
120	Detection of 3,5-Diiodothyronine in Sera of Patients with Altered Thyroid Status Using a New Monoclonal Antibody-Based Chemiluminescence Immunoassay. <i>Thyroid</i> , 2014, 24, 1350-1360.	4.5	64
121	Hepatic metabolite profiles in mice with a suboptimal selenium status. <i>Journal of Nutritional Biochemistry</i> , 2014, 25, 914-922.	4.2	20
122	Soy isoflavones interfere with thyroid hormone homeostasis in orchidectomized middle-aged rats. <i>Toxicology and Applied Pharmacology</i> , 2014, 278, 124-134.	2.8	28
123	Role of 3,5-diiodothyronine in chronic kidney disease. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, .	1.2	3
124	Hormone des Hypothalamus und der Hypophyse. <i>Springer-Lehrbuch</i> , 2014, , 483-494.	0.0	0
125	Steroidhormone – Produkte von Nebennierenrinde und Keimdrüsen. <i>Springer-Lehrbuch</i> , 2014, , 495-511.	0.0	0
126	Schilddrüsenhormone – Zentrale Regulatoren von Entwicklung, Wachstum, Grundumsatz, Stoffwechsel und Zelldifferenzierung. <i>Springer-Lehrbuch</i> , 2014, , 512-527.	0.0	0

#	ARTICLE	IF	CITATIONS
127	Screening assays on inhibitors, modulators and substrates of deiodinase and dehalogenase activities. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, .	1.2	0
128	Functional and metabolic responses of thyrocytes to 3-Iodothyronamine (3-TIAM). <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, .	1.2	0
129	Evidence for thymimetic action of 3,5-T2 in diet-induced obese mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, .	1.2	0
130	The mouse Leydig cell line MLTC-1 prefers L-thyroxine over 3,3',5-triiodo-L-thyronine in transport across plasma membrane and shows steroidogenesis response to thyroid hormone treatment. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, .	1.2	0
131	Aspects of 3-iodothyronamine (3TIAM) induced signaling by human and mouse trace amine-associated receptor 5 (TAAR5). <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, .	1.2	0
132	Function of thyroid hormone transporters in the central nervous system. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2013, 1830, 3965-3973.	2.4	48
133	Mechanism-based testing strategy using in vitro approaches for identification of thyroid hormone disrupting chemicals. <i>Toxicology in Vitro</i> , 2013, 27, 1320-1346.	2.4	165
134	Serum selenium is low in newly diagnosed Graves' disease: a population-based study. <i>Clinical Endocrinology</i> , 2013, 79, 584-590.	2.4	84
135	Autoantibodies to the IGF1 Receptor in Graves' Orbitopathy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 752-760.	3.6	77
136	Evaluation of the Association between Persistent Organic Pollutants (POPs) and Diabetes in Epidemiological Studies: A National Toxicology Program Workshop Review. <i>Environmental Health Perspectives</i> , 2013, 121, 774-783.	6.0	280
137	Selenium and the thyroid. <i>Current Opinion in Endocrinology, Diabetes and Obesity</i> , 2013, 20, 441-448.	2.3	60
138	Serum 25-Hydroxyvitamin D and Cancer Risk in Older Adults: Results from a Large German Prospective Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2013, 22, 905-916.	2.5	61
139	Se- and S-Based Thiouracil and Methimazole Analogues Exert Different Inhibitory Mechanisms on Type 1 and Type 2 Deiodinases. <i>European Thyroid Journal</i> , 2013, 2, 252-258.	2.4	18
140	Strong associations of 25-hydroxyvitamin D concentrations with all-cause, cardiovascular, cancer, and respiratory disease mortality in a large cohort study. <i>American Journal of Clinical Nutrition</i> , 2013, 97, 782-793.	4.7	238
141	Establishment of a competitive chemiluminescence immunoassay to detect 3,5-T2 in human serum. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2013, 121, .	1.2	1
142	Strong associations of 25-hydroxyvitamin D levels with all-cause, cardiovascular, cancer and respiratory disease mortality in a large cohort study. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2013, 121, .	1.2	1
143	Thyroid hormone transporters during testicular development in rodents. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2013, 121, .	1.2	0
144	Orchidectomy of middle-aged rats decreases liver deiodinase 1 and pituitary deiodinase 2 activity. <i>Journal of Endocrinology</i> , 2012, 215, 247-256.	2.6	21

#	ARTICLE	IF	CITATIONS
145	Tyrosine Kinase Inhibitors Noncompetitively Inhibit MCT8-Mediated Iodothyronine Transport. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, E100-E105.	3.6	84
146	Identification of Iopanoic Acid as Substrate of Type 1 Deiodinase by a Novel Nonradioactive Iodide-Release Assay. <i>Endocrinology</i> , 2012, 153, 2506-2513.	2.8	56
147	Thyroid Function Is Maintained Despite Increased Oxidative Stress in Mice Lacking Selenoprotein Biosynthesis in Thyroid Epithelial Cells. <i>Antioxidants and Redox Signaling</i> , 2012, 17, 902-913.	5.4	37
148	LC-MS/MS detection of thyroid hormone metabolites in tissue samples. <i>Expert Review of Endocrinology and Metabolism</i> , 2012, 7, 511-513.	2.4	2
149	Plasma Bile Acids Are Associated with Energy Expenditure and Thyroid Function in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, 535-542.	3.6	57
150	Bone Turnover and Bone Mineral Density Are Independently Related to Selenium Status in Healthy Euthyroid Postmenopausal Women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, 4061-4070.	3.6	91
151	Standardization of Misleading Immunoassay Based 25-Hydroxyvitamin D Levels with Liquid Chromatography Tandem-Mass Spectrometry in a Large Cohort Study. <i>PLoS ONE</i> , 2012, 7, e48774.	2.5	46
152	Inhibition of xanthine oxidase reduces wasting and improves outcome in a rat model of cancer cachexia. <i>International Journal of Cancer</i> , 2012, 131, 2187-2196.	5.1	51
153	Does the aromatic l-amino acid decarboxylase contribute to thyronamine biosynthesis?. <i>Molecular and Cellular Endocrinology</i> , 2012, 349, 195-201.	3.2	37
154	Differential Modulation of Beta-Adrenergic Receptor Signaling by Trace Amine-Associated Receptor 1 Agonists. <i>PLoS ONE</i> , 2011, 6, e27073.	2.5	54
155	Circulating Levels of Peroxiredoxin 4 as a Novel Biomarker of Oxidative Stress in Patients with Sepsis. <i>Shock</i> , 2011, 35, 460-465.	2.1	40
156	Aminoaciduria, but normal thyroid hormone levels and signalling, in mice lacking the amino acid and thyroid hormone transporter Slc7a8. <i>Biochemical Journal</i> , 2011, 439, 249-255.	3.7	55
157	Copeptin and Peroxiredoxin-4 Independently Predict Mortality in Patients With Nonspecific Complaints Presenting to the Emergency Department. <i>Academic Emergency Medicine</i> , 2011, 18, 851-859.	1.8	38
158	Comparison of different selenocompounds with respect to nutritional value vs. toxicity using liver cells in culture. <i>Journal of Nutritional Biochemistry</i> , 2011, 22, 945-955.	4.2	102
159	Selenium Transport in Mammals: Selenoprotein P and Its Receptors. , 2011, , 205-219.		2
160	Insights into molecular properties of the human monocarboxylate transporter 8 by combining functional with structural information. <i>Thyroid Research</i> , 2011, 4, S4.	1.5	27
161	Developmental and cell type-specific expression of thyroid hormone transporters in the mouse brain and in primary brain cells. <i>Glia</i> , 2011, 59, 463-471.	4.9	106
162	Evidence for Extrathyroidal Formation of 3-Iodothyronamine in Humans as Provided by a Novel Monoclonal Antibody-Based Chemiluminescent Serum Immunoassay. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, 1864-1872.	3.6	86

#	ARTICLE	IF	CITATIONS
163	Monocarboxylate transporter 8 deficiency: altered thyroid morphology and persistent high triiodothyronine/thyroxine ratio after thyroidectomy. <i>European Journal of Endocrinology</i> , 2011, 165, 555-561.	3.7	42
164	Thyronamines—Past, Present, and Future. <i>Endocrine Reviews</i> , 2011, 32, 64-80.	20.1	118
165	Selenium status, thyroid volume, and multiple nodule formation in an area with mild iodine deficiency. <i>European Journal of Endocrinology</i> , 2011, 164, 585-590.	3.7	103
166	Environmental factors. , 2011, , 385-395.		0
167	Mutation of megalin leads to urinary loss of selenoprotein P and selenium deficiency in serum, liver, kidneys and brain. <i>Biochemical Journal</i> , 2010, 431, 103-111.	3.7	70
168	Midregional Proenkephalin A and N-terminal Protachykinin A are decreased in the cerebrospinal fluid of patients with dementia disorders and acute neuroinflammation. <i>Journal of Neuroimmunology</i> , 2010, 221, 62-67.	2.3	16
169	Xanthohumol, a prenylated chalcone from hops, modulates hepatic expression of genes involved in thyroid hormone distribution and metabolism. <i>Molecular Nutrition and Food Research</i> , 2010, 54, S225-35.	3.3	24
170	Increased risk of thyroid pathology in patients with thyroid hemiagenesis: results of a large cohort case—control study. <i>European Journal of Endocrinology</i> , 2010, 162, 153-160.	3.7	57
171	Essential Molecular Determinants for Thyroid Hormone Transport and First Structural Implications for Monocarboxylate Transporter 8. <i>Journal of Biological Chemistry</i> , 2010, 285, 28054-28063.	3.4	97
172	Effect of mutations of the human serpin protein corticosteroid-binding globulin on cortisol-binding, thermal and protease sensitivity. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2010, 120, 30-37.	2.5	7
173	Proteomic approach reveals novel targets for retinoic acid-mediated therapy of thyroid carcinoma. <i>Molecular and Cellular Endocrinology</i> , 2010, 325, 110-117.	3.2	23
174	Immunoluminometric assay for quantification of peroxiredoxin 4 in human serum. <i>Clinica Chimica Acta</i> , 2010, 411, 1258-1263.	1.1	21
175	Neuronal selenoprotein expression is required for interneuron development and prevents seizures and neurodegeneration. <i>FASEB Journal</i> , 2010, 24, 844-852.	0.5	193
176	Selenium Supplementation Fails to Correct the Selenoprotein Synthesis Defect in Subjects with SBP2 Gene Mutations. <i>Thyroid</i> , 2009, 19, 277-281.	4.5	69
177	Retinoic acid-mediated down-regulation of ENO1/MBP-1 gene products caused decreased invasiveness of the follicular thyroid carcinoma cell lines. <i>Journal of Molecular Endocrinology</i> , 2009, 42, 249-260.	2.5	32
178	Neuronal 3,5-Triiodothyronine (T ₃) Uptake and Behavioral Phenotype of Mice Deficient in <i>Mct8</i> , the Neuronal T ₃ Transporter Mutated in Allan—Herndon—Dudley Syndrome. <i>Journal of Neuroscience</i> , 2009, 29, 9439-9449.	3.6	172
179	Reduced Serum Selenoprotein P Concentrations in German Prostate Cancer Patients. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009, 18, 2386-2390.	2.5	25
180	Interference of Endocrine Disrupters with Thyroid Hormone Receptor—Dependent Transactivation. <i>Toxicological Sciences</i> , 2009, 110, 125-137.	3.1	84

#	ARTICLE	IF	CITATIONS
181	Surface translocation and tri-iodothyronine uptake of mutant MCT8 proteins are cell type-dependent. <i>Journal of Molecular Endocrinology</i> , 2009, 43, 263-271.	2.5	56
182	Downregulation of the hepatic selenoprotein biosynthesis machinery impairs selenium metabolism during the acute phase response in mice. <i>FASEB Journal</i> , 2009, 23, 1758-1765.	0.5	128
183	Selenium and thyroid. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2009, 23, 815-827.	4.7	169
184	Welcome to Thyroid Research. <i>Thyroid Research</i> , 2008, 1, 1.	1.5	4
185	Development of a validated liquid chromatography/tandem mass spectrometry method for the distinction of thyronine and thyronamine constitutional isomers and for the identification of new deiodinase substrates. <i>Rapid Communications in Mass Spectrometry</i> , 2008, 22, 3286-3296.	1.5	54
186	On the importance of selenium and iodine metabolism for thyroid hormone biosynthesis and human health. <i>Molecular Nutrition and Food Research</i> , 2008, 52, 1235-1246.	3.3	220
187	New assay for the measurement of selenoprotein P as a sepsis biomarker from serum. <i>Journal of Trace Elements in Medicine and Biology</i> , 2008, 22, 24-32.	3.0	114
188	Peroxides and Peroxide-Degrading Enzymes in the Thyroid. <i>Antioxidants and Redox Signaling</i> , 2008, 10, 1577-1592.	5.4	65
189	Detection of stable N-terminal protachykinin A immunoreactivity in human plasma and cerebrospinal fluid. <i>Peptides</i> , 2008, 29, 1201-1206.	2.4	14
190	Environment and endocrinology: The case of thyroidology. <i>Annales D'Endocrinologie</i> , 2008, 69, 116-122.	1.4	52
191	Thyronamines Are Isozyme-Specific Substrates of Deiodinases. <i>Endocrinology</i> , 2008, 149, 3037-3045.	2.8	72
192	V2 vasopressin receptor deficiency causes changes in expression and function of renal and hypothalamic components involved in electrolyte and water homeostasis. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F1177-F1190.	2.7	19
193	Hepatic selenoprotein P (SePP) expression restores selenium transport and prevents infertility and motor-incoordination in <i>Sepp</i> -knockout mice. <i>Biochemical Journal</i> , 2008, 409, 741-749.	3.7	132
194	Development of a validated liquid chromatography/tandem mass spectrometry method for the distinction of thyronine and thyronamine constitutional isomers and for the identification of new deiodinase substrates. , 2008, 22, 3286.		1
195	Selenoproteins – biochemistry and clinical relevance. <i>Biological Chemistry</i> , 2007, 388, 985-6.	2.5	11
196	The Ultraviolet Filter Benzophenone 2 Interferes with the Thyroid Hormone Axis in Rats and Is a Potent in Vitro Inhibitor of Human Recombinant Thyroid Peroxidase. <i>Endocrinology</i> , 2007, 148, 2835-2844.	2.8	97
197	Selenoproteins of the thyroid gland: expression, localization and possible function of glutathione peroxidase 3. <i>Biological Chemistry</i> , 2007, 388, 1053-1059.	2.5	105
198	Thyroid hormone transporters in health and disease: advances in thyroid hormone deiodination. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2007, 21, 173-191.	4.7	63

#	ARTICLE	IF	CITATIONS
199	Endocrine Disruptors and the Thyroid Gland—A Combined in Vitro and in Vivo Analysis of Potential New Biomarkers. <i>Environmental Health Perspectives</i> , 2007, 115, 77-83.	6.0	125
200	Effects of a 5-day treatment with the UV-filter octyl-methoxycinnamate (OMC) on the function of the hypothalamo-pituitary—thyroid function in rats. <i>Toxicology</i> , 2007, 238, 192-199.	4.2	75
201	Procalcitonin is elevated in the cerebrospinal fluid of patients with dementia and acute neuroinflammation. <i>Journal of Neuroimmunology</i> , 2007, 189, 169-174.	2.3	15
202	Neuronal and ependymal expression of selenoprotein P in the human brain. <i>Journal of Neural Transmission</i> , 2007, 114, 877-884.	2.8	84
203	Effects of isoflavonoids and other plant-derived compounds on the hypothalamus—pituitary—thyroid hormone axis. <i>Maturitas</i> , 2006, 55, S14-S25.	2.4	25
204	Proenkephalin A 119—159, a stable proenkephalin A precursor fragment identified in human circulation. <i>Peptides</i> , 2006, 27, 1835-1840.	2.4	50
205	Genistein and other soya isoflavones are potent ligands for transthyretin in serum and cerebrospinal fluid. <i>British Journal of Nutrition</i> , 2006, 95, 1171-1176.	2.3	54
206	Selenium-Dependent Pre- and Posttranscriptional Mechanisms Are Responsible for Sexual Dimorphic Expression of Selenoproteins in Murine Tissues. <i>Endocrinology</i> , 2006, 147, 5883-5892.	2.8	109
207	Synthesis and Metabolism of Thyroid Hormones Is Preferentially Maintained in Selenium-Deficient Transgenic Mice. <i>Endocrinology</i> , 2006, 147, 1306-1313.	2.8	73
208	Selenium and goiter prevalence in borderline iodine sufficiency. <i>European Journal of Endocrinology</i> , 2006, 155, 807-812.	3.7	39
209	Grundlagen der parakrinen, autokrinen und intrakrinen Regulation endokriner Organe. , 2006, , 3-24.		0
210	Hepatically derived selenoprotein P is a key factor for kidney but not for brain selenium supply. <i>Biochemical Journal</i> , 2005, 386, 221-226.	3.7	183
211	Free-flow isoelectric focusing of proteins remaining in cell fragments following sonication of thyroid carcinoma cells. <i>Electrophoresis</i> , 2005, 26, 2109-2116.	2.4	41
212	Xanthohumol stimulates iodide uptake in rat thyroid-derived FRTL-5 cells. <i>Molecular Nutrition and Food Research</i> , 2005, 49, 832-836.	3.3	17
213	Developmentally regulated thyroid hormone distributor proteins in marsupials, a reptile, and fish. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2005, 288, R1264-R1272.	1.8	37
214	Letter re: Id1 Gene Expression in Hyperplastic and Neoplastic Thyroid Tissues. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005, 90, 5906-5906.	3.6	12
215	Selenoprotein Expression in H ₁₄ Cell Carcinomas and in the Human H ₁₄ Cell Carcinoma Line XTC.UC1. <i>Thyroid</i> , 2005, 15, 405-416.	4.5	11
216	Selenium, the Thyroid, and the Endocrine System. <i>Endocrine Reviews</i> , 2005, 26, 944-984.	20.1	444

#	ARTICLE	IF	CITATIONS
217	Hepatic deiodinase activity is dispensable for the maintenance of normal circulating thyroid hormone levels in mice. <i>Biochemical and Biophysical Research Communications</i> , 2005, 337, 739-745.	2.1	41
218	Selenium and the Control of Thyroid Hormone Metabolism. <i>Thyroid</i> , 2005, 15, 841-853.	4.5	128
219	Thyroid Hormone Metabolism. , 2004, , 480-489.		0
220	Guard Your Master: Thyroid Hormone Receptors Protect Their Gland of Origin from Thyroid Cancer. <i>Endocrinology</i> , 2004, 145, 4427-4429.	2.8	4
221	Contribution of Interleukin-12 to the Pathogenesis of Non-thyroidal Illness. <i>Hormone and Metabolic Research</i> , 2004, 36, 101-106.	1.5	32
222	Human thyroid carcinoma cell lines show different retinoic acid receptor repertoires and retinoid responses. <i>European Journal of Endocrinology</i> , 2004, 150, 547-556.	3.7	38
223	Expression Profiling and Genetic Alterations of the Selenoproteins GI-GPx and SePP in Colorectal Carcinogenesis. <i>Nutrition and Cancer</i> , 2004, 48, 6-14.	2.0	111
224	Interleukin-18, a proinflammatory cytokine, contributes to the pathogenesis of non-thyroidal illness mainly via the central part of the hypothalamus-pituitary-thyroid axis. <i>European Journal of Endocrinology</i> , 2004, 151, 497-502.	3.7	34
225	55th Mosbach Colloquium: How nutrients influence gene activity. <i>Biological Chemistry</i> , 2004, 385, 569-569.	2.5	0
226	CD82, and CD63 in thyroid cancer.. <i>International Journal of Molecular Medicine</i> , 2004, 14, 517.	4.0	3
227	Endocrine active compounds affect thyrotropin and thyroid hormone levels in serum as well as endpoints of thyroid hormone action in liver, heart and kidney. <i>Toxicology</i> , 2004, 205, 95-102.	4.2	140
228	Selenium and brain function: a poorly recognized liaison. <i>Brain Research Reviews</i> , 2004, 45, 164-178.	9.0	281
229	Association between mutations in a thyroid hormone transporter and severe X-linked psychomotor retardation. <i>Lancet, The</i> , 2004, 364, 1435-1437.	13.7	615
230	Efficient selenium transfer from mother to offspring in selenoprotein-P-deficient mice enables dose-dependent rescue of phenotypes associated with selenium deficiency. <i>Biochemical Journal</i> , 2004, 378, 21-26.	3.7	95
231	Glutathione peroxidase isoforms as part of the local antioxidative defense system in normal and Barrett's esophagus. <i>International Journal of Cancer</i> , 2003, 105, 300-304.	5.1	50
232	Gene disruption discloses role of selenoprotein P in selenium delivery to target tissues. <i>Biochemical Journal</i> , 2003, 370, 397-402.	3.7	378
233	Selenium Supplementation Enhances Low Selenium Levels and Stimulates Glutathione Peroxidase Activity in Peripheral Blood and Distal Colon Mucosa in Past and Present Carriers of Colon Adenomas. <i>Nutrition and Cancer</i> , 2003, 46, 125-130.	2.0	30
234	Expression of telomerase genes in thyroid carcinoma. <i>International Journal of Oncology</i> , 2002, 21, 265.	3.3	11

#	ARTICLE	IF	CITATIONS
235	Thyroid hormone receptors and type I iodothyronine 5 α -deiodinase activity of human thyroid toxic adenomas and benign cold nodules. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2002, 110, 166-170.	1.2	21
236	Proinflammatory cytokines inhibit the expression and function of human type I 5'-deiodinase in HepG2 hepatocarcinoma cells. <i>European Journal of Endocrinology</i> , 2002, 146, 559-566.	3.7	62
237	Iodothyronine Deiodinases. <i>Methods in Enzymology</i> , 2002, 347, 125-167.	1.0	157
238	The Impact of Iron and Selenium Deficiencies on Iodine and Thyroid Metabolism: Biochemistry and Relevance to Public Health. <i>Thyroid</i> , 2002, 12, 867-878.	4.5	274
239	Spectral karyotype analysis of colon cancer cell lines of the tumor suppressor and mutator pathway. <i>Cytogenetic and Genome Research</i> , 2002, 98, 22-28.	1.1	28
240	The promoter of the human sodium/iodide-symporter gene responds to retinoic acid. <i>Molecular and Cellular Endocrinology</i> , 2002, 189, 145-155.	3.2	52
241	Clinical impact of retinoids in redifferentiation therapy of advanced thyroid cancer: final results of a pilot study. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2002, 29, 775-782.	6.4	167
242	A complex DNA-repeat structure within the Selenoprotein P promoter contains a functionally relevant polymorphism and is genetically unstable under conditions of mismatch repair deficiency. <i>European Journal of Human Genetics</i> , 2002, 10, 499-504.	2.8	42
243	Identification of an element within the promoter of human selenoprotein P responsive to transforming growth factor- β ² . <i>FEBS Journal</i> , 2001, 268, 6176-6181.	0.2	27
244	Modulation of selenoprotein P expression by TGF β ¹ is mediated by Smad proteins. <i>BioFactors</i> , 2001, 14, 135-142.	5.4	21
245	Innovative strategies for the treatment of thyroid cancer. <i>European Journal of Endocrinology</i> , 2000, 143, 15-24.	3.7	64
246	Retinoic Acid Redifferentiation Therapy for Thyroid Cancer. <i>Thyroid</i> , 2000, 10, 393-406.	4.5	137
247	Selenium in Biology: Facts and Medical Perspectives. <i>Biological Chemistry</i> , 2000, 381, 849-64.	2.5	241
248	Effects of proinflammatory cytokines on anterior pituitary 5'-deiodinase type I and type II. <i>Journal of Endocrinology</i> , 2000, 167, 505-515.	2.6	61
249	Inverse mRNA Expression of the Selenocysteine-Containing Proteins GI-GPx and SeP in Colorectal Adenomas Compared With Adjacent Normal Mucosa. <i>Nutrition and Cancer</i> , 2000, 37, 108-116.	2.0	64
250	Regulation of CD97 Protein in Thyroid Carcinoma ¹ . <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999, 84, 1104-1109.	3.6	56
251	Type 1 deiodinase is stimulated by iodothyronines and involved in thyroid hormone metabolism in human somatomammotroph GX cells. <i>European Journal of Endocrinology</i> , 1999, 140, 367-370.	3.7	11
252	The selenoprotein thioredoxin reductase is expressed in peripheral blood monocytes and THP1 human myeloid leukemia cells \bullet regulation by 1,25 α -dihydroxyvitamin D ₃ and selenite. <i>BioFactors</i> , 1999, 10, 329-338.	5.4	36

#	ARTICLE	IF	CITATIONS
253	The trace element selenium and the thyroid gland. <i>Biochimie</i> , 1999, 81, 527-533.	2.6	139
254	Local activation and inactivation of thyroid hormones: the deiodinase family. <i>Molecular and Cellular Endocrinology</i> , 1999, 151, 103-119.	3.2	316
255	Transforming growth factor- β 1 inhibits expression of selenoprotein P in cultured human liver cells. <i>FEBS Letters</i> , 1999, 460, 23-26.	2.8	29
256	Increased I-131 Uptake in Local Recurrence and Distant Metastases After Second Treatment With Retinoic Acid. <i>Clinical Nuclear Medicine</i> , 1999, 24, 849.	1.3	12
257	Functional retinoid and thyroid hormone receptors in human thyroid-carcinoma cell lines and tissues. , 1998, 76, 368-376.		42
258	Redifferentiation Therapy with Retinoids: Therapeutic Option for Advanced Follicular and Papillary Thyroid Carcinoma. <i>World Journal of Surgery</i> , 1998, 22, 569-574.	1.6	133
259	Selenoproteins Are Expressed in Fetal Human Osteoblast-like Cells. <i>Biochemical and Biophysical Research Communications</i> , 1998, 245, 101-107.	2.1	72
260	Reverse Transcriptase-Polymerase Chain Reaction Analysis of Thyrocyte-Relevant Genes in Fine-Needle Aspiration Biopsies of the Human Thyroid. <i>Thyroid</i> , 1998, 8, 981-987.	4.5	27
261	Expression pattern of gastrointestinal selenoproteinsâ€” targets for selenium supplementation. <i>Nutrition and Cancer</i> , 1998, 32, 64-70.	2.0	42
262	Functional retinoid and thyroid hormone receptors in human thyroidâ€”carcinoma cell lines and tissues. <i>International Journal of Cancer</i> , 1998, 76, 368-376.	5.1	1
263	Cloning and Characterization of the Human Selenoprotein P Promoter. <i>Journal of Biological Chemistry</i> , 1997, 272, 29364-29371.	3.4	94
264	Consensus statement on lung cancer. <i>European Journal of Cancer Prevention</i> , 1997, 6, 316-322.	1.3	7
265	Retinoic Acid Increases Sodium/Iodide Symporter mRNA Levels in Human Thyroid Cancer Cell Lines and Suppresses Expression of Functional Symporter in Nontransformed FRTL-5 Rat Thyroid Cells. <i>Biochemical and Biophysical Research Communications</i> , 1997, 240, 832-838.	2.1	196
266	Structure of the Human Type I Iodothyronine 5â€”Deiodinase Gene and Localization to Chromosome 1p32â€”p33. <i>Genomics</i> , 1997, 42, 361-363.	2.9	38
267	Local estradiol metabolism in osteoblast- and osteoclast-like cells. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1997, 61, 167-174.	2.5	59
268	The Promoter of the Human Type I 5'-Deiodinase Gene - Mapping of the Transcription Start Site and Identification of a DR+4 Thyroid-Hormone-Responsive Element. <i>FEBS Journal</i> , 1997, 247, 288-297.	0.2	77
269	Different Tissue Distribution, Elimination, and Kinetics of Thyroxine and Its Conformational Analog, the Synthetic Flavonoid EMD 49209 in the Rat. <i>Endocrinology</i> , 1997, 138, 79-84.	2.8	8
270	3,5-Diiodo-L-Thyronine Stimulates Type 1 5'-Deiodinase Activity in Rat Anterior Pituitaries in Vivo and in Reaggregate Cultures and GH3 Cells in Vitro. <i>Endocrinology</i> , 1997, 138, 3242-3248.	2.8	11

#	ARTICLE	IF	CITATIONS
271	Local Estradiol Metabolism in Osteoblast- and Osteoclast-like Cells. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1997, 61, 167-174.	2.5	1
272	Selected Novel Flavones Inhibit the DNA Binding or the DNA Religation Step of Eukaryotic Topoisomerase I. <i>Journal of Biological Chemistry</i> , 1996, 271, 2262-2270.	3.4	200
273	Rapid stimulation of type I 5 α -deiodinase in rat pituitaries by 3,3 α ,5-triiodo-L-thyronine. <i>Molecular and Cellular Endocrinology</i> , 1995, 108, 17-21.	3.2	49
274	Expression and regulation of aromatase cytochrome P450 in THP 1 human myeloid leukaemia cells. <i>Molecular and Cellular Endocrinology</i> , 1995, 110, 27-33.	3.2	32
275	Binding of Thyroxine to Pig Transthyretin, its cDNA Structure, and Other Properties. <i>FEBS Journal</i> , 1995, 230, 977-986.	0.2	1
276	Binding of Thyroxine to Pig Transthyretin, its cDNA Structure, and Other Properties. <i>FEBS Journal</i> , 1995, 230, 977-986.	0.2	25
277	Topoisomerase I Inhibition Enhances Vitamin D-Responsive Expression of the Receptor for Lipopolysaccharide Binding Protein CD 14. <i>Biochemical and Biophysical Research Communications</i> , 1994, 199, 531-539.	2.1	15
278	Early adaptation of thyrotropin and thyroglobulin secretion to experimentally decreased iodine supply in man. <i>Metabolism: Clinical and Experimental</i> , 1992, 41, 1093-1096.	3.4	77
279	Human thyrotropin receptor gene: Expression in thyroid tumors and correlation to markers of thyroid differentiation and dedifferentiation. <i>Molecular and Cellular Endocrinology</i> , 1991, 82, R7-R12.	3.2	131
280	Effect of Biological Alterations of Type I 5' Deiodinase Activity on Affinity Labeled Membrane Proteins in Rat Liver and Kidney*. <i>Endocrinology</i> , 1990, 126, 826-831.	2.8	27
281	5 α -Deiodination in Rat Hepatocytes: Effects of Specific Flavonoid Inhibitors*. <i>Endocrinology</i> , 1990, 126, 1660-1667.	2.8	22
282	Flavonoid Administration Immediately Displaces Thyroxine (T ₄) from Serum Transthyretin, Increases Serum Free T ₄ , and Decreases Serum Thyrotropin in the Rat*. <i>Endocrinology</i> , 1990, 126, 2890-2895.	2.8	52
283	Identification of type I iodothyronine 5 α -deiodinase as a selenoenzyme. <i>Biochemical and Biophysical Research Communications</i> , 1990, 173, 1143-1149.	2.1	373
284	Rapid Effects of the Flavonoid EMD 21388 on Serum Thyroid Hormone Binding and Thyrotropin Regulation in the Rat*. <i>Endocrinology</i> , 1989, 125, 532-537.	2.8	59
285	Thyroid hormone effect on rat heart mitochondrial proteins and affinity labeling with N-bromoacetyl-3,3 α ,5-triiodo-L-thyronine Lack of direct effect on the adenine nucleotide translocase. <i>FEBS Letters</i> , 1989, 255, 385-390.	2.8	24
286	Circadian and Pulsatile Thyrotropin Secretion in Euthyroid Man Under the Influence of Thyroid Hormone and Glucocorticoid Administration*. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1987, 65, 83-88.	3.6	166
287	Crystal structure of phlorizin and the iodothyronine deiodinase inhibitory activity of phloretin analogues. <i>Biochemical Pharmacology</i> , 1986, 35, 2221-2227.	4.4	21
288	Subcellular localization of thyroxine-5-deiodinase in rat liver. <i>Journal of Endocrinological Investigation</i> , 1980, 3, 73-76.	3.3	13

#	ARTICLE	IF	CITATIONS
289	Conversion of T3 and rT3 to 3,3 ² -T2: pH dependency. Clinica Chimica Acta, 1978, 90, 45-51.	1.1	9
290	Darstellung von [¹⁴ C]- ³ P¹, <i>P</i>⁵- ³ P⁵-Di(adenosin ⁵ - ² - ⁵ -pentaphosphat durch direkte Umsetzung von [¹⁴ C]-Adenosin ⁵ - ² - ⁵ -diphosphat mit aktiviertem Adenosin ⁵ - ² - ⁵ -triphosphat. Justus Liebigs Annalen Der Chemie, 1977, 1977, 1160-1166.	0.5	2
291	³ P¹, ³ P⁵-Bis-(5 ² -adenosyl)pentaphosphate: Is this Adenylate Kinase Inhibitor Substrate for Mitochondrial Processes?. Zeitschrift Fur Naturforschung - Section C Journal of Biosciences, 1977, 32, 786-791.	1.4	2
292	3,5-Diiodo-L-Thyronine (3,5-T2) Exerts Thyromimetic Effects on Hypothalamus-Pituitary-Thyroid Axis, Body Composition, and Energy Metabolism in Male Diet-Induced Obese Mice. , 0, .		1
293	Selenoprotein biosynthesis – importance of structural elements in the 3' untranslated region. , 0, 2005, .		0
294	Purkinje cell degeneration and loss of cerebellar interneurons during development of mice lacking neuronal selenoprotein biosynthesis. , 0, 2005, .		0
295	Hepatic deiodinase activity is dispensable for the maintenance of normal circulating thyroid hormone levels, but not for thyroid hormone-dependent gene expression. , 0, 2005, .		0
296	Distinct roles for neuronal selenoproteins in cerebellar development. , 0, 2006, .		0
297	3,5-T2 might play a crucial role in the rodent heart. Endocrine Abstracts, 0, , .	0.0	0
298	3-T1AM signaling in the rat thyrocyte cell line PCCL3. Endocrine Abstracts, 0, , .	0.0	0
299	Circulating 3-T1AM and 3,5-T2 in critically ill patients: a cross-sectional observational study. Endocrine Abstracts, 0, , .	0.0	0