Timo A Lakka

List of Publications by Citations

Source: https://exaly.com/author-pdf/6797455/timo-a-lakka-publications-by-citations.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

80 180 283 33,246 h-index g-index citations papers 6.11 8.9 39,084 305 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
283	The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. <i>JAMA - Journal of the American Medical Association</i> , 2002 , 288, 2709-16	27.4	3341
282	Genetic studies of body mass index yield new insights for obesity biology. <i>Nature</i> , 2015 , 518, 197-206	50.4	2687
281	Discovery and refinement of loci associated with lipid levels. <i>Nature Genetics</i> , 2013 , 45, 1274-1283	36.3	1904
280	Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. <i>Nature Genetics</i> , 2012 , 44, 981-90	36.3	1482
279	Defining the role of common variation in the genomic and biological architecture of adult human height. <i>Nature Genetics</i> , 2014 , 46, 1173-86	36.3	1339
278	New genetic loci link adipose and insulin biology to body fat distribution. <i>Nature</i> , 2015 , 518, 187-196	50.4	920
277	Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. <i>Nature Genetics</i> , 2014 , 46, 234-44	36.3	784
276	Metabolic syndrome and development of diabetes mellitus: application and validation of recently suggested definitions of the metabolic syndrome in a prospective cohort study. <i>American Journal of Epidemiology</i> , 2002 , 156, 1070-7	3.8	624
275	Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. <i>Nature Genetics</i> , 2012 , 44, 991-1005	36.3	621
274	Relation of leisure-time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction. <i>New England Journal of Medicine</i> , 1994 , 330, 1549-54	59.2	619
273	A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. <i>Nature Genetics</i> , 2012 , 44, 659-69	36.3	615
272	Common variants associated with plasma triglycerides and risk for coronary artery disease. <i>Nature Genetics</i> , 2013 , 45, 1345-52	36.3	597
271	Risk of acute coronary events and serum concentration of asymmetrical dimethylarginine. <i>Lancet, The,</i> 2001 , 358, 2127-8	40	490
270	Low levels of leisure-time physical activity and cardiorespiratory fitness predict development of the metabolic syndrome. <i>Diabetes Care</i> , 2002 , 25, 1612-8	14.6	476
269	Uric acid level as a risk factor for cardiovascular and all-cause mortality in middle-aged men: a prospective cohort study. <i>Archives of Internal Medicine</i> , 2004 , 164, 1546-51		474
268	Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. <i>Nature Genetics</i> , 2013 , 45, 501-12	36.3	437
267	Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study. <i>Diabetes</i> , 2005 , 54, 158-65	0.9	434

266	Rare and low-frequency coding variants alter human adult height. <i>Nature</i> , 2017 , 542, 186-190	50.4	412
265	Physical activity attenuates the influence of FTO variants on obesity risk: a meta-analysis of 218,166 adults and 19,268 children. <i>PLoS Medicine</i> , 2011 , 8, e1001116	11.6	379
264	Rare variant in scavenger receptor BI raises HDL cholesterol and increases risk of coronary heart disease. <i>Science</i> , 2016 , 351, 1166-71	33.3	325
263	Exome-wide association study of plasma lipids in >300,000 individuals. <i>Nature Genetics</i> , 2017 , 49, 1758-	1366	310
262	Genetic fine mapping and genomic annotation defines causal mechanisms at type 2 diabetes susceptibility loci. <i>Nature Genetics</i> , 2015 , 47, 1415-25	36.3	292
261	Physical activity in prevention and treatment of the metabolic syndrome. <i>Applied Physiology, Nutrition and Metabolism</i> , 2007 , 32, 76-88	3	287
260	Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. <i>PLoS Genetics</i> , 2013 , 9, e1003500	6	277
259	Sedentary lifestyle, poor cardiorespiratory fitness, and the metabolic syndrome. <i>Medicine and Science in Sports and Exercise</i> , 2003 , 35, 1279-86	1.2	277
258	Genome-wide associations for birth weight and correlations with adult disease. <i>Nature</i> , 2016 , 538, 248-	2 5 2.4	266
257	Effects of exercise training on glucose homeostasis: the HERITAGE Family Study. <i>Diabetes Care</i> , 2005 , 28, 108-14	14.6	265
256	Antioxidant Supplementation in Atherosclerosis Prevention (ASAP) study: a randomized trial of the effect of vitamins E and C on 3-year progression of carotid atherosclerosis. <i>Journal of Internal Medicine</i> , 2000 , 248, 377-86	10.8	258
255	Relationship of physical activity and body mass index to the risk of hypertension: a prospective study in Finland. <i>Hypertension</i> , 2004 , 43, 25-30	8.5	255
254	The genetics of blood pressure regulation and its target organs from association studies in 342,415 individuals. <i>Nature Genetics</i> , 2016 , 48, 1171-1184	36.3	251
253	Fish oil-derived fatty acids, docosahexaenoic acid and docosapentaenoic acid, and the risk of acute coronary events: the Kuopio ischaemic heart disease risk factor study. <i>Circulation</i> , 2000 , 102, 2677-9	16.7	251
252	Abdominal obesity is associated with increased risk of acute coronary events in men. <i>European Heart Journal</i> , 2002 , 23, 706-13	9.5	242
251	BDNF is a novel marker of cognitive function in ageing women: the DR@EXTRA Study. <i>Neurobiology of Learning and Memory</i> , 2008 , 90, 596-603	3.1	237
250	New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. <i>Nature Genetics</i> , 2013 , 45, 76-82	36.3	232
249	The Influence of Age and Sex on Genetic Associations with Adult Body Size and Shape: A Large-Scale Genome-Wide Interaction Study. <i>PLoS Genetics</i> , 2015 , 11, e1005378	6	220

248	Mercury accumulation and accelerated progression of carotid atherosclerosis: a population-based prospective 4-year follow-up study in men in eastern Finland. <i>Atherosclerosis</i> , 2000 , 148, 265-73	3.1	217
247	Genome-wide association analysis identifies three new susceptibility loci for childhood body mass index. <i>Human Molecular Genetics</i> , 2016 , 25, 389-403	5.6	202
246	Risk of acute coronary events according to serum concentrations of enterolactone: a prospective population-based case-control study. <i>Lancet, The</i> , 1999 , 354, 2112-5	40	202
245	Increased risk of acute myocardial infarction in carriers of the hemochromatosis gene Cys282Tyr mutation: a prospective cohort study in men in eastern Finland. <i>Circulation</i> , 1999 , 100, 1274-9	16.7	196
244	Physical activity, cardiovascular risk factors, and mortality among Finnish adults with diabetes. <i>Diabetes Care</i> , 2005 , 28, 799-805	14.6	195
243	Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. <i>Nature Genetics</i> , 2018 , 50, 26-41	36.3	186
242	Cardiovascular fitness as a predictor of mortality in men. Archives of Internal Medicine, 2001, 161, 825-3	1	183
241	Occupational, commuting, and leisure-time physical activity in relation to total and cardiovascular mortality among Finnish subjects with type 2 diabetes. <i>Circulation</i> , 2004 , 110, 666-73	16.7	182
240	Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic risk factors. <i>Nature Genetics</i> , 2019 , 51, 804-814	36.3	181
239	Trans-ancestry meta-analyses identify rare and common variants associated with blood pressure and hypertension. <i>Nature Genetics</i> , 2016 , 48, 1151-1161	36.3	181
238	Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. <i>Circulation</i> , 2001 , 103, 2674-80	16.7	173
237	Low physical activity as a predictor for total and cardiovascular disease mortality in middle-aged men and women in Finland. <i>European Heart Journal</i> , 2004 , 25, 2204-11	9.5	170
236	Abdominal obesity is associated with accelerated progression of carotid atherosclerosis in men. <i>Atherosclerosis</i> , 2001 , 154, 497-504	3.1	156
235	Effect of exercise training on plasma levels of C-reactive protein in healthy adults: the HERITAGE Family Study. <i>European Heart Journal</i> , 2005 , 26, 2018-25	9.5	145
234	Serum fatty acid composition predicts development of impaired fasting glycaemia and diabetes in middle-aged men. <i>Diabetic Medicine</i> , 2002 , 19, 456-64	3.5	141
233	Stress-induced blood pressure reactivity and incident stroke in middle-aged men. <i>Stroke</i> , 2001 , 32, 126	3 <i>-</i> ₹.₱	137
232	Low serum lycopene concentration is associated with an excess incidence of acute coronary events and stroke: the Kuopio Ischaemic Heart Disease Risk Factor Study. <i>British Journal of Nutrition</i> , 2001 , 85, 749-54	3.6	127
231	A central role for GRB10 in regulation of islet function in man. <i>PLoS Genetics</i> , 2014 , 10, e1004235	6	124

(2001-2007)

230	Occupational, commuting and leisure-time physical activity in relation to coronary heart disease among middle-aged Finnish men and women. <i>Atherosclerosis</i> , 2007 , 194, 490-7	3.1	123
229	Life-Course Genome-wide Association Study Meta-analysis of Total Body BMD and Assessment of Age-Specific Effects. <i>American Journal of Human Genetics</i> , 2018 , 102, 88-102	11	119
228	Fine mapping of five loci associated with low-density lipoprotein cholesterol detects variants that double the explained heritability. <i>PLoS Genetics</i> , 2011 , 7, e1002198	6	118
227	Intra-person variability of various physical activity assessments in the Kuopio Ischaemic Heart Disease Risk Factor Study. <i>International Journal of Epidemiology</i> , 1992 , 21, 467-72	7.8	116
226	Depression and early retirement: prospective population based study in middle aged men. <i>Journal of Epidemiology and Community Health</i> , 2005 , 59, 70-4	5.1	115
225	Cardiorespiratory fitness and the progression of carotid atherosclerosis in middle-aged men. <i>Annals of Internal Medicine</i> , 2001 , 134, 12-20	8	114
224	Genome-wide meta-analysis of 241,258 adults accounting for smoking behaviour identifies novel loci for obesity traits. <i>Nature Communications</i> , 2017 , 8, 14977	17.4	105
223	Blood pressure and the progression of carotid atherosclerosis in middle-aged men. <i>Hypertension</i> , 1999 , 34, 51-6	8.5	104
222	Genome-wide physical activity interactions in adiposity - A meta-analysis of 200,452 adults. <i>PLoS Genetics</i> , 2017 , 13, e1006528	6	103
221	Metabolic syndrome and cognitive function: a population-based follow-up study in elderly women. <i>Dementia and Geriatric Cognitive Disorders</i> , 2007 , 23, 29-34	2.6	101
220	Aerobic exercise and the lipid profile in type 1 diabetic men: a randomized controlled trial. <i>Medicine and Science in Sports and Exercise</i> , 2000 , 32, 1541-8	1.2	100
219	Delaying decline in pulmonary function with physical activity: a 25-year follow-up. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2003 , 168, 494-9	10.2	99
218	Effects of aerobic physical exercise on inflammation and atherosclerosis in men: the DNASCO Study: a six-year randomized, controlled trial. <i>Annals of Internal Medicine</i> , 2004 , 140, 1007-14	8	99
217	An insertion/deletion polymorphism in the alpha2B-adrenergic receptor gene is a novel genetic risk factor for acute coronary events. <i>Journal of the American College of Cardiology</i> , 2001 , 37, 1516-22	15.1	98
216	Weight gain and the risk of developing insulin resistance syndrome. <i>Diabetes Care</i> , 1998 , 21, 1637-43	14.6	95
215	Trans-ethnic fine-mapping of lipid loci identifies population-specific signals and allelic heterogeneity that increases the trait variance explained. <i>PLoS Genetics</i> , 2013 , 9, e1003379	6	94
214	Cardiorespiratory fitness and the risk for stroke in men. Archives of Internal Medicine, 2003, 163, 1682-8		94
213	Leucine7 to proline7 polymorphism in the preproneuropeptide Y is associated with the progression of carotid atherosclerosis, blood pressure and serum lipids in Finnish men. <i>Atherosclerosis</i> , 2001 , 159, 145-51	3.1	93

212	Exercise-induced silent myocardial ischemia and coronary morbidity and mortality in middle-aged men. <i>Journal of the American College of Cardiology</i> , 2001 , 38, 72-9	15.1	91
211	Determinants of cardiorespiratory fitness in men aged 42 to 60 years with and without cardiovascular disease. <i>American Journal of Cardiology</i> , 2009 , 103, 1598-604	3	90
210	Physical activity and sedentary behaviour in relation to cardiometabolic risk in children: cross-sectional findings from the Physical Activity and Nutrition in Children (PANIC) Study. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2014 , 11, 55	8.4	89
209	Systemic immune mediators and lifestyle changes in the prevention of type 2 diabetes: results from the Finnish Diabetes Prevention Study. <i>Diabetes</i> , 2006 , 55, 2340-6	0.9	87
208	Dyslipidaemia as a predictor of hypertension in middle-aged men. <i>European Heart Journal</i> , 2008 , 29, 25	6 9. 8	86
207	Hyperinsulinemia and the risk of cardiovascular death and acute coronary and cerebrovascular events in men: the Kuopio Ischaemic Heart Disease Risk Factor Study. <i>Archives of Internal Medicine</i> , 2000 , 160, 1160-8		86
206	Epidemiology and treatment of the metabolic syndrome. <i>Annals of Medicine</i> , 2004 , 36, 332-46	1.5	82
205	The genetic regulatory signature of type 2 diabetes in human skeletal muscle. <i>Nature Communications</i> , 2016 , 7, 11764	17.4	82
204	Systolic blood pressure during recovery from exercise and the risk of acute myocardial infarction in middle-aged men. <i>Hypertension</i> , 2004 , 44, 820-5	8.5	81
203	Plasma vitamin C modifies the association between hypertension and risk of stroke. <i>Stroke</i> , 2002 , 33, 1568-73	6.7	79
202	A novel common variant in DCST2 is associated with length in early life and height in adulthood. <i>Human Molecular Genetics</i> , 2015 , 24, 1155-68	5.6	77
201	Perceived health as a predictor of early retirement. <i>Scandinavian Journal of Work, Environment and Health</i> , 2004 , 30, 287-92	4.3	77
200	Validation of metabolic syndrome score by confirmatory factor analysis in children and adults and prediction of cardiometabolic outcomes in adults. <i>Diabetologia</i> , 2014 , 57, 940-9	10.3	73
199	Physical activity and diabetes complications in patients with type 1 diabetes: the Finnish Diabetic Nephropathy (FinnDiane) Study. <i>Diabetes Care</i> , 2008 , 31, 230-2	14.6	71
198	Anger expression and incident stroke: prospective evidence from the Kuopio ischemic heart disease study. <i>Stroke</i> , 1999 , 30, 523-8	6.7	71
197	Moderate to high intensity conditioning leisure time physical activity and high cardiorespiratory fitness are associated with reduced plasma fibrinogen in eastern Finnish men. <i>Journal of Clinical Epidemiology</i> , 1993 , 46, 1119-27	5.7	71
196	Heart rate response during exercise test and cardiovascular mortality in middle-aged men. <i>European Heart Journal</i> , 2006 , 27, 582-8	9.5	70
195	Cardiorespiratory fitness and vigorous leisure-time physical activity modify the association of small size at birth with the metabolic syndrome. <i>Diabetes Care</i> , 2003 , 26, 2156-64	14.6	70

(2006-2007)

194	Carotid intima-media thickness and cognitive function in elderly women: a population-based study. <i>Neuroepidemiology</i> , 2007 , 28, 207-13	5.4	69	
193	Dietary factors associated with overweight and body adiposity in Finnish children aged 6-8 years: the PANIC Study. <i>International Journal of Obesity</i> , 2012 , 36, 950-5	5.5	68	
192	Dietary Intake, FTO Genetic Variants, and Adiposity: A Combined Analysis of Over 16,000 Children and Adolescents. <i>Diabetes</i> , 2015 , 64, 2467-76	0.9	66	
191	Associations of motor and cardiovascular performance with academic skills in children. <i>Medicine and Science in Sports and Exercise</i> , 2014 , 46, 1016-24	1.2	66	
190	Leisure-time physical activity and the metabolic syndrome in the Finnish diabetes prevention study. <i>Diabetes Care</i> , 2010 , 33, 1610-7	14.6	66	
189	Serum high sensitivity C-reactive protein and cognitive function in elderly women. <i>Age and Ageing</i> , 2007 , 36, 443-8	3	65	
188	Leptin and leptin receptor gene polymorphisms and changes in glucose homeostasis in response to regular exercise in nondiabetic individuals: the HERITAGE family study. <i>Diabetes</i> , 2004 , 53, 1603-8	0.9	65	
187	Leisure time physical activity is associated with poor glycemic control in type 1 diabetic women: the FinnDiane study. <i>Diabetes Care</i> , 2005 , 28, 777-82	14.6	62	
186	Physical workload and risk of early retirement: prospective population-based study among middle-aged men. <i>Journal of Occupational and Environmental Medicine</i> , 2002 , 44, 930-9	2	61	
185	Low serum folate concentrations are associated with an excess incidence of acute coronary events: the Kuopio Ischaemic Heart Disease Risk Factor Study. <i>European Journal of Clinical Nutrition</i> , 2000 , 54, 424-8	5.2	61	
184	Multi-ancestry genome-wide gene-smoking interaction study of 387,272 individuals identifies new loci associated with serum lipids. <i>Nature Genetics</i> , 2019 , 51, 636-648	36.3	59	
183	Assessment of body composition by dual-energy X-ray absorptiometry, bioimpedance analysis and anthropometrics in children: the Physical Activity and Nutrition in Children study. <i>Clinical Physiology and Functional Imaging</i> , 2015 , 35, 21-33	2.4	58	
182	Polymorphism in high density lipoprotein paraoxonase gene and risk of acute myocardial infarction in men: prospective nested case-control study. <i>BMJ: British Medical Journal</i> , 1999 , 319, 487-9; discussion 490		56	
181	Integrative analysis of gene expression, DNA methylation, physiological traits, and genetic variation in human skeletal muscle. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 10883-10888	11.5	54	
180	Cardiorespiratory fitness as a feature of metabolic syndrome in older men and women: the Dose-Responses to Exercise Training study (DRQ EXTRA). <i>Diabetes Care</i> , 2008 , 31, 1242-7	14.6	54	
179	Systolic blood pressure response to exercise testing is related to the risk of acute myocardial infarction in middle-aged men. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 421-428		51	
178	Food consumption, nutrient intake and the risk of having metabolic syndrome: the DRQ EXTRA Study. European Journal of Clinical Nutrition, 2011, 65, 368-77	5.2	49	
177	High dietary methionine intake increases the risk of acute coronary events in middle-aged men. Nutrition, Metabolism and Cardiovascular Diseases, 2006, 16, 113-20	4.5	48	

176	Cross-Sectional Associations of Objectively-Measured Physical Activity and Sedentary Time with Body Composition and Cardiorespiratory Fitness in Mid-Childhood: The PANIC Study. <i>Sports Medicine</i> , 2017 , 47, 769-780	10.6	47
175	Epidemiological studies of exercise in diabetes prevention. <i>Applied Physiology, Nutrition and Metabolism</i> , 2007 , 32, 583-95	3	47
174	A principal component meta-analysis on multiple anthropometric traits identifies novel loci for body shape. <i>Nature Communications</i> , 2016 , 7, 13357	17.4	46
173	The trans-ancestral genomic architecture of glycemic traits. <i>Nature Genetics</i> , 2021 , 53, 840-860	36.3	44
172	Protein-coding variants implicate novel genes related to lipid homeostasis contributing to body-fat distribution. <i>Nature Genetics</i> , 2019 , 51, 452-469	36.3	44
171	Cardiopulmonary fitness is a function of lean mass, not total body weight: The DR@ EXTRA study. <i>European Journal of Preventive Cardiology</i> , 2015 , 22, 1171-9	3.9	43
170	Dietary factors and their associations with socioeconomic background in Finnish girls and boys 6-8 years of age: the PANIC Study. <i>European Journal of Clinical Nutrition</i> , 2011 , 65, 1211-8	5.2	43
169	Associations of physical activity and sedentary behavior with academic skillsa follow-up study among primary school children. <i>PLoS ONE</i> , 2014 , 9, e107031	3.7	41
168	Metabolic syndrome and the progression of carotid intima-media thickness in elderly women. <i>Archives of Internal Medicine</i> , 2006 , 166, 444-9		41
167	Asymmetrical dimethylarginine (ADMA) and risk of acute coronary events. Does statin treatment influence plasma ADMA levels?. <i>Atherosclerosis Supplements</i> , 2003 , 4, 19-22	1.7	41
166	G-protein beta3 subunit C825T polymorphism: no association with risk for hypertension and obesity. <i>Journal of Hypertension</i> , 2001 , 19, 2149-55	1.9	41
165	Systolic blood pressure response to exercise testing is related to the risk of acute myocardial infarction in middle-aged men. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 421-8		40
164	Conditioning leisure time physical activity and cardiorespiratory fitness in sociodemographic groups of middle-ages men in eastern Finland. <i>International Journal of Epidemiology</i> , 1996 , 25, 86-93	7.8	40
163	GWAS on longitudinal growth traits reveals different genetic factors influencing infant, child, and adult BMI. <i>Science Advances</i> , 2019 , 5, eaaw3095	14.3	39
162	Multiancestry Genome-Wide Association Study of Lipid Levels Incorporating Gene-Alcohol Interactions. <i>American Journal of Epidemiology</i> , 2019 , 188, 1033-1054	3.8	39
161	Analysis by LC-MS/MS of endogenous steroids from human serum, plasma, endometrium and endometriotic tissue. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018 , 152, 165-172	3.5	39
160	Chronotropic incompetence and mortality in middle-aged men with known or suspected coronary heart disease. <i>European Heart Journal</i> , 2008 , 29, 1896-902	9.5	39
159	Physical activity and sedentary time in relation to academic achievement in children. <i>Journal of Science and Medicine in Sport</i> , 2017 , 20, 583-589	4.4	38

158	Normal values for heart rate variability parameters in children 6-8 years of age: the PANIC Study. <i>Clinical Physiology and Functional Imaging</i> , 2014 , 34, 290-6	2.4	38
157	The presentation of adrenarche is sexually dimorphic and modified by body adiposity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 3889-94	5.6	37
156	Low physical activity as a predictor for antihypertensive drug treatment in 25-64-year-old populations in eastern and south-western Finland. <i>Journal of Hypertension</i> , 2005 , 23, 293-9	1.9	37
155	Physical activity modifies the effect of SNPs in the SLC2A2 (GLUT2) and ABCC8 (SUR1) genes on the risk of developing type 2 diabetes. <i>Physiological Genomics</i> , 2007 , 31, 264-72	3.6	36
154	A quantitative trait locus on 7q31 for the changes in plasma insulin in response to exercise training: the HERITAGE Family Study. <i>Diabetes</i> , 2003 , 52, 1583-7	0.9	36
153	Determinants of serum 25-hydroxyvitamin D concentration in Finnish children: the Physical Activity and Nutrition in Children (PANIC) study. <i>British Journal of Nutrition</i> , 2016 , 115, 1080-91	3.6	36
152	Association of MBOAT7 gene variant with plasma ALT levels in children: the PANIC study. <i>Pediatric Research</i> , 2016 , 80, 651-655	3.2	34
151	New Blood Pressure-Associated Loci Identified in Meta-Analyses of 475 000 Individuals. <i>Circulation: Cardiovascular Genetics</i> , 2017 , 10,		33
150	Interaction of single nucleotide polymorphisms in ADRB2, ADRB3, TNF, IL6, IGF1R, LIPC, LEPR, and GHRL with physical activity on the risk of type 2 diabetes mellitus and changes in characteristics of the metabolic syndrome: The Finnish Diabetes Prevention Study. <i>Metabolism: Clinical and</i>	12.7	33
149	Experimental, 2008, 57, 428-36 SNPs in PPARG associate with type 2 diabetes and interact with physical activity. <i>Medicine and Science in Sports and Exercise</i> , 2008, 40, 25-33	1.2	33
148	Simultaneous analysis by LC-MS/MS of 22 ketosteroids with hydroxylamine derivatization and underivatized estradiol from human plasma, serum and prostate tissue. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2019 , 164, 642-652	3.5	33
147	Cardiorespiratory fitness in aging men and women: the DR@ EXTRA study. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2011 , 21, 679-87	4.6	32
146	Physical exercise and blood pressure with reference to the angiotensinogen M235T polymorphism. <i>Physiological Genomics</i> , 2002 , 10, 71-7	3.6	32
145	Leisure-time physical activity and rate of bone loss among peri- and postmenopausal women: a longitudinal study. <i>Bone</i> , 2001 , 29, 442-6	4.7	32
144	Novel genetic associations for blood pressure identified via gene-alcohol interaction in up to 570K individuals across multiple ancestries. <i>PLoS ONE</i> , 2018 , 13, e0198166	3.7	31
143	Multi-ancestry sleep-by-SNP interaction analysis in 126,926 individuals reveals lipid loci stratified by sleep duration. <i>Nature Communications</i> , 2019 , 10, 5121	17.4	31
142	Exercise, fitness and cognition A randomised controlled trial in older individuals: The DRQ EXTRA study. <i>European Geriatric Medicine</i> , 2010 , 1, 266-272	3	31
141	A trans-ancestral meta-analysis of genome-wide association studies reveals loci associated with childhood obesity. <i>Human Molecular Genetics</i> , 2019 , 28, 3327-3338	5.6	30

140	Cardiorespiratory fitness and metabolic syndrome in older men and women: the dose responses to Exercise Training (DR@ EXTRA) study. <i>Diabetes Care</i> , 2010 , 33, 1655-7	14.6	30
139	The current standard measure of cardiorespiratory fitness introduces confounding by body mass: the DR@ EXTRA study. <i>International Journal of Obesity</i> , 2012 , 36, 1135-40	5.5	30
138	Different forms of physical activity and cardiovascular risk factors among 24-64-year-old men and women in Finland. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 51-9		30
137	Sex-dimorphic genetic effects and novel loci for fasting glucose and insulin variability. <i>Nature Communications</i> , 2021 , 12, 24	17.4	30
136	A Low-Frequency Inactivating Variant Enriched in the Finnish Population Is Associated With Fasting Insulin Levels and Type 2 Diabetes Risk. <i>Diabetes</i> , 2017 , 66, 2019-2032	0.9	29
135	The effects of a 2-year individualized and family-based lifestyle intervention on physical activity, sedentary behavior and diet in children. <i>Preventive Medicine</i> , 2016 , 87, 81-88	4.3	28
134	Obesity and weight gain are associated with increased incidence of hyperinsulinemia in non-diabetic men. <i>Hormone and Metabolic Research</i> , 2002 , 34, 492-8	3.1	28
133	Associations of diet quality with cognition in children - the Physical Activity and Nutrition in Children Study. <i>British Journal of Nutrition</i> , 2015 , 114, 1080-7	3.6	27
132	Physical activity in adolescence and bone density in peri- and postmenopausal women: a population-based study. <i>Bone</i> , 1997 , 21, 363-7	4.7	27
131	Craniofacial morphology but not excess body fat is associated with risk of having sleep-disordered breathingthe PANIC Study (a questionnaire-based inquiry in 6-8-year-olds). <i>European Journal of Pediatrics</i> , 2012 , 171, 1747-52	4.1	26
130	Discovery of rare variants associated with blood pressure regulation through meta-analysis of 1.3 million individuals. <i>Nature Genetics</i> , 2020 , 52, 1314-1332	36.3	26
129	Measures of cardiorespiratory fitness in relation to measures of body size and composition among children. <i>Clinical Physiology and Functional Imaging</i> , 2015 , 35, 469-77	2.4	25
128	Body fat mass, lean body mass and associated biomarkers as determinants of bone mineral density in children 6-8years of age - The Physical Activity and Nutrition in Children (PANIC) study. <i>Bone</i> , 2018 , 108, 106-114	4.7	25
127	Clustering of metabolic risk factors is associated with high-normal levels of liver enzymes among 6-to 8-year-old children: the PANIC study. <i>Metabolic Syndrome and Related Disorders</i> , 2012 , 10, 337-43	2.6	25
126	Association of age and education with different types of leisure-time physical activity among 4437 Finnish adults. <i>Journal of Physical Activity and Health</i> , 2008 , 5, 242-51	2.5	25
125	Novel loci for childhood body mass index and shared heritability with adult cardiometabolic traits. <i>PLoS Genetics</i> , 2020 , 16, e1008718	6	25
124	Association of waist and hip circumference with 12-year progression of carotid intima-media thickness in elderly women. <i>International Journal of Obesity</i> , 2007 , 31, 1406-11	5.5	24
123	Longitudinal associations of physical activity and sedentary time with cardiometabolic risk factors in children. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2019 , 29, 113-123	4.6	24

122	Cardiovascular fitness and haemodynamic responses to maximal cycle ergometer exercise test in children 6-8 years of age. <i>Journal of Sports Sciences</i> , 2014 , 32, 652-9	3.6	22	
121	Low-grade inflammation and depressive symptoms as predictors of abdominal obesity. <i>Scandinavian Journal of Public Health</i> , 2012 , 40, 674-80	3	22	
120	The TNF-alpha G-308A polymorphism is associated with C-reactive protein levels: the HERITAGE Family Study. <i>Vascular Pharmacology</i> , 2006 , 44, 377-83	5.9	22	
119	C-reactive protein and metabolic syndrome in elderly women: a 12-year follow-up study. <i>Diabetes Care</i> , 2006 , 29, 931-2	14.6	22	
118	Arginine intake, blood pressure, and the incidence of acute coronary events in men: the Kuopio Ischaemic Heart Disease Risk Factor Study. <i>American Journal of Clinical Nutrition</i> , 2002 , 76, 359-64	7	21	
117	Consortium-based genome-wide meta-analysis for childhood dental caries traits. <i>Human Molecular Genetics</i> , 2018 , 27, 3113-3127	5.6	20	
116	Diet quality and academic achievement: a prospective study among primary school children. <i>European Journal of Nutrition</i> , 2017 , 56, 2299-2308	5.2	20	
115	Protein tyrosine phosphatase 1B variant associated with fat distribution and insulin metabolism. <i>Obesity</i> , 2005 , 13, 829-34		19	
114	Development of cortical motor circuits between childhood and adulthood: A navigated TMS-HdEEG study. <i>Human Brain Mapping</i> , 2017 , 38, 2599-2615	5.9	18	
113	Physical activity, sedentary behaviour, and socioeconomic status among Finnish girls and boys aged 6-8 years. <i>European Journal of Sport Science</i> , 2017 , 17, 462-472	3.9	18	
112	Determinants of pain and functioning in hip osteoarthritis - a two-year prospective study. <i>Clinical Rehabilitation</i> , 2013 , 27, 281-7	3.3	18	
111	Leisure-time physical activity, cardiorespiratory fitness and feelings of hopelessness in men. <i>BMC Public Health</i> , 2009 , 9, 204	4.1	18	
110	Associations of Physical Performance and Adiposity with Cognition in Children. <i>Medicine and Science in Sports and Exercise</i> , 2015 , 47, 2166-74	1.2	17	
109	Diet, fitness and metabolic syndromethe DR@ EXTRA study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2012 , 22, 553-60	4.5	17	
108	Usefulness of chronotropic incompetence in response to exercise as a predictor of myocardial infarction in middle-aged men without cardiovascular disease. <i>American Journal of Cardiology</i> , 2008 , 101, 992-8	3	17	
107	Religious affiliation and all-cause mortality: a prospective population study in middle-aged men in eastern Finland. <i>International Journal of Epidemiology</i> , 1996 , 25, 1244-9	7.8	17	
106	Abdominal adiposity and cardiometabolic risk factors in children and adolescents: a Mendelian randomization analysis. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1079-1087	7	16	
105	Prevalence and associated factors of abnormal liver values in children with celiac disease. <i>Digestive</i> and Liver Disease, 2016 , 48, 1023-9	3.3	16	

104	Cross-sectional associations of food consumption with plasma fatty acid composition and estimated desaturase activities in Finnish children. <i>Lipids</i> , 2014 , 49, 467-79	1.6	16
103	Clinical signs of temporomandibular disorders and various pain conditions among children 6 to 8 years of age: the PANIC study. <i>Journal of Orofacial Pain</i> , 2012 , 26, 17-25		16
102	Cardiorespiratory fitness and exercise-induced ST segment depression in assessing the risk of sudden cardiac death in men. <i>Heart</i> , 2017 , 103, 383-389	5.1	15
101	Associations of physical activity, sedentary time, and cardiorespiratory fitness with heart rate variability in 6- to 9-year-old children: the PANIC study. <i>European Journal of Applied Physiology</i> , 2019 , 119, 2487-2498	3.4	15
100	Cardiorespiratory fitness, respiratory function and hemodynamic responses to maximal cycle ergometer exercise test in girls and boys aged 9-11 years: the PANIC Study. <i>European Journal of Applied Physiology</i> , 2015 , 115, 235-43	3.4	15
99	Development of corticospinal motor excitability and cortical silent period from mid-childhood to adulthood - a navigated TMS study. <i>Neurophysiologie Clinique</i> , 2018 , 48, 65-75	2.7	15
98	Metabolic risk factors are associated with stiffness index, reflection index and finger skin temperature in childrenPhysical Activity and Nutrition in Children (PANIC) study. <i>Circulation Journal</i> , 2013 , 77, 1281-8	2.9	15
97	Association of exercise-induced, silent ST-segment depression with the risk of stroke and cardiovascular diseases in men. <i>Stroke</i> , 2003 , 34, 1760-5	6.7	15
96	A multi-ancestry genome-wide study incorporating gene-smoking interactions identifies multiple new loci for pulse pressure and mean arterial pressure. <i>Human Molecular Genetics</i> , 2019 , 28, 2615-2633	5.6	14
95	The 148 M allele of the PNPLA3 is associated with plasma irisin levels in a population sample of Caucasian children: The PANIC Study. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 793-6	12.7	14
94	Two-minute heart rate recovery after cycle ergometer exercise and all-cause mortality in middle-aged men. <i>Journal of Internal Medicine</i> , 2011 , 270, 589-96	10.8	14
93	Association of leisure time physical activity and abdominal obesity with fasting serum insulin and 2-h postchallenge plasma glucose levels. <i>Diabetic Medicine</i> , 2006 , 23, 1025-8	3.5	14
92	Associations of Sedentary Behavior, Physical Activity, Cardiorespiratory Fitness, and Body Fat Content With Pain Conditions in Children: The Physical Activity and Nutrition in Children Study. <i>Journal of Pain</i> , 2016 , 17, 845-53	5.2	14
91	Associations of Objectively Measured Physical Activity and Sedentary Time With Arterial Stiffness in Pre-Pubertal Children. <i>Pediatric Exercise Science</i> , 2017 , 29, 326-335	2	13
90	The rs1800629 polymorphism in the TNF gene interacts with physical activity on the changes in C-reactive protein levels in the Finnish Diabetes Prevention Study. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2010 , 118, 757-9	2.3	13
89	Cardiorespiratory fitness and risk of disability pension: a prospective population based study in Finnish men. <i>Occupational and Environmental Medicine</i> , 2003 , 60, 765-9	2.1	13
88	Cardiac adrenergic denervation in patients with non-Q-wave versus Q-wave myocardial infarction. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2000 , 27, 816-21	8.8	13
87	Associations between estimated aerobic fitness and cardiovascular risk factors in adults with different levels of abdominal obesity. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2005 , 12, 126-31		13

(2005-2019)

86	Peak oxygen uptake cut-points to identify children at increased cardiometabolic risk - The PANIC Study. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2019 , 29, 16-24	4.6	13
85	Digitally supported program for type 2 diabetes risk identification and risk reduction in real-world setting: protocol for the StopDia model and randomized controlled trial. <i>BMC Public Health</i> , 2019 , 19, 255	4.1	12
84	An Overview on the Associations between Health Behaviors and Brain Health in Children and Adolescents with Special Reference to Diet Quality. <i>International Journal of Environmental Research and Public Health</i> , 2020 , 17,	4.6	12
83	Associations of Self-rated Fitness and Different Types of Leisure Time Physical Activity With Predicted Aerobic Fitness in 5979 Finnish Adults. <i>Journal of Physical Activity and Health</i> , 2004 , 1, 142-1	53 ^{2.5}	12
82	VO2max/kg is expected to be lower in obese individuals!. <i>International Journal of Cardiology</i> , 2015 , 189, 234	3.2	11
81	The value of cardiorespiratory fitness and exercise-induced ST segment depression in predicting death from coronary heart disease. <i>International Journal of Cardiology</i> , 2015 , 196, 31-3	3.2	11
80	Associations of TM6SF2 167K allele with liver enzymes and lipid profile in children: the PANIC Study. <i>Pediatric Research</i> , 2016 , 79, 684-8	3.2	11
79	Cross-sectional associations of plasma fatty acid composition and estimated desaturase and elongase activities with cardiometabolic risk in Finnish childrenThe PANIC study. <i>Journal of Clinical Lipidology</i> , 2016 , 10, 82-91	4.9	11
78	Secular trends affect timing of emergence of permanent teeth. <i>Angle Orthodontist</i> , 2016 , 86, 53-8	2.6	11
77	Associations of Dehydroepiandrosterone Sulfate With Cardiometabolic Risk Factors in Prepubertal Children. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 2592-2600	5.6	11
76	Leisure-time physical activity is associated with the metabolic syndrome in type 1 diabetes: effect of the PPARgamma Pro12Ala polymorphism: the FinnDiane Study. <i>Diabetes Care</i> , 2007 , 30, 1618-20	14.6	11
75	Quantitative trait locus on chromosome 20q13 for plasma levels of C-reactive protein in healthy whites: the HERITAGE Family Study. <i>Physiological Genomics</i> , 2006 , 27, 103-7	3.6	11
74	Cardiac adrenergic innervation within the first 3 months after acute myocardial infarction. <i>Clinical Physiology</i> , 2000 , 20, 366-73		11
73	Determinants for craniofacial pains in children 6-8 years of age: the PANIC study. <i>Acta Odontologica Scandinavica</i> , 2017 , 75, 453-460	2.2	10
72	High Leisure-Time Physical Activity Is Associated With Reduced Risk of Sudden Cardiac Death Among Men With Low Cardiorespiratory Fitness. <i>Canadian Journal of Cardiology</i> , 2018 , 34, 288-294	3.8	9
71	Adiposity, physical activity and neuromuscular performance in children. <i>Journal of Sports Sciences</i> , 2016 , 34, 1699-706	3.6	9
70	Effectiveness of workload at the heart rate of 100 beats/min in predicting cardiovascular mortality in men aged 42, 48, 54, or 60 years at baseline. <i>American Journal of Cardiology</i> , 2007 , 100, 563-8	3	9
69	Associations of Body Composition and Physical Activity with Balance and Walking Ability in the Elderly. <i>Journal of Physical Activity and Health</i> , 2005 , 2, 298-306	2.5	9

68	Cardiorespiratory Fitness, Physical Activity, and Insulin Resistance in Children. <i>Medicine and Science in Sports and Exercise</i> , 2020 , 52, 1144-1152	1.2	9
67	Interactions between genetic variation and cellular environment in skeletal muscle gene expression. <i>PLoS ONE</i> , 2018 , 13, e0195788	3.7	9
66	Lateral facial profile may reveal the risk for sleep disordered breathing in childrenthe PANIC-study. <i>Acta Odontologica Scandinavica</i> , 2015 , 73, 550-5	2.2	8
65	Sedentary lifestyle and emergence of hopelessness in middle-aged men. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2010 , 17, 524-9		8
64	Lifestyle management in the metabolic syndrome. <i>Metabolic Syndrome and Related Disorders</i> , 2006 , 4, 270-86	2.6	8
63	Functional and structural asymmetry in primary motor cortex in Asperger syndrome: a navigated TMS and imaging study. <i>Brain Topography</i> , 2019 , 32, 504-518	4.3	7
62	Effect of a 2-y dietary and physical activity intervention on plasma fatty acid composition and estimated desaturase and elongase activities in children: the Physical Activity and Nutrition in Children Study. <i>American Journal of Clinical Nutrition</i> , 2016 , 104, 964-972	7	7
61	Food sources of energy and nutrients in Finnish girls and boys 6-8 years of age - the PANIC study. <i>Food and Nutrition Research</i> , 2016 , 60, 32444	3.1	7
60	Physical activity, physical fitness, and risk of type 2 diabetes mellitus. <i>Metabolic Syndrome and Related Disorders</i> , 2005 , 3, 35-44	2.6	7
59	Associations between apolipoprotein E phenotype, glucose metabolism and cognitive function in men. An explorative study in a population sample. <i>Diabetic Medicine</i> , 2001 , 18, 991-7	3.5	7
58	Exercise, genetics and prevention of type 2 diabetes. Essays in Biochemistry, 2006, 42, 177-92	7.6	7
57	Maturation changes the excitability and effective connectivity of the frontal lobe: A developmental TMS-EEG study. <i>Human Brain Mapping</i> , 2019 , 40, 2320-2335	5.9	7
56	Associations of Cardiorespiratory Fitness and Adiposity With Arterial Stiffness and Arterial Dilatation Capacity in Response to a Bout of Exercise in Children. <i>Pediatric Exercise Science</i> , 2019 , 31, 238-247	2	6
55	Longitudinal Associations of Fitness, Motor Competence, and Adiposity with Cognition. <i>Medicine and Science in Sports and Exercise</i> , 2019 , 51, 465-471	1.2	6
54	Mediating effects of motor performance, cardiorespiratory fitness, physical activity, and sedentary behaviour on the associations of adiposity and other cardiometabolic risk factors with academic achievement in children. <i>Journal of Sports Sciences</i> , 2018 , 36, 2296-2303	3.6	6
53	Predictors of sleep disordered breathing in children: the PANIC study. <i>European Journal of Orthodontics</i> , 2018 , 40, 268-272	3.3	6
52	Cardiorespiratory fitness modifies the association between dietary fat intake and plasma fatty acids. <i>European Journal of Clinical Nutrition</i> , 2003 , 57, 810-5	5.2	6
51	Exercise, diet, and cognition in a 4-year randomized controlled trial: Dose-Responses to Exercise Training (DR@ EXTRA). <i>American Journal of Clinical Nutrition</i> , 2021 , 113, 1428-1439	7	6

(2018-2019)

50	Adiposity Criteria in Assessing Increased Cardiometabolic Risk in Prepubertal Children. <i>Frontiers in Endocrinology</i> , 2019 , 10, 410	5.7	5
49	Genetic predisposition to higher body fat yet lower cardiometabolic risk in children and adolescents. <i>International Journal of Obesity</i> , 2019 , 43, 2007-2016	5.5	5
48	Reproducibility of pulse contour analysis in children before and after maximal exercise stress test: the Physical Activity and Nutrition in Children (PANIC) study. <i>Clinical Physiology and Functional Imaging</i> , 2011 , 31, 132-8	2.4	5
47	A 21year physical activity and dietary intervention attenuates the increase in insulin resistance in a general population of children: the PANIC study. <i>Diabetologia</i> , 2020 , 63, 2270-2281	10.3	5
46	Health-related correlates of psychological well-being among girls and boys 6-8 years of age: The Physical Activity and Nutrition in Children study. <i>Journal of Paediatrics and Child Health</i> , 2018 , 54, 506-50	1 9 3	4
45	Tissue-Specific Alteration of Metabolic Pathways Influences Glycemic Regulation		4
44	Eating Competence Is Associated with Lower Prevalence of Obesity and Better Insulin Sensitivity in Finnish Adults with Increased Risk for Type 2 Diabetes: The StopDia Study. <i>Nutrients</i> , 2019 , 12,	6.7	4
43	Associations of cardiometabolic risk factors with heart rate variability in 6- to 8-year-old children: The PANIC Study. <i>Pediatric Diabetes</i> , 2020 , 21, 251-258	3.6	4
42	Associations of lifestyle factors with serum dehydroepiandrosterone sulphate and insulin-like growth factor-1 concentration in prepubertal children. <i>Clinical Endocrinology</i> , 2018 , 88, 234-242	3.4	4
41	Gene-educational attainment interactions in a multi-ancestry genome-wide meta-analysis identify novel blood pressure loci. <i>Molecular Psychiatry</i> , 2021 , 26, 2111-2125	15.1	3
40	Determinants of Cardiorespiratory Fitness in a Population Sample of Girls and Boys Aged 6 to 8 Years. <i>Journal of Physical Activity and Health</i> , 2016 , 13, 1149-1155	2.5	3
39	Obesity is an important source of bias in the assessment of cardiorespiratory fitness. <i>American Heart Journal</i> , 2015 , 170, e7-8	4.9	3
38	Moderators of Maintained Increase in Aerobic Exercise Among Aging Men and Women in a 4-Year Randomized Controlled Trial: The DRQ EXTRA Study. <i>Journal of Physical Activity and Health</i> , 2015 , 12, 1477-84	2.5	3
37	Associations of physical activity, sedentary time, and diet quality with biomarkers of inflammation in children. <i>European Journal of Sport Science</i> , 2021 , 1-10	3.9	3
36	Multi-ancestry genome-wide gene-sleep interactions identify novel loci for blood pressure. <i>Molecular Psychiatry</i> , 2021 ,	15.1	3
35	Response: food fortification as a means to increase vitamin D intake. <i>British Journal of Nutrition</i> , 2016 , 116, 1134-5	3.6	3
34	Associations of dietary carbohydrate and fatty acid intakes with cognition among children. <i>Public Health Nutrition</i> , 2020 , 23, 1657-1663	3.3	2
33	Serum 25-Hydroxyvitamin D, Plasma Lipids, and Associated Gene Variants in Prepubertal Children. Journal of Clinical Endocrinology and Metabolism, 2018, 103, 2670-2679	5.6	2

32	Associations of IGF-1 and Adrenal Androgens with Cognition in Childhood. <i>Hormone Research in Paediatrics</i> , 2019 , 91, 329-335	3.3	2
31	Different forms of physical activity and cardiovascular risk factors among 24 B 4-year-old men and women in Finland. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 51-59		2
30	Cross-country skiing and the risk of acute myocardial infarction: A prospective cohort study. <i>European Journal of Preventive Cardiology</i> , 2020 , 27, 1108-1111	3.9	2
29	Primary hand motor representation areas in healthy children, preadolescents, adolescents, and adults. <i>NeuroImage</i> , 2021 , 228, 117702	7.9	2
28	Comparison of Communication Channels for Large-Scale Type 2 Diabetes Risk Screening and Intervention Recruitment: Empirical Study. <i>JMIR Diabetes</i> , 2021 , 6, e21356	2.7	2
27	Child-related and parental predictors for thelarche in a general population of girls: the PANIC study. <i>Pediatric Research</i> , 2020 , 88, 676-680	3.2	1
26	Birth weight is associated with dietary factors at the age of 6-8 years: the Physical Activity and Nutrition in Children (PANIC) study. <i>Public Health Nutrition</i> , 2018 , 21, 1278-1285	3.3	1
25	Comparison between parameters from maximal cycle ergometer test first without respiratory gas analysis and thereafter with respiratory gas analysis among healthy prepubertal children. <i>Applied Physiology, Nutrition and Metabolism</i> , 2016 , 41, 624-30	3	1
24	Exercise and inflammation: reply. European Heart Journal, 2006, 27, 1385-1386	9.5	1
23	Protein-Coding Variants Implicate Novel Genes Related to Lipid Homeostasis Contributing to Body Fat Distribution		1
22	Multi-ancestry analysis of gene-sleep interactions in 126,926 individuals identifies multiple novel blood lipid loci that contribute to our understanding of sleep-associated adverse blood lipid profile		1
21	Longitudinal associations of physical activity, sedentary time, and cardiorespiratory fitness with arterial health in children - the PANIC study. <i>Journal of Sports Sciences</i> , 2021 , 39, 1980-1987	3.6	1
20	The effects of a 2-year physical activity and dietary intervention on plasma lipid concentrations in children: the PANIC Study. <i>European Journal of Nutrition</i> , 2021 , 60, 425-434	5.2	1
19	Relation of oxygen uptake to work rate in prepubertal healthy children - reference for VO /W-slope and effect on cardiorespiratory fitness assessment. <i>Clinical Physiology and Functional Imaging</i> , 2018 , 38, 645-651	2.4	1
18	Digitally Supported Lifestyle Intervention to Prevent Type 2 Diabetes Through Healthy Habits: Secondary Analysis of Long-Term User Engagement Trajectories in a Randomized Controlled Trial <i>Journal of Medical Internet Research</i> , 2022 , 24, e31530	7.6	1
17	LongITools: Dynamic longitudinal exposome trajectories in cardiovascular and metabolic noncommunicable diseases <i>Environmental Epidemiology</i> , 2022 , 6, e184	0.2	1
16	Associations between cardiorespiratory fitness, motor competence, and adiposity in children. <i>Translational Sports Medicine</i> , 2021 , 4, 56-64	1.3	0
15	A quantitative ultra-performance liquid chromatography high-resolution mass spectrometry analysis of steroids from human scalp hair <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2022 , 215, 114768	3.5	O

LIST OF PUBLICATIONS

14	The Composition and Functional Capacities of Saliva Microbiota Differ Between Children With Low and High Sweet Treat Consumption <i>Frontiers in Nutrition</i> , 2022 , 9, 864687	6.2	О
13	Cardiorespiratory Fitness as a Prognostic Factor in Heart Failure Needs to be Standardized for Body Composition. <i>American Journal of Cardiology</i> , 2015 , 116, 164	3	
12	Finger skin temperatures in 8- to 11-year-old children: determinants including physical characteristics and seasonal variation. The Physical Activity and Nutrition in Children (PANIC) Study. <i>European Journal of Applied Physiology</i> , 2016 , 116, 405-13	3.4	
11	Accuracy of Cardiorespiratory Fitness and Adiposity to Discriminate Elevated Cardiometabolic Risk Among Prepubertal Children. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 731	1.2	
10	Sedentary Lifestyle And Emergence Of Hopelessness In Middle-aged Men. <i>Medicine and Science in Sports and Exercise</i> , 2010 , 42, 58	1.2	
9	Enterolactone and coronary events. <i>Lancet, The</i> , 2000 , 355, 1643	40	
8	PHYSICAL ACTIVITY AS PREDICTOR OF TREATED HYPERTENSION IN MIDDLE-AGED POPULATIONS IN EASTERN FINLAND. <i>Medicine and Science in Sports and Exercise</i> , 2003 , 35, S67	1.2	
7	Aerobic Physical Exercise and Atherosclerosis in Men. <i>Annals of Internal Medicine</i> , 2004 , 141, 890	8	
6	Is there a duty to participate in a health research? A viewpoint of children 68 years of age and their parents. <i>International Diabetes Nursing</i> , 2016 , 13, 49-54		
5	Associations of Genetic Susceptibility to Alzheimer@ Disease with Adiposity and Cardiometabolic Risk Factors among Children in a 2-Year Follow-up Study. <i>Journal of Alzheimerm Disease</i> , 2018 , 64, 587-5	9 5 3	
4	Dental caries among Finnish teenagers participating in physical activity and diet intervention: association with anthropometrics and behavioural factors. <i>BMC Oral Health</i> , 2021 , 21, 333	3.7	
3	Cost-effectiveness of physical activity intervention in children - results based on the Physical Activity and Nutrition in Children (PANIC) study. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2021 , 18, 116	8.4	
2	The Mediating Role of Endocrine Factors in the Positive Relationship Between Fat Mass and Bone Mineral Content in Children Aged 9-11 Years: The Physical Activity and Nutrition in Children Study <i>Frontiers in Endocrinology</i> , 2022 , 13, 850448	5.7	
1	Definitions of the Metabolic Syndrome R eply. <i>JAMA - Journal of the American Medical Association</i> , 2003 , 289, 1241	27.4	