

Fiona Furlong

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6800704/publications.pdf>

Version: 2024-02-01

27
papers

1,191
citations

516561

16
h-index

501076

28
g-index

28
all docs

28
docs citations

28
times ranked

2068
citing authors

#	ARTICLE	IF	CITATIONS
1	The fate of chemoresistance in triple negative breast cancer (TNBC). <i>BBA Clinical</i> , 2015, 3, 257-275.	4.1	293
2	Overexpression of the microRNA miR-433 promotes resistance to paclitaxel through the induction of cellular senescence in ovarian cancer cells. <i>Cancer Medicine</i> , 2015, 4, 745-758.	1.3	132
3	Cisplatin induces the release of extracellular vesicles from ovarian cancer cells that can induce invasiveness and drug resistance in bystander cells. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2018, 373, 20170065.	1.8	90
4	Connective tissue growth factor [CTGF]/CCN2 stimulates mesangial cell migration through integrated dissolution of focal adhesion complexes and activation of cell polarization. <i>FASEB Journal</i> , 2004, 18, 1541-1543.	0.2	71
5	Low MAD2 expression levels associate with reduced progression-free survival in patients with high-grade serous epithelial ovarian cancer. <i>Journal of Pathology</i> , 2012, 226, 746-755.	2.1	64
6	Organization of mammary epithelial cells into 3D acinar structures requires glucocorticoid and JNK signaling. <i>Journal of Cell Biology</i> , 2004, 166, 133-143.	2.3	59
7	IHG-1 Amplifies TGF- β 1 Signaling and Is Increased in Renal Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2008, 19, 1672-1680.	3.0	57
8	FKBP1 and its peptide derivatives inhibit endocrine therapy resistant cancer stem cells and breast cancer metastasis by downregulating DLL4 and Notch4. <i>BMC Cancer</i> , 2019, 19, 351.	1.1	45
9	Cellular senescence induced by aberrant MAD2 levels impacts on paclitaxel responsiveness in vitro. <i>British Journal of Cancer</i> , 2009, 101, 1900-1908.	2.9	44
10	Autophagic secretion of HMGB1 from cancer-associated fibroblasts promotes metastatic potential of non-small cell lung cancer cells via NF- κ B signaling. <i>Cell Death and Disease</i> , 2021, 12, 858.	2.7	39
11	Connective tissue growth factor/CCN2 stimulates actin disassembly through Akt/protein kinase B-mediated phosphorylation and cytoplasmic translocation of p27 Kip1. <i>FASEB Journal</i> , 2006, 20, 1712-1714.	0.2	38
12	FKBP1-based peptide, ALM201, targets angiogenesis and cancer stem cells in ovarian cancer. <i>British Journal of Cancer</i> , 2020, 122, 361-371.	2.9	38
13	IHG-1 Promotes Mitochondrial Biogenesis by Stabilizing PGC-1 α . <i>Journal of the American Society of Nephrology: JASN</i> , 2011, 22, 1475-1485.	3.0	36
14	HDAC6 Degradation Inhibits the Growth of High-Grade Serous Ovarian Cancer Cells. <i>Cancers</i> , 2020, 12, 3734.	1.7	22
15	Dysregulated intracellular signaling impairs CTGF-stimulated responses in human mesangial cells exposed to high extracellular glucose. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 292, F1691-F1700.	1.3	19
16	Too MAD or not MAD enough: The duplicitous role of the spindle assembly checkpoint protein MAD2 in cancer. <i>Cancer Letters</i> , 2020, 469, 11-21.	3.2	18
17	IHG-1 Increases Mitochondrial Fusion and Bioenergetic Function. <i>Diabetes</i> , 2014, 63, 4314-4325.	0.3	17
18	IHG-1 must be localised to mitochondria to decrease Smad7 expression and amplify TGF- β 1-induced fibrotic responses. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013, 1833, 1969-1978.	1.9	16

#	ARTICLE	IF	CITATIONS
19	Spindle assembly checkpoint protein expression correlates with cellular proliferation and shorter time to recurrence in ovarian cancer. <i>Human Pathology</i> , 2014, 45, 1509-1519.	1.1	16
20	FKBPL: a marker of good prognosis in breast cancer. <i>Oncotarget</i> , 2015, 6, 12209-12223.	0.8	13
21	Recurrence of Urothelial Carcinoma of the Bladder: A Role for Insulin-Like Growth Factor-II Loss of Imprinting and Cytoplasmic E-Cadherin Immunolocalization. <i>Clinical Cancer Research</i> , 2008, 14, 6829-6838.	3.2	11
22	Hypoxia-targeted cupric-tirapazamine liposomes potentiate radiotherapy in prostate cancer spheroids. <i>International Journal of Pharmaceutics</i> , 2021, 607, 121018.	2.6	11
23	PTPase inhibition restores ERK1/2 phosphorylation and protects mammary epithelial cells from apoptosis. <i>Biochemical and Biophysical Research Communications</i> , 2005, 336, 1292-1299.	1.0	10
24	MAD2 downregulation in hypoxia is independent of promoter hypermethylation. <i>Cell Cycle</i> , 2010, 9, 2928-2937.	1.3	9
25	The association between MAD2 and prognosis in cancer: a systematic review and meta-analyses. <i>Oncotarget</i> , 2017, 8, 102223-102234.	0.8	9
26	MAD2 downregulation in hypoxia is independent of promoter hypermethylation. <i>Cell Cycle</i> , 2010, 9, 2856-65.	1.3	7
27	BRCA1 and MAD2 Are Coexpressed and Are Prognostic Indicators in Tubo-ovarian High-Grade Serous Carcinoma. <i>International Journal of Gynecological Cancer</i> , 2018, 28, 472-478.	1.2	6