# Mitsuhiro Takeno

## List of Publications by Year
in descending order

| 72 | 4,405 | 236925 | 118850 |
|---|---|---|---|
| papers | citations | 25 | 62 |
| | | h-index | g-index |

| 72 | 72 | 72 | 4372 |
|---|---|---|---|
| all docs | docs citations | times ranked | citing authors |

| # | Article | IF | Citations |
|---|---------|-----|-----------|
| 1 | Behçet's Disease. New England Journal of Medicine, 1999, 341, 1284-1291. | 27.0 | 1,636 |
| 2 | Genome-wide association analysis identifies new susceptibility loci for Behçet's disease and epistasis between HLA-B*51 and ERAP1. Nature Genetics, 2013, 45, 202-207. | 21.4 | 483 |
| 3 | Genome-wide association studies identify IL23R-IL12RB2 and IL10 as Behçet's disease susceptibility loci. Nature Genetics, 2010, 42, 703-706. | 21.4 | 476 |
| 4 | Incidence and Risk Factors for Serious Infection in Patients with Rheumatoid Arthritis Treated with Tumor Necrosis Factor Inhibitors: A Report from the Registry of Japanese Rheumatoid Arthritis Patients for Longterm Safety. Journal of Rheumatology, 2011, 38, 1258-1264. | 2.0 | 141 |
| 5 | Targeted resequencing implicates the familial Mediterranean fever gene <i>MEFV</i> and the toll-like receptor 4 gene <i>TLR4</i> in Behçet disease. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 8134-8139. | 7.1 | 140 |
| 6 | Behçet Disease. Medicine (United States), 2011, 90, 125-132. | 1.0 | 129 |
| 7 | Trial of Apremilast for Oral Ulcers in Behçet's Syndrome. New England Journal of Medicine, 2019, 381, 1918-1928. | 27.0 | 125 |
| 8 | The 2nd edition of consensus statements for the diagnosis and management of intestinal Behçet's disease: indication of anti-TNFα monoclonal antibodies. Journal of Gastroenterology, 2014, 49, 156-162. | 5.1 | 105 |
| 9 | Clinical manifestations of Behçet's disease depending on sex and age: results from Japanese nationwide registration. Rheumatology, 2017, 56, 1918-1927. | 1.9 | 60 |
| 10 | JCS 2017 Guideline on Management of Vasculitis Syndrome―― Digest Version ―. Circulation Journal, 2020, 84, 299-359. | 1.6 | 59 |
| 11 | The predictive prognostic factors for polymyositis/dermatomyositis-associated interstitial lung disease. Arthritis Research and Therapy, 2018, 20, 7. | 3.5 | 54 |
| 12 | Heme oxygenase 1 attenuates the development of atopic dermatitis―like lesions in mice: Implications for human disease. Journal of Allergy and Clinical Immunology, 2008, 122, 290-297.e8. | 2.9 | 51 |
| 13 | Continuous evolution of clinical phenotype in 578 Japanese patients with Behçet's disease: a retrospective observational study. Arthritis Research and Therapy, 2016, 18, 217. | 3.5 | 51 |
| 14 | Immune Checkpoint Inhibitor-Induced Myositis: a Case Report and Literature Review. Current Rheumatology Reports, 2019, 21, 10. | 4.7 | 49 |
| 15 | Clinical characteristics of neuro-Behcet's disease in Japan: a multicenter retrospective analysis. Modern Rheumatology, 2012, 22, 405-413. | 1.8 | 48 |
| 16 | Power Doppler ultrasonography is useful for assessing disease activity and predicting joint destruction in rheumatoid arthritis patients receiving tocilizumab―preliminary data. Rheumatology International, 2012, 32, 1327-1333. | 3.0 | 42 |
| 17 | Dysregulated heme oxygenase-1low M2-like macrophages augment lupus nephritis via Bach1 induced by type I interferons. Arthritis Research and Therapy, 2018, 20, 64. | 3.5 | 42 |
| 18 | Evidence-based diagnosis and clinical practice guidelines for intestinal Behçet's disease 2020 edited by Intractable Diseases, the Health and Labour Sciences Research Grants. Journal of Gastroenterology, 2020, 55, 679-700. | 5.1 | 39 |

| # | Article | IF | Citations |
|---|---------|----|-----------|
| 19 | Association of reduced heme oxygenase-1 with excessive Toll-like receptor 4 expression in peripheral blood mononuclear cells in Behçet's disease. Arthritis Research and Therapy, 2008, 10, R16. | 3.5 | 38 |
| 20 | Changes in the proportion of clinical clusters contribute to the phenotypic evolution of Behçet's disease in Japan. Arthritis Research and Therapy, 2021, 23, 49. | 3.5 | 35 |
| 21 | Gastrointestinal manifestations of Behçet's disease in Japan: a study of 43 patients. Rheumatology International, 2014, 34, 851-856. | 3.0 | 33 |
| 22 | Retrospective analysis of long-term outcome of chronic progressive neurological manifestations in Behcet's disease. Journal of the Neurological Sciences, 2015, 349, 143-148. | 0.6 | 32 |
| 23 | [18]F-FDG and [18]F-NaF PET/CT demonstrate coupling of inflammation and accelerated bone turnover in rheumatoid arthritis. Modern Rheumatology, 2016, 26, 180-187. | 1.8 | 31 |
| 24 | GWAS-identified CCR1 and IL10 loci contribute to M1 macrophage-predominant inflammation in Behçet's disease. Arthritis Research and Therapy, 2018, 20, 124. | 3.5 | 30 |
| 25 | Ultrasonography is a potent tool for the prediction of progressive joint destruction during clinical remission of rheumatoid arthritis. Modern Rheumatology, 2013, 23, 456-465. | 1.8 | 30 |
| 26 | Deep-inspiration breath-hold [18]F-FDG-PET/CT is useful for assessment of connective tissue disease associated interstitial pneumonia. Modern Rheumatology, 2016, 26, 121-127. | 1.8 | 29 |
| 27 | The association of Behçet's syndrome with HLA-B51 as understood in 2021. Current Opinion in Rheumatology, 2022, 34, 4-9. | 4.3 | 27 |
| 28 | Distinct clinical features between acute and chronic progressive parenchymal neuro-Behçet disease: meta-analysis. Scientific Reports, 2017, 7, 10196. | 3.3 | 26 |
| 29 | Bach1 regulates osteoclastogenesis in a mouse model via both heme oxygenase 1–dependent and heme oxygenase 1–independent pathways. Arthritis and Rheumatism, 2012, 64, 1518-1528. | 6.7 | 25 |
| 30 | Clinical value of capsule endoscopy for detecting small bowel lesions in patients with intestinal Behçet's disease. Digestive Endoscopy, 2016, 28, 179-185. | 2.3 | 22 |
| 31 | Thymoma–associated multi–organ autoimmunity: A case of graft–versus–host disease–like erythroderma complicated by Good syndrome successfully treated by thymectomy. Journal of Dermatology, 2017, 44, 830-835. | 1.2 | 22 |
| 32 | Long-term safety and efficacy of rituximab in 7 Japanese patients with ANCA-associated vasculitis. Modern Rheumatology, 2015, 25, 603-608. | 1.8 | 21 |
| 33 | A novel 8-joint ultrasound score is useful in daily practice for rheumatoid arthritis. Modern Rheumatology, 2015, 25, 379-385. | 1.8 | 20 |
| 34 | Single center study on ethnic and clinical features of Behcet's disease in Moscow, Russia. Clinical Rheumatology, 2015, 34, 321-327. | 2.2 | 20 |
| 35 | Investigation of Electronic Resistance in Lithium-Ion Batteries by AC Impedance Spectroscopy. Journal of the Electrochemical Society, 2017, 164, A3862-A3867. | 2.9 | 20 |
| 36 | Anti-glutamate receptor ε2 antibodies in psychiatric patients with anti-thyroid autoantibodies – A prevalence study in Japan. Neuroscience Letters, 2013, 534, 217-222. | 2.1 | 16 |

| # | Article | IF | Citations |
|---|---------|-----|-----------|
| 37 | Dysfunction of TRIM21 in interferon signature of systemic lupus erythematosus. Modern Rheumatology, 2018, 28, 993-1003. | 1.8 | 15 |
| 38 | The influence of HLA-B51 on clinical manifestations among Japanese patients with Behçet's disease: A nationwide survey. Modern Rheumatology, 2020, 30, 708-714. | 1.8 | 15 |
| 39 | The ocular involvement did not accompany with the genital ulcer or the gastrointestinal symptoms at the early stage of Behçet's disease. Modern Rheumatology, 2019, 29, 357-362. | 1.8 | 14 |
| 40 | Clinical Course before and after Cataract and Glaucoma Surgery under Systemic Infliximab Therapy in Patients with Behçet's Disease. Case Reports in Ophthalmology, 2011, 2, 189-192. | 0.7 | 13 |
| 41 | Positioning of apremilast in treatment of Behçet's disease. Modern Rheumatology, 2020, 30, 219-224. | 1.8 | 13 |
| 42 | Apremilast for oral ulcers associated with active Behçet's syndrome over 68 weeks: long-term results from a phase 3 randomised clinical trial. Clinical and Experimental Rheumatology, 2021, 39, 80-87. | 0.8 | 13 |
| 43 | Recommendations for the Management of Neuro-Behçet's Disease by the Japanese National Research Committee for Behçet's Disease. Internal Medicine, 2020, 59, 2359-2367. | 0.7 | 11 |
| 44 | HLA-A26 is a risk factor for Behçet's disease ocular lesions. Modern Rheumatology, 2021, 31, 214-218. | 1.8 | 11 |
| 45 | Efficacy and safety of apremilast for 3 months in Behçet's disease: A prospective observational study. Modern Rheumatology, 2021, 31, 856-861. | 1.8 | 10 |
| 46 | Clinical features of early-stage possible Behçet's disease patients with a variant-type major organ involvement in Japan. Modern Rheumatology, 2019, 29, 640-646. | 1.8 | 9 |
| 47 | Analysis of various factors on the relapse of acute neurological attacks in Behçet's disease. Modern Rheumatology, 2014, 24, 961-965. | 1.8 | 8 |
| 48 | Ultrasonography predicts achievement of Boolean remission after DAS28-based clinical remission of rheumatoid arthritis. Modern Rheumatology, 2014, 24, 590-598. | 1.8 | 8 |
| 49 | Predicting joint destruction in rheumatoid arthritis with power Doppler, anti-citrullinated peptide antibody, and joint swelling. Modern Rheumatology, 2015, 25, 842-848. | 1.8 | 7 |
| 50 | Adalimumab in the management of Beh&ccedil;et&rsquo;s disease. Therapeutics and Clinical Risk Management, 2015, 11, 611. | 2.0 | 6 |
| 51 | miR-1 is a novel biomarker for polymyositis/dermatomyositis-associated interstitial lung disease. Modern Rheumatology, 2020, 30, 878-883. | 1.8 | 6 |
| 52 | Should we reconsider the definition of elderly-onset rheumatoid arthritis in an ageing society?. Modern Rheumatology, 2022, 32, 323-329. | 1.8 | 6 |
| 53 | Vascular Involvement of Behçet's Disease. , 2015, , 79-100. | | 5 |
| 54 | Preferences of Japanese rheumatoid arthritis patients in treatment decision-making. Modern Rheumatology, 2013, 23, 891-896. | 1.8 | 5 |

| # | Article | IF | Citations |
|---|---------|-----|-----------|
| 55 | Beneficial effects of apremilast on genital ulcers, skin lesions, and arthritis in patients with Behçet's disease: A systematic review and meta-analysis. Modern Rheumatology, 2022, 32, 1153-1162. | 1.8 | 5 |
| 56 | Apremilast in a Japanese subgroup with Behçet's syndrome: Results from a Phase 3, randomised, double-blind, placebo-controlled study. Modern Rheumatology, 2022, 32, 413-421. | 1.8 | 5 |
| 57 | Non-IgG4-related Multifocal Fibrosclerosis. Internal Medicine, 2016, 55, 2497-2502. | 0.7 | 3 |
| 58 | On-demand ultrasonography assessment in the most symptomatic joint supports the 8-joint score system for management of rheumatoid arthritis patients. Modern Rheumatology, 2017, 27, 257-265. | 1.8 | 2 |
| 59 | Challenges to expanding the clinical application of musculoskeletal ultrasonography (MSUS) among rheumatologists: from a second survey in Japan. Modern Rheumatology, 2012, 22, 202-208. | 1.8 | 2 |
| 60 | Comparability of European League Against Rheumatology-Recommended Pharmacological Treatments of Oral Ulcers Associated with Behçet's Disease: A Systematic Literature Review of Randomized Controlled Trials. Open Access Rheumatology: Research and Reviews, 2020, Volume 12, 323-335. | 1.6 | 2 |
| 61 | Lymphopenia Helps Early Diagnosis of Systemic Lupus Erythematosus for Patients With Psychosis as an Initial Symptom. Psychosomatics, 2015, 56, 85-88. | 2.5 | 1 |
| 62 | <p>A Cost per Responder Model for Abatacept versus Adalimumab Among Rheumatoid Arthritis Patients with Seropositivity</p>. ClinicoEconomics and Outcomes Research, 2020, Volume 12, 589-594. | 1.9 | 1 |
| 63 | Behçet's disease and activities of daily living. Rheumatology, 2022, 61, 1133-1140. | 1.9 | 1 |
| 64 | Clinical features of Behçet's disease patients with joint symptoms in Japan: A national multicenter study. Modern Rheumatology, 2022, 32, 1146-1152. | 1.8 | 1 |
| 65 | Behçet's Disease and Colon Cancer. Internal Medicine, 2011, 50, 537-538. | 0.7 | 0 |
| 66 | Successful treatment of extensive venous thrombosis with combination of immunosuppressants and a direct Xa inhibitor in Behçet's disease. Modern Rheumatology Case Reports, 2018, 2, 204-208. | 0.7 | 0 |
| 67 | 249. IMPROVEMENTS AND CORRELATIONS IN ORAL ULCERS, DISEASE ACTIVITY, AND QOL IN BEHÇET'S SYNDROME PATIENTS TREATED WITH APREMILAST: A PHASE 3 RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED STUDY. Rheumatology, 2019, 58, . | 1.9 | 0 |
| 68 | 250. IMPROVEMENTS IN QUALITY OF LIFE IN BEHÇET'S SYNDROME PATIENTS TREATED WITH APREMILAST: A PHASE 3 RANDOMIZED, DOUBLE-BLIND, PLACEBO-CONTROLLED STUDY. Rheumatology, 2019, 58, . | 1.9 | 0 |
| 69 | Rapid progression to end-stage renal disease in a young female with mixed immunopathological features of lupus and ANCA-associated vasculitis. International Journal of Rheumatic Diseases, 2019, 22, 956-958. | 1.9 | 0 |
| 70 | Relationships between self-efficacy on health behavior and patient's assessment of rheumatoid arthritis conditions. An Official Journal of the Japan Primary Care Association, 2013, 36, 308-314. | 0.1 | 0 |
| 71 | Apremilast for oral ulcers associated with active Behçet's syndrome over 68 weeks: long-term results from a phase 3 randomised clinical trial. Clinical and Experimental Rheumatology, 2021, 39 Suppl 132, 80-87. | 0.8 | 0 |
| 72 | Severe digital ischemia as an unrecognized manifestation in patients with antisynthetase autoantibodies: Case series and systematic literature review. Journal of Scleroderma and Related Disorders, 0, , 239719832210908. | 1.7 | 0 |