

Mitsuhiro Takeno

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6803786/publications.pdf>

Version: 2024-02-01

72
papers

4,405
citations

236925

25
h-index

118850

62
g-index

72
all docs

72
docs citations

72
times ranked

4372
citing authors

#	ARTICLE	IF	CITATIONS
1	Behçet's disease and activities of daily living. <i>Rheumatology</i> , 2022, 61, 1133-1140.	1.9	1
2	The association of Behçet's syndrome with HLA-B51 as understood in 2021. <i>Current Opinion in Rheumatology</i> , 2022, 34, 4-9.	4.3	27
3	Clinical features of Behçet's disease patients with joint symptoms in Japan: A national multicenter study. <i>Modern Rheumatology</i> , 2022, 32, 1146-1152.	1.8	1
4	Beneficial effects of apremilast on genital ulcers, skin lesions, and arthritis in patients with Behçet's disease: A systematic review and meta-analysis. <i>Modern Rheumatology</i> , 2022, 32, 1153-1162.	1.8	5
5	Apremilast in a Japanese subgroup with Behçet's syndrome: Results from a Phase 3, randomised, double-blind, placebo-controlled study. <i>Modern Rheumatology</i> , 2022, 32, 413-421.	1.8	5
6	Should we reconsider the definition of elderly-onset rheumatoid arthritis in an ageing society?. <i>Modern Rheumatology</i> , 2022, 32, 323-329.	1.8	6
7	Efficacy and safety of apremilast for 3 months in Behçet's disease: A prospective observational study. <i>Modern Rheumatology</i> , 2021, 31, 856-861.	1.8	10
8	HLA-A26 is a risk factor for Behçet's disease ocular lesions. <i>Modern Rheumatology</i> , 2021, 31, 214-218.	1.8	11
9	Changes in the proportion of clinical clusters contribute to the phenotypic evolution of Behçet's disease in Japan. <i>Arthritis Research and Therapy</i> , 2021, 23, 49.	3.5	35
10	Apremilast for oral ulcers associated with active Behçet's syndrome over 68 weeks: long-term results from a phase 3 randomised clinical trial. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 132, 80-87.	0.8	0
11	Apremilast for oral ulcers associated with active Behçet's syndrome over 68 weeks: long-term results from a phase 3 randomised clinical trial. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 80-87.	0.8	13
12	The influence of HLA-B51 on clinical manifestations among Japanese patients with Behçet's disease: A nationwide survey. <i>Modern Rheumatology</i> , 2020, 30, 708-714.	1.8	15
13	miR-1 is a novel biomarker for polymyositis/dermatomyositis-associated interstitial lung disease. <i>Modern Rheumatology</i> , 2020, 30, 878-883.	1.8	6
14	Positioning of apremilast in treatment of Behçet's disease. <i>Modern Rheumatology</i> , 2020, 30, 219-224.	1.8	13
15	<p>A Cost per Responder Model for Abatacept versus Adalimumab Among Rheumatoid Arthritis Patients with Seropositivity</p>. <i>ClinicoEconomics and Outcomes Research</i> , 2020, Volume 12, 589-594.	1.9	1
16	Evidence-based diagnosis and clinical practice guidelines for intestinal Behçet's disease 2020 edited by Intractable Diseases, the Health and Labour Sciences Research Grants. <i>Journal of Gastroenterology</i> , 2020, 55, 679-700.	5.1	39
17	Recommendations for the Management of Neuro-Behçet's Disease by the Japanese National Research Committee for Behçet's Disease. <i>Internal Medicine</i> , 2020, 59, 2359-2367.	0.7	11
18	JCS 2017 Guideline on Management of Vasculitis Syndromeâ€• Digest Version â€•. <i>Circulation Journal</i> , 2020, 84, 299-359.	1.6	59

#	ARTICLE	IF	CITATIONS
19	Comparability of European League Against Rheumatology-Recommended Pharmacological Treatments of Oral Ulcers Associated with Behçet's Disease: A Systematic Literature Review of Randomized Controlled Trials. <i>Open Access Rheumatology: Research and Reviews</i> , 2020, Volume 12, 323-335.	1.6	2
20	Clinical features of early-stage possible Behçet's disease patients with a variant-type major organ involvement in Japan. <i>Modern Rheumatology</i> , 2019, 29, 640-646.	1.8	9
21	Trial of Apremilast for Oral Ulcers in Behçet's Syndrome. <i>New England Journal of Medicine</i> , 2019, 381, 1918-1928.	27.0	125
22	249. Improvements and Correlations in Oral Ulcers, Disease Activity, and QOL in Behçet's Syndrome Patients Treated with Apremilast: A Phase 3 Randomized, Double-Blind, Placebo-Controlled Study. <i>Rheumatology</i> , 2019, 58, .	1.9	0
23	250. Improvements in Quality of Life in Behçet's Syndrome Patients Treated with Apremilast: A Phase 3 Randomized, Double-Blind, Placebo-Controlled Study. <i>Rheumatology</i> , 2019, 58, .	1.9	0
24	Rapid progression to end-stage renal disease in a young female with mixed immunopathological features of lupus and ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 956-958.	1.9	0
25	Immune Checkpoint Inhibitor-Induced Myositis: a Case Report and Literature Review. <i>Current Rheumatology Reports</i> , 2019, 21, 10.	4.7	49
26	The ocular involvement did not accompany with the genital ulcer or the gastrointestinal symptoms at the early stage of Behçet's disease. <i>Modern Rheumatology</i> , 2019, 29, 357-362.	1.8	14
27	Dysfunction of TRIM21 in interferon signature of systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2018, 28, 993-1003.	1.8	15
28	Successful treatment of extensive venous thrombosis with combination of immunosuppressants and a direct Xa inhibitor in Behçet's disease. <i>Modern Rheumatology Case Reports</i> , 2018, 2, 204-208.	0.7	0
29	GWAS-identified CCR1 and IL10 loci contribute to M1 macrophage-predominant inflammation in Behçet's disease. <i>Arthritis Research and Therapy</i> , 2018, 20, 124.	3.5	30
30	Dysregulated heme oxygenase-1low M2-like macrophages augment lupus nephritis via Bach1 induced by type I interferons. <i>Arthritis Research and Therapy</i> , 2018, 20, 64.	3.5	42
31	The predictive prognostic factors for polymyositis/dermatomyositis-associated interstitial lung disease. <i>Arthritis Research and Therapy</i> , 2018, 20, 7.	3.5	54
32	Thymoma-associated multi-organ autoimmunity: A case of graft-versus-host disease-like erythroderma complicated by Good syndrome successfully treated by thymectomy. <i>Journal of Dermatology</i> , 2017, 44, 830-835.	1.2	22
33	Distinct clinical features between acute and chronic progressive parenchymal neuro-Behçet disease: meta-analysis. <i>Scientific Reports</i> , 2017, 7, 10196.	3.3	26
34	Clinical manifestations of Behçet's disease depending on sex and age: results from Japanese nationwide registration. <i>Rheumatology</i> , 2017, 56, 1918-1927.	1.9	60
35	On-demand ultrasonography assessment in the most symptomatic joint supports the 8-joint score system for management of rheumatoid arthritis patients. <i>Modern Rheumatology</i> , 2017, 27, 257-265.	1.8	2
36	Investigation of Electronic Resistance in Lithium-Ion Batteries by AC Impedance Spectroscopy. <i>Journal of the Electrochemical Society</i> , 2017, 164, A3862-A3867.	2.9	20

#	ARTICLE	IF	CITATIONS
37	Non-IgG4-related Multifocal Fibrosclerosis. <i>Internal Medicine</i> , 2016, 55, 2497-2502.	0.7	3
38	Continuous evolution of clinical phenotype in 578 Japanese patients with Behçet's disease: a retrospective observational study. <i>Arthritis Research and Therapy</i> , 2016, 18, 217.	3.5	51
39	Clinical value of capsule endoscopy for detecting small bowel lesions in patients with intestinal Behçet's disease. <i>Digestive Endoscopy</i> , 2016, 28, 179-185.	2.3	22
40	Deep-inspiration breath-hold ¹⁸ F-FDG-PET/CT is useful for assessment of connective tissue disease associated interstitial pneumonia. <i>Modern Rheumatology</i> , 2016, 26, 121-127.	1.8	29
41	¹⁸ F-FDG and ¹⁸ F-NaF PET/CT demonstrate coupling of inflammation and accelerated bone turnover in rheumatoid arthritis. <i>Modern Rheumatology</i> , 2016, 26, 180-187.	1.8	31
42	Adalimumab in the management of Behçet's disease. <i>Therapeutics and Clinical Risk Management</i> , 2015, 11, 611.	2.0	6
43	A novel 8-joint ultrasound score is useful in daily practice for rheumatoid arthritis. <i>Modern Rheumatology</i> , 2015, 25, 379-385.	1.8	20
44	Retrospective analysis of long-term outcome of chronic progressive neurological manifestations in Behçet's disease. <i>Journal of the Neurological Sciences</i> , 2015, 349, 143-148.	0.6	32
45	Predicting joint destruction in rheumatoid arthritis with power Doppler, anti-citrullinated peptide antibody, and joint swelling. <i>Modern Rheumatology</i> , 2015, 25, 842-848.	1.8	7
46	Long-term safety and efficacy of rituximab in 7 Japanese patients with ANCA-associated vasculitis. <i>Modern Rheumatology</i> , 2015, 25, 603-608.	1.8	21
47	Single center study on ethnic and clinical features of Behçet's disease in Moscow, Russia. <i>Clinical Rheumatology</i> , 2015, 34, 321-327.	2.2	20
48	Lymphopenia Helps Early Diagnosis of Systemic Lupus Erythematosus for Patients With Psychosis as an Initial Symptom. <i>Psychosomatics</i> , 2015, 56, 85-88.	2.5	1
49	Vascular Involvement of Behçet's Disease. , 2015, , 79-100.		5
50	Analysis of various factors on the relapse of acute neurological attacks in Behçet's disease. <i>Modern Rheumatology</i> , 2014, 24, 961-965.	1.8	8
51	Ultrasonography predicts achievement of Boolean remission after DAS28-based clinical remission of rheumatoid arthritis. <i>Modern Rheumatology</i> , 2014, 24, 590-598.	1.8	8
52	The 2nd edition of consensus statements for the diagnosis and management of intestinal Behçet's disease: indication of anti-TNF± monoclonal antibodies. <i>Journal of Gastroenterology</i> , 2014, 49, 156-162.	5.1	105
53	Gastrointestinal manifestations of Behçet's disease in Japan: a study of 43 patients. <i>Rheumatology International</i> , 2014, 34, 851-856.	3.0	33
54	Genome-wide association analysis identifies new susceptibility loci for Behçet's disease and epistasis between HLA-B*51 and ERAP1. <i>Nature Genetics</i> , 2013, 45, 202-207.	21.4	483

#	ARTICLE	IF	CITATIONS
55	Anti-glutamate receptor É2 antibodies in psychiatric patients with anti-thyroid autoantibodies â€“ A prevalence study in Japan. <i>Neuroscience Letters</i> , 2013, 534, 217-222.	2.1	16
56	Targeted resequencing implicates the familial Mediterranean fever gene <i>MEFV</i> and the toll-like receptor 4 gene <i>TLR4</i> in BehÃ§et disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 8134-8139.	7.1	140
57	Ultrasonography is a potent tool for the prediction of progressive joint destruction during clinical remission of rheumatoid arthritis. <i>Modern Rheumatology</i> , 2013, 23, 456-465.	1.8	30
58	Relationships between self-efficacy on health behavior and patient's assessment of rheumatoid arthritis conditions. <i>An Official Journal of the Japan Primary Care Association</i> , 2013, 36, 308-314.	0.1	0
59	Preferences of Japanese rheumatoid arthritis patients in treatment decision-making. <i>Modern Rheumatology</i> , 2013, 23, 891-896.	1.8	5
60	Bach1 regulates osteoclastogenesis in a mouse model via both heme oxygenase 1â€“dependent and heme oxygenase 1â€“independent pathways. <i>Arthritis and Rheumatism</i> , 2012, 64, 1518-1528.	6.7	25
61	Power Doppler ultrasonography is useful for assessing disease activity and predicting joint destruction in rheumatoid arthritis patients receiving tocilizumabâ€“preliminary data. <i>Rheumatology International</i> , 2012, 32, 1327-1333.	3.0	42
62	Challenges to expanding the clinical application of musculoskeletal ultrasonography (MSUS) among rheumatologists: from a second survey in Japan. <i>Modern Rheumatology</i> , 2012, 22, 202-208.	1.8	2
63	Clinical characteristics of neuro-Behcetâ€™s disease in Japan: a multicenter retrospective analysis. <i>Modern Rheumatology</i> , 2012, 22, 405-413.	1.8	48
64	Incidence and Risk Factors for Serious Infection in Patients with Rheumatoid Arthritis Treated with Tumor Necrosis Factor Inhibitors: A Report from the Registry of Japanese Rheumatoid Arthritis Patients for Longterm Safety. <i>Journal of Rheumatology</i> , 2011, 38, 1258-1264.	2.0	141
65	BehÃ§et Disease. <i>Medicine (United States)</i> , 2011, 90, 125-132.	1.0	129
66	BehÃ§et's Disease and Colon Cancer. <i>Internal Medicine</i> , 2011, 50, 537-538.	0.7	0
67	Clinical Course before and after Cataract and Glaucoma Surgery under Systemic Infliximab Therapy in Patients with BehÃ§etâ€™s Disease. <i>Case Reports in Ophthalmology</i> , 2011, 2, 189-192.	0.7	13
68	Genome-wide association studies identify IL23R-IL12RB2 and IL10 as BehÃ§et's disease susceptibility loci. <i>Nature Genetics</i> , 2010, 42, 703-706.	21.4	476
69	Association of reduced heme oxygenase-1 with excessive Toll-like receptor 4 expression in peripheral blood mononuclear cells in BehÃ§et's disease. <i>Arthritis Research and Therapy</i> , 2008, 10, R16.	3.5	38
70	Heme oxygenase 1 attenuates the development of atopic dermatitisâ€“like lesions in mice: Implications for human disease. <i>Journal of Allergy and Clinical Immunology</i> , 2008, 122, 290-297.e8.	2.9	51
71	BehÃ§et's Disease. <i>New England Journal of Medicine</i> , 1999, 341, 1284-1291.	27.0	1,636
72	Severe digital ischemia as an unrecognized manifestation in patients with antisynthetase autoantibodies: Case series and systematic literature review. <i>Journal of Scleroderma and Related Disorders</i> , 0, , 239719832210908.	1.7	0