

Sabrina Flor

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6804098/publications.pdf>

Version: 2024-02-01

16
papers

246
citations

932766

10
h-index

940134

16
g-index

17
all docs

17
docs citations

17
times ranked

324
citing authors

#	ARTICLE	IF	CITATIONS
1	Miniaturized imprinted solid phase extraction to the selective analysis of Coenzyme Q10 in urine. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2019, 1116, 24-29.	1.2	6
2	Development of an enantioselective capillary electrophoretic method for the simultaneous determination of montelukast enantiomeric and diastereoisomeric forms and its main degradation product. <i>Electrophoresis</i> , 2016, 37, 2420-2428.	1.3	16
3	Development and Validation of a CD-MEKC System for the Simultaneous Determination of Dihydrostreptomycin Sulfate and Two Benzylpenicillin Salts. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2015, 38, 955-962.	0.5	2
4	New Analytical Strategies Applied to the Determination of Coenzyme Q10 in Biological Matrix. <i>Methods in Molecular Biology</i> , 2015, 1208, 409-420.	0.4	9
5	Novel nelfinavir mesylate loaded d- α -tocopheryl polyethylene glycol 1000 succinate micelles for enhanced pediatric anti HIV therapy: In vitro characterization and in vivo evaluation. <i>Colloids and Surfaces B: Biointerfaces</i> , 2014, 123, 302-310.	2.5	34
6	Development and validation of a capillary electrophoresis method for determination of enantiomeric purity and related substances of esomeprazole in raw material and pellets. <i>Electrophoresis</i> , 2014, 35, 804-810.	1.3	20
7	The use of coenzyme Q0 as a template in the development of a molecularly imprinted polymer for the selective recognition of coenzyme Q10. <i>Analytica Chimica Acta</i> , 2014, 807, 67-74.	2.6	23
8	Development and Validation of a Highly Sensitive HPLC Method for Determination of Paclitaxel in Pharmaceutical Dosage forms and Biological Samples. <i>Current Pharmaceutical Analysis</i> , 2014, 10, 185-192.	0.3	10
9	Molecularly Imprinted Solid Phase Extraction Before Capillary Electrophoresis for the Analysis of Estrogens in Serum Samples. <i>Current Analytical Chemistry</i> , 2014, 10, 235-240.	0.6	5
10	MINIATURIZED HPLC-UV METHOD FOR ANALYSIS OF COENZYME Q10 IN HUMAN PLASMA. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2011, 34, 2485-2494.	0.5	11
11	Simultaneous determination of nine endogenous steroids in human urine by polymeric mixed micelle capillary electrophoresis. <i>Electrophoresis</i> , 2010, 31, 3305-3313.	1.3	32
12	Novel and highly sensitive mixed polymeric electrokinetic chromatography system for determination of contaminants and impurities of heparin samples. <i>Electrophoresis</i> , 2010, 31, 3606-3612.	1.3	7
13	FAST AND SENSITIVE NEW HPLC-UV METHOD FOR DETERMINATION OF OMEPRAZOLE AND MAJOR RELATED SUBSTANCES IN PHARMACEUTICAL FORMULATION. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2010, 33, 1666-1678.	0.5	16
14	A capillary electrophoretic system based on a novel microemulsion for the analysis of coenzyme Q10 in human plasma by electrokinetic chromatography. <i>Electrophoresis</i> , 2009, 30, 1899-1905.	1.3	26
15	Simple, Highly Sensitive Micro HPLC Method for the Determination of Coenzyme Q10 and its Major Related Substances. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2009, 32, 860-873.	0.5	6
16	Simultaneous determination of natural and synthetic estrogens by EKC using a novel microemulsion. <i>Electrophoresis</i> , 2006, 27, 4431-4438.	1.3	21