Hiroyasu Iso

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6817594/publications.pdf

Version: 2024-02-01

592 papers 23,849 citations

79 h-index 125 g-index

609 all docs 609 docs citations

609 times ranked 22473 citing authors

#	Article	IF	CITATIONS
1	Serum Cholesterol Levels and Six-Year Mortality from Stroke in 350,977 Men Screened for the Multiple Risk Factor Intervention Trial. New England Journal of Medicine, 1989, 320, 904-910.	13.9	1,043
2	Intake of Fish and n3 Fatty Acids and Risk of Coronary Heart Disease Among Japanese. Circulation, 2006, 113, 195-202.	1.6	496
3	Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. Clinical and Experimental Nephrology, 2007, 11, 41-50.	0.7	488
4	Intake of Fish and Omega-3 Fatty Acids and Risk of Stroke in Women. JAMA - Journal of the American Medical Association, 2001, 285, 304.	3.8	408
5	The Relationship between Green Tea and Total Caffeine Intake and Risk for Self-Reported Type 2 Diabetes among Japanese Adults. Annals of Internal Medicine, 2006, 144, 554.	2.0	389
6	What has made the population of Japan healthy?. Lancet, The, 2011, 378, 1094-1105.	6.3	381
7	Association of Sleep Duration with Mortality from Cardiovascular Disease and Other Causes for Japanese Men and Women: the JACC Study. Sleep, 2009, 32, 295-301.	0.6	311
8	Serum Triglycerides and Risk of Coronary Heart Disease among Japanese Men and Women. American Journal of Epidemiology, 2001, 153, 490-499.	1.6	294
9	Prospective Study of Calcium, Potassium, and Magnesium Intake and Risk of Stroke in Women. Stroke, 1999, 30, 1772-1779.	1.0	293
10	Carotid Intima-Media Thickness and Plaque Characteristics as a Risk Factor for Stroke in Japanese Elderly Men. Stroke, 2004, 35, 2788-2794.	1.0	262
11	Fish, I‰-3 Polyunsaturated Fatty Acids, and Mortality From Cardiovascular Diseases in a Nationwide Community-Based Cohort of Japanese Men and Women. Journal of the American College of Cardiology, 2008, 52, 988-996.	1.2	251
12	Association of Dietary Intake of Soy, Beans, and Isoflavones With Risk of Cerebral and Myocardial Infarctions in Japanese Populations. Circulation, 2007, 116, 2553-2562.	1.6	247
13	Perceived Mental Stress and Mortality From Cardiovascular Disease Among Japanese Men and Women. Circulation, 2002, 106, 1229-1236.	1.6	237
14	Cardiovascular Disease Epidemiology in Asia. Circulation Journal, 2013, 77, 1646-1652.	0.7	217
15	Linoleic Acid, Other Fatty Acids, and the Risk of Stroke. Stroke, 2002, 33, 2086-2093.	1.0	205
16	The joint impact on being overweight of self reported behaviours of eating quickly and eating until full: cross sectional survey. BMJ: British Medical Journal, 2008, 337, a2002-a2002.	2.4	196
17	Adult Mortality Attributable to Preventable Risk Factors for Non-Communicable Diseases and Injuries in Japan: A Comparative Risk Assessment. PLoS Medicine, 2012, 9, e1001160.	3.9	196
18	Coffee, green tea, black tea and oolong tea consumption and risk of mortality from cardiovascular disease in Japanese men and women. Journal of Epidemiology and Community Health, 2011, 65, 230-240.	2.0	186

#	Article	IF	Citations
19	Metabolic Syndrome and the Risk of Ischemic Heart Disease and Stroke Among Japanese Men and Women. Stroke, 2007, 38, 1744-1751.	1.0	184
20	Alcohol Consumption and Risk of Stroke Among Middle-Aged Men: The JPHC Study Cohort I. Stroke, 2004, 35, 1124-1129.	1.0	175
21	A Prospective Cohort Study of Shift Work and Risk of Ischemic Heart Disease in Japanese Male Workers. American Journal of Epidemiology, 2006, 164, 128-135.	1.6	168
22	A Low Prevalence of Coronary Heart Disease among Subjects with Increased High-Density Lipoprotein Cholesterol Levels, Including Those with Plasma Cholesteryl Ester Transfer Protein Deficiency. Preventive Medicine, 1998, 27, 659-667.	1.6	163
23	Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. American Journal of Clinical Nutrition, 2008, 88, 195-202.	2.2	162
24	Prospective Study of Fat and Protein Intake and Risk of Intraparenchymal Hemorrhage in Women. Circulation, 2001, 103, 856-863.	1.6	153
25	Trends in the Incidence of Coronary Heart Disease and Stroke and Their Risk Factors in Japan, 1964 to 2003. Journal of the American College of Cardiology, 2008, 52, 71-79.	1.2	152
26	Randomized Controlled Trial for an Effect of Green Tea Consumption on Insulin Resistance and Inflammation Markers. Journal of Nutritional Science and Vitaminology, 2005, 51, 335-342.	0.2	148
27	Daily Total Physical Activity Level and Premature Death in Men and Women: Results From a Large-Scale Population-Based Cohort Study in Japan (JPHC Study). Annals of Epidemiology, 2008, 18, 522-530.	0.9	147
28	Fruit and Vegetable Intake and Risk of Total Cancer and Cardiovascular Disease: Japan Public Health Center-based Prospective Study. American Journal of Epidemiology, 2007, 167, 59-70.	1.6	145
29	Low-Density Lipoprotein Cholesterol Concentrations and Death Due to Intraparenchymal Hemorrhage. Circulation, 2009, 119, 2136-2145.	1.6	144
30	Walking and Sports Participation and Mortality From Coronary Heart Disease and Stroke. Journal of the American College of Cardiology, 2005, 46, 1761-1767.	1.2	139
31	Smoking Cessation and Mortality from Cardiovascular Disease among Japanese Men and Women: The JACC Study. American Journal of Epidemiology, 2005, 161, 170-179.	1.6	138
32	Profile of the JACC Study. Journal of Epidemiology, 2005, 15, S4-S8.	1.1	137
33	Cohort Profile of the Japan Collaborative Cohort Study at Final Follow-up. Journal of Epidemiology, 2013, 23, 227-232.	1.1	134
34	Prospective Study of Depressive Symptoms and Risk of Stroke Among Japanese. Stroke, 2001, 32, 903-908.	1.0	133
35	Serum Total Homocysteine Concentrations and Risk of Stroke and Its Subtypes in Japanese. Circulation, 2004, 109, 2766-2772.	1.6	133
36	Fruit, vegetable and bean intake and mortality from cardiovascular disease among Japanese men and women: the JACC Study. British Journal of Nutrition, 2009, 102, 285-292.	1.2	132

#	Article	IF	CITATIONS
37	Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. European Journal of Cancer Prevention, 2009, 18, 240-247.	0.6	131
38	Lifestyle and Cardiovascular Disease in Japan. Journal of Atherosclerosis and Thrombosis, 2011, 18, 83-88.	0.9	131
39	Cigarette Smoking and Risk of Stroke and its Subtypes Among Middle-Aged Japanese Men and Women. Stroke, 2004, 35, 1248-1253.	1.0	129
40	Associations of dietary magnesium intake with mortality from cardiovascular disease: The JACC study. Atherosclerosis, 2012, 221, 587-595.	0.4	129
41	Dietary Intake of Calcium in Relation to Mortality From Cardiovascular Disease. Stroke, 2006, 37, 20-26.	1.0	125
42	Obstructive sleep apnea and incident type 2 diabetes. Sleep Medicine, 2016, 25, 156-161.	0.8	125
43	Alcohol Intake and the Risk of Cardiovascular Disease in Middle-Aged Japanese Men. Stroke, 1995, 26, 767-773.	1.0	124
44	Body Mass Index and Mortality From Cardiovascular Disease Among Japanese Men and Women. Stroke, 2005, 36, 1377-1382.	1.0	123
45	The Impact of Green Tea and Coffee Consumption on the Reduced Risk of Stroke Incidence in Japanese Population. Stroke, 2013, 44, 1369-1374.	1.0	123
46	Dietary Fiber Intake Is Associated with Reduced Risk of Mortality from Cardiovascular Disease among Japanese Men and Women , ,. Journal of Nutrition, 2010, 140, 1445-1453.	1.3	119
47	Serum Fatty Acids and Fish Intake in Rural Japanese, Urban Japanese, Japanese American and Caucasian American Men. International Journal of Epidemiology, 1989, 18, 374-381.	0.9	118
48	Serum and Dietary Magnesium and Risk of Ischemic Stroke: The Atherosclerosis Risk in Communities Study. American Journal of Epidemiology, 2009, 169, 1437-1444.	1.6	118
49	Serum cholesterol levels in relation to the incidence of cancer: The JPHC study cohorts. International Journal of Cancer, 2009, 125, 2679-2686.	2.3	118
50	Alcohol Consumption and Mortality From Stroke and Coronary Heart Disease Among Japanese Men and Women. Stroke, 2008, 39, 2936-2942.	1.0	112
51	Blood Pressure and the Risk of Stroke, Cardiovascular Disease, and All-Cause Mortality Among Japanese: The JPHC Study. American Journal of Hypertension, 2009, 22, 273-280.	1.0	105
52	Relationships of Age at Menarche and Menopause, and Reproductive Year with Mortality from Cardiovascular Disease in Japanese Postmenopausal Women: The JACC Study. Journal of Epidemiology, 2006, 16, 177-184.	1.1	104
53	Trends for Blood Pressure and Its Contribution to Stroke Incidence in the Middle-Aged Japanese Population. Stroke, 2009, 40, 1571-1577.	1.0	104
54	Blood pressure-lowering effect of Shinrin-yoku (Forest bathing): a systematic review and meta-analysis. BMC Complementary and Alternative Medicine, 2017, 17, 409.	3.7	104

#	Article	IF	CITATIONS
55	Dietary Calcium Intake and Risks of Stroke, Its Subtypes, and Coronary Heart Disease in Japanese. Stroke, 2008, 39, 2449-2456.	1.0	103
56	Egg consumption, serum total cholesterol concentrations and coronary heart disease incidence: Japan Public Health Center-based prospective study. British Journal of Nutrition, 2006, 96, 921-928.	1.2	102
57	Dietary Folate and Vitamin B ₆ and B ₁₂ Intake in Relation to Mortality From Cardiovascular Diseases. Stroke, 2010, 41, 1285-1289.	1.0	102
58	Fat and Protein Intakes and Risk of Intraparenchymal Hemorrhage among Middle-aged Japanese. American Journal of Epidemiology, 2003, 157, 32-39.	1.6	101
59	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCH–JAPAN study. Journal of Clinical Lipidology, 2018, 12, 674-684.e5.	0.6	101
60	Trends in the incidence of coronary heart disease and stroke and the prevalence of cardiovascular risk factors among Japanese men from 1963 to 1994. American Journal of Medicine, 2002, 112, 104-109.	0.6	100
61	The impact of the metabolic syndrome and its components on the incidence of ischemic heart disease and stroke: the Japan public health center-based study. Hypertension Research, 2009, 32, 289-298.	1.5	100
62	Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan Public Health Centerâ€"based Prospective Study. American Journal of Clinical Nutrition, 2010, 91, 456-464.	2.2	100
63	Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. Clinical Nutrition, 2013, 32, 300-308.	2.3	98
64	Effects of Habitual Alcohol Intake on Ambulatory Blood Pressure, Heart Rate, and Its Variability Among Japanese Men. Hypertension, 2009, 53, 13-19.	1.3	97
65	The criteria for metabolic syndrome and the national health screening and education system in Japan. Epidemiology and Health, 2017, 39, e2017003.	0.8	96
66	Dietary patterns and all-cause, cancer, and cardiovascular disease mortality in Japanese men and women: The Japan public health center-based prospective study. PLoS ONE, 2017, 12, e0174848.	1.1	96
67	Impact of the revision of a nutrient database on the validity of a self-administered food frequency questionnaire (FFQ). Journal of Epidemiology, 2006, 16, 107-116.	1.1	92
68	Changes in Coronary Heart Disease Risk Among Japanese. Circulation, 2008, 118, 2725-2729.	1.6	89
69	Mild Retinopathy Is a Risk Factor for Cardiovascular Mortality in Japanese With and Without Hypertension. Circulation, 2011, 124, 2502-2511.	1.6	89
70	Healthy lifestyle behaviours and cardiovascular mortality among Japanese men and women: the Japan collaborative cohort study. European Heart Journal, 2012, 33, 467-477.	1.0	88
71	Soft drink intake in relation to incident ischemic heart disease, stroke, and stroke subtypes in Japanese men and women: the Japan Public Health Centre–based study cohort I. American Journal of Clinical Nutrition, 2012, 96, 1390-1397.	2.2	88
72	Alcohol Consumption and Mortality among Middle-aged and Elderly Japanese Men and Women. Annals of Epidemiology, 2005, 15, 590-597.	0.9	87

#	Article	IF	CITATIONS
73	Dietary intake of saturated fatty acids and mortality from cardiovascular disease in Japanese: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. American Journal of Clinical Nutrition, 2010, 92, 759-765.	2.2	87
74	ACCURACY OF DEATH CERTIFICATE DIAGNOSIS OF INTRACRANIAL HEMORRHAGE AND NONHEMORRHAGIC STROKE. American Journal of Epidemiology, 1990, 132, 993-998.	1.6	86
75	Risk and Protective Factors Related to Mortality from Pneumonia among Middleaged and Elderly Community Residents: The JACC Study. Journal of Epidemiology, 2007, 17, 194-202.	1.1	86
76	Serum total cholesterol levels and risk of mortality from stroke and coronary heart disease in Japanese: The JACC study. Atherosclerosis, 2007, 194, 415-420.	0.4	86
77	Serum Total and Non-High-Density Lipoprotein Cholesterol and the Risk Prediction of Cardiovascular Events - The JALS-ECC Circulation Journal, 2010, 74, 1346-1356.	0.7	85
78	Application of Computer Tomography-Oriented Criteria for Stroke Subtype Classification in a Prospective Study. Annals of Epidemiology, 2000, 10, 81-87.	0.9	84
79	Relation Between Serum Total Cholesterol Level and Cardiovascular Disease Stratified by Sex and Age Group: A Pooled Analysis of 65Â594 Individuals From 10 Cohort Studies in Japan. Journal of the American Heart Association, 2012, 1, e001974.	1.6	84
80	Fasting and non-fasting triglycerides and risk of ischemic cardiovascular disease in Japanese men and women: The Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2014, 237, 361-368.	0.4	83
81	Associations between dietary intakes of iron, copper and zinc with risk of type 2 diabetes mellitus: A large population-based prospective cohort study. Clinical Nutrition, 2018, 37, 667-674.	2.3	83
82	Low-density lipoprotein cholesterol and risk of coronary heart disease among Japanese men and women: The Circulatory Risk in Communities Study (CIRCS). Preventive Medicine, 2011, 52, 381-386.	1.6	79
83	The effects of smoking and smoking cessation on mortality from cardiovascular disease among Japanese: pooled analysis of three large-scale cohort studies in Japan. Tobacco Control, 2010, 19, 50-57.	1.8	76
84	Metabolic Syndrome and All-Cause and Cardiovascular Disease Mortality Japan Public Health Center-Based Prospective (JPHC) Study. Circulation Journal, 2009, 73, 878-884.	0.7	75
85	Social Support and Stroke and Coronary Heart Disease. Stroke, 2008, 39, 768-775.	1.0	73
86	Perceived Level of Life Enjoyment and Risks of Cardiovascular Disease Incidence and Mortality. Circulation, 2009, 120, 956-963.	1.6	72
87	Diabetes Mellitus and Risk of Stroke and Its Subtypes Among Japanese. Stroke, 2011, 42, 2611-2614.	1.0	72
88	Dietary Intakes of Antioxidant Vitamins and Mortality From Cardiovascular Disease. Stroke, 2011, 42, 1665-1672.	1.0	70
89	Association of Breakfast Intake With Incident Stroke and Coronary Heart Disease. Stroke, 2016, 47, 477-481.	1.0	69
90	Underweight as a Predictor of Diabetes in Older Adults. Diabetes Care, 2008, 31, 583-584.	4.3	68

#	Article	IF	CITATIONS
91	Social support and cancer incidence and mortality: the JPHC study cohort II. Cancer Causes and Control, 2013, 24, 847-860.	0.8	68
92	Dietary intake of saturated fatty acids and incident stroke and coronary heart disease in Japanese communities: the JPHC Study. European Heart Journal, 2013, 34, 1225-1232.	1.0	66
93	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). Annals of Epidemiology, 2015, 25, 512-518.e3.	0.9	66
94	Proportions of Stroke Subtypes Among Men and Women ≥40 Years of Age in an Urban Japanese City in 1992, 1997, and 2002. Stroke, 2006, 37, 1374-1378.	1.0	65
95	Associations of All-Cause Mortality with Census-Based Neighbourhood Deprivation and Population Density in Japan: A Multilevel Survival Analysis. PLoS ONE, 2014, 9, e97802.	1.1	65
96	Effects of a Long-term Hypertension Control Program on Stroke Incidence and Prevalence in a Rural Community in Northeastern Japan. Stroke, 1998, 29, 1510-1518.	1.0	64
97	Risk Factors for Fatal Subarachnoid Hemorrhage. Stroke, 2003, 34, 2781-2787.	1.0	64
98	Usual Alcohol Consumption and Arterial Oxygen Desaturation During Sleep. JAMA - Journal of the American Medical Association, 2004, 292, 923-925.	3.8	64
99	Genetic Predisposition to Ischemic Stroke. Stroke, 2017, 48, 253-258.	1.0	64
100	Cigarette smoking and risk of coronary heart disease incidence among middle-aged Japanese men and women: the JPHC Study Cohort I. European Journal of Cardiovascular Prevention and Rehabilitation, 2006, 13, 207-213.	3.1	63
101	C-reactive protein levels and risk of mortality from cardiovascular disease in Japanese: The JACC Study. Atherosclerosis, 2009, 207, 291-297.	0.4	63
102	Body Mass Index and Risk of Stroke and Myocardial Infarction in a Relatively Lean Population. Circulation: Cardiovascular Quality and Outcomes, 2010, 3, 498-505.	0.9	62
103	Social Network, Social Support, and Risk of Incident Stroke. Stroke, 2014, 45, 2868-2873.	1.0	62
104	The Dose-Response Relationship Between Body Mass Index and the Risk of Incident Stage ^ ^ge;3 Chronic Kidney Disease in a General Japanese Population: The Ibaraki Prefectural Health Study (IPHS). Journal of Epidemiology, 2014, 24, 444-451.	1.1	62
105	Associations between copper and zinc intakes from diet and mortality from cardiovascular disease in a large population-based prospective cohort study. Journal of Nutritional Biochemistry, 2018, 56, 126-132.	1.9	62
106	Smoking Raises the Risk of Total and Ischemic Strokes in Hypertensive Men Hypertension Research, 2003, 26, 209-217.	1.5	61
107	Community-Based Education Classes for Hypertension Control. Hypertension, 1996, 27, 968-974.	1.3	61
108	Serum total cholesterol and mortality in a Japanese population. Journal of Clinical Epidemiology, 1994, 47, 961-969.	2.4	59

#	Article	IF	CITATIONS
109	Relationship between Sleep-Disordered Breathing and Blood Pressure Levels in Community-Based Samples of Japanese Men. Hypertension Research, 2004, 27, 479-484.	1.5	59
110	Rice Intake Is Associated with Reduced Risk of Mortality from Cardiovascular Disease in Japanese Men but Not Women1–3. Journal of Nutrition, 2011, 141, 595-602.	1.3	58
111	High serum total cholesterol levels is a risk factor of ischemic stroke for general Japanese population: The JPHC study. Atherosclerosis, 2012, 221, 565-569.	0.4	58
112	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2015, 101, 1029-1037.	2.2	58
113	Effect of coffee consumption on all-cause and total cancer mortality: findings from the JACC study. European Journal of Epidemiology, 2011, 26, 285-293.	2.5	55
114	Daily Total Physical Activity and Incident Stroke. Stroke, 2017, 48, 1730-1736.	1.0	55
115	Prognostic impact of supraventricular premature complexes in community-based health checkups: The Ibaraki Prefectural Health Study. European Heart Journal, 2015, 36, 170-178.	1.0	54
116	Cigarette Smoking and Risk of Disabling Dementia in a Japanese Rural Community: A Nested Case-Control Study. Cerebrovascular Diseases, 2008, 25, 324-331.	0.8	53
117	Intake of Folate, Vitamin B ₆ and Vitamin B ₁₂ and the Risk of CHD: The Japan Public Health Center-Based Prospective Study Cohort I. Journal of the American College of Nutrition, 2008, 27, 127-136.	1.1	53
118	Chronic Kidney Disease and Drinking Status in Relation to Risks of Stroke and Its Subtypes. Stroke, 2011, 42, 2531-2537.	1.0	53
119	The Association of Having a Late Dinner or Bedtime Snack and Skipping Breakfast with Overweight in Japanese Women. Journal of Obesity, 2019, 2019, 1-5.	1.1	53
120	Relations between protein intake and blood pressure in Japanese men and women: the Circulatory Risk in Communities Study (CIRCS). American Journal of Clinical Nutrition, 2009, 90, 377-384.	2.2	52
121	A low level of C-reactive protein in Japanese adults and its association with cardiovascular risk factors: The Japan NCVC-Collaborative Inflammation Cohort (JNIC) Study. Atherosclerosis, 2007, 194, 238-244.	0.4	51
122	Prognostic value of plasma high-sensitivity C-reactive protein levels in Japanese patients with stable coronary artery disease: The Japan NCVC-Collaborative Inflammation Cohort (JNIC) Study. Atherosclerosis, 2009, 207, 272-276.	0.4	51
123	Incidences of Herpes Zoster and Postherpetic Neuralgia in Japanese Adults Aged 50 Years and Older From a Community-based Prospective Cohort Study: The SHEZ Study. Journal of Epidemiology, 2015, 25, 617-625.	1.1	51
124	Anemia and Reduced Kidney Function as Risk Factors forÂNew Onset of Atrial Fibrillation (from the) Tj ETQq0 0	0 rgBT /O	verlock 10 Tf 5
125	Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. Journal of Nutrition, 2017, 147, 1749-1756.	1.3	51
126	Identification of dementia using standard clinical assessments by primary care physicians in Japan. Geriatrics and Gerontology International, 2018, 18, 738-744.	0.7	51

#	Article	IF	CITATIONS
127	Frequency of Food Intake and Estimated Nutrient Intake among Men and Women: The JACC Study Journal of Epidemiology, 2005, 15, S24-S42.	1.1	50
128	Serum total homocysteine concentrations and risk of mortality from stroke and coronary heart disease in Japanese: The JACC study. Atherosclerosis, 2008, 198, 412-418.	0.4	50
129	Effects of a laughter and exercise program on physiological and psychological health among communityâ€dwelling elderly in Japan: Randomized controlled trial. Geriatrics and Gerontology International, 2013, 13, 152-160.	0.7	50
130	Population profile and residential environment of an urban poor community in Dhaka, Bangladesh. Environmental Health and Preventive Medicine, 2017, 22, 1.	1.4	49
131	Effects of serum B vitamins on elevated plasma homocysteine levels associated with the mutation of methylenetetrahydrofolate reductase gene in Japanese. Atherosclerosis, 2002, 164, 321-328.	0.4	48
132	Adult height and the risk of cardiovascular disease among middle aged men and women in Japan. European Journal of Epidemiology, 2011, 26, 13-21.	2.5	48
133	Age―and Genderâ€specific BMI in Terms of the Lowest Mortality in Japanese General Population. Obesity, 2008, 16, 2348-2355.	1.5	47
134	Body Fat Distribution and the Risk of Hypertension and Diabetes among Japanese Men and Women. Hypertension Research, 2008, 31, 851-857.	1.5	47
135	Self-reported stroke and myocardial infarction had adequate sensitivity in a population-based prospective study JPHC (Japan Public Health Center)–based Prospective Study. Journal of Clinical Epidemiology, 2009, 62, 667-673.	2.4	47
136	High-density Lipoprotein Subclasses and Risk of Stroke and its Subtypes in Japanese Population. Stroke, 2013, 44, 327-333.	1.0	47
137	Association between mortality and incidence rates of coronary heart disease and stroke: The Japan Public Health Center-based prospective (JPHC) study. International Journal of Cardiology, 2016, 222, 281-286.	0.8	47
138	The Circulatory Risk in Communities Study (CIRCS): A Long-Term Epidemiological Study for Lifestyle-Related Disease Among Japanese Men and Women Living in Communities. Journal of Epidemiology, 2019, 29, 83-91.	1.1	47
139	Influenza (H1N1) 2009 Outbreak and School Closure, Osaka Prefecture, Japan. Emerging Infectious Diseases, 2009, 15, 1685-1685.	2.0	46
140	Association between Non-High-Density Lipoprotein Cholesterol Levels and the Incidence of Coronary Heart Disease among Japanese: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2011, 18, 454-463.	0.9	46
141	Prevalence of non-communicable disease risk factors among poor shantytown residents in Dhaka, Bangladesh: a community-based cross-sectional survey. BMJ Open, 2017, 7, e014710.	0.8	46
142	Coping strategies and risk of cardiovascular disease incidence and mortality: the Japan Public Health Center-based prospective Study. European Heart Journal, 2016, 37, 890-899.	1.0	45
143	Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. BMJ, The, 2020, 368, m34.	3.0	45
144	Physician–Pharmacist Cooperation Program for Blood Pressure Control in Patients With Hypertension: A Randomized-Controlled Trial. American Journal of Hypertension, 2010, 23, 1144-1152.	1.0	44

#	Article	IF	Citations
145	Carotid Artery Wall Thickness and Risk of Stroke Subtypes. Stroke, 2011, 42, 397-403.	1.0	44
146	Development of a Point-based Prediction Model for the Incidence of Total Stroke. Stroke, 2013, 44, 1295-1302.	1.0	44
147	Combined Effect of Blood Pressure and Total Cholesterol Levels on Long-Term Risks of Subtypes of Cardiovascular Death. Hypertension, 2015, 65, 517-524.	1.3	44
148	Magnesium Intake and Risk of Self-Reported Type 2 Diabetes among Japanese. Journal of the American College of Nutrition, 2010, 29, 99-106.	1.1	43
149	Relationship Between Cold Temperature and Cardiovascular Mortality, With Assessment of Effect Modification by Individual Characteristics. Circulation Journal, 2013, 77, 1854-1861.	0.7	43
150	Association of high-density lipoprotein cholesterol concentration with different types of stroke and coronary heart disease: The Japan Public Health Center-based prospective (JPHC) study. Atherosclerosis, 2017, 265, 147-154.	0.4	43
151	The Japanese food score and risk of all-cause, CVD and cancer mortality: the Japan Collaborative Cohort Study. British Journal of Nutrition, 2018, 120, 464-471.	1.2	43
152	Nutrition and disease in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). Asian Pacific Journal of Cancer Prevention, 2007, 8 Suppl, 35-80.	0.5	43
153	Cardiovascular Disease Epidemiology of Cerebrovascular Disease : Stroke Epidemic in Japan. Journal of Epidemiology, 1996, 6, 43-47.	1.1	42
154	Age-Specific Relationship between Blood Pressure and the Risk of Total and Cardiovascular Mortality in Japanese Men and Women. Hypertension Research, 2005, 28, 901-909.	1.5	42
155	Metabolic Syndrome and the Risk of Ischemic Heart Disease and Stroke among Middle-Aged Japanese. Hypertension Research, 2008, 31, 1887-1894.	1.5	42
156	Trends for coronary heart disease and its risk factors in Japan: Epidemiologic and pathologic studies Japanese Circulation Journal, 1990, 54, 428-435.	1.0	41
157	Angiotensinogen T174M and M235T variants, sodium intake and hypertension among non-drinking, lean Japanese men and women. Journal of Hypertension, 2000, 18, 1197-1206.	0.3	41
158	The trend of coronary heart disease and its risk factors based on epidemiological investigations Japanese Circulation Journal, 1987, 51, 319-324.	1.0	40
159	Type A behaviour and risk of coronary heart disease: The JPHC Study. International Journal of Epidemiology, 2008, 37, 1395-1405.	0.9	40
160	Î ³ -Glutamyltranspeptidase and Incident Stroke Among Japanese Men and Women. Stroke, 2010, 41, 385-388.	1.0	40
161	Rice consumption is not associated with risk of cardiovascular disease morbidity or mortality in Japanese men and women: a large population-based, prospective cohort study. American Journal of Clinical Nutrition, 2014, 100, 199-207.	2.2	40
162	Impact of Neighborhood Socioeconomic Conditions on the Risk of Stroke in Japan. Journal of Epidemiology, 2015, 25, 254-260.	1.1	40

#	Article	IF	Citations
163	Impact of seaweed intake on health. European Journal of Clinical Nutrition, 2021, 75, 877-889.	1.3	40
164	Associations Between Metabolic Syndrome and Mortality From Cardiovascular Disease in Japanese General Population, Findings on Overweight and Non-Overweight Individuals Ibaraki Prefectural Health Study. Circulation Journal, 2009, 73, 1635-1642.	0.7	39
165	Heavy Alcohol Consumption and Risk of Atrial Fibrillation. Circulation Journal, 2014, 78, 955-961.	0.7	39
166	Alcohol consumption patterns in Thailand and their relationship with non-communicable disease. BMC Public Health, 2015, 15, 1297.	1.2	39
167	Bowel Movement Frequency, Laxative Use, and Mortality From Coronary Heart Disease and Stroke Among Japanese Men and Women: The Japan Collaborative Cohort (JACC) Study. Journal of Epidemiology, 2016, 26, 242-248.	1.1	39
168	Impact of Alcohol Intake and Drinking Patterns on Mortality From All Causes and Major Causes of Death in a Japanese Population. Journal of Epidemiology, 2018, 28, 140-148.	1.1	39
169	Seaweed intake and risk of cardiovascular disease: the Japan Public Health Center–based Prospective (JPHC) Study. American Journal of Clinical Nutrition, 2019, 110, 1449-1455.	2.2	39
170	An inverse correlation of VZV skin-test reaction, but not antibody, with severity of herpes zoster skin symptoms and zoster-associated pain. Journal of Dermatological Science, 2013, 69, 243-249.	1.0	38
171	Milk Drinking and Mortality: Findings From the Japan Collaborative Cohort Study. Journal of Epidemiology, 2015, 25, 66-73.	1.1	38
172	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. American Journal of Clinical Nutrition, 2020, 111, 1027-1035.	2.2	38
173	Antigens of Tissue Plasminogen Activator and Plasminogen Activator Inhibitor 1: Correlates in Nonsmoking Japanese and Caucasian Men and Women. Thrombosis and Haemostasis, 1993, 70, 475-480.	1.8	38
174	Associations of Dietary Iron Intake With Mortality From Cardiovascular Disease: The JACC Study. Journal of Epidemiology, 2012, 22, 484-493.	1.1	37
175	Development of a Risk Equation for the Incidence of Coronary Artery Disease and Ischemic Stroke for Middle-Aged Japanese – Japan Public Health Center-Based Prospective Study –. Circulation Journal, 2016, 80, 1386-1395.	0.7	37
176	Non-communicable disease risk factor profile among public employees in a regional city in northern Ethiopia. Scientific Reports, 2018, 8, 9298.	1.6	37
177	Alkaline Phosphatase and Risk of Stroke Among Japanese: The Circulatory Risk in Communities Study (CIRCS). Journal of Stroke and Cerebrovascular Diseases, 2013, 22, 1046-1055.	0.7	36
178	Calcium Intake and Blood Pressure in Seven Japanese Populations. American Journal of Epidemiology, 1991, 133, 776-783.	1.6	35
179	Relationships between Sleep-Disordered Breathing and Blood Pressure and Excessive Daytime Sleepiness among Truck Drivers. Hypertension Research, 2006, 29, 605-610.	1.5	35
180	Alcohol Consumption, Social Support, and Risk of Stroke and Coronary Heart Disease Among Japanese Men: The JPHC Study. Alcoholism: Clinical and Experimental Research, 2009, 33, 1025-1032.	1.4	35

#	Article	IF	Citations
181	Smoking Cessation: Predictive Factors Among Middle-Aged Japanese. Nicotine and Tobacco Research, 2010, 12, 1050-1054.	1.4	35
182	Incidence of metabolic syndrome according to combinations of lifestyle factors among middle-aged Japanese male workers. Preventive Medicine, 2010, 51, 118-122.	1.6	35
183	C-reactive protein levels and risk of stroke and its subtype in Japanese: The Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2011, 217, 187-193.	0.4	35
184	Dietary magnesium intake and risk of incident coronary heart disease in men: A prospective cohort study. Clinical Nutrition, 2018, 37, 1602-1608.	2.3	35
185	The Relation of Body Fat Distribution and Body Mass with Haemoglobin A1c, Blood Pressure and Blood Lipids in Urban Japanese Men. International Journal of Epidemiology, 1991, 20, 88-94.	0.9	34
186	Pulse pressure is an independent risk factor for stroke among middle-aged Japanese with normal systolic blood pressure: the JPHC study. Journal of Hypertension, 2011, 29, 319-324.	0.3	34
187	A Japanese health success story: trends in cardiovascular diseases, their risk factors, and the contribution of public health and personalized approaches. EPMA Journal, 2011, 2, 49-57.	3.3	34
188	Serum coenzyme Q10 and risk of disabling dementia: The Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2014, 237, 400-403.	0.4	34
189	Polymorphisms of the Beta Fibrinogen Gene and Plasma Fibrinogen Concentration in Caucasian and Japanese Population Samples. Thrombosis and Haemostasis, 1995, 73, 106-111.	1.8	34
190	Ankle-Arm Blood Pressure Index and Cardiovascular Risk Factors in Elderly Japanese Men. Hypertension Research, 2003, 26, 377-382.	1.5	33
191	Risk factors for snoring among Japanese men and women: a community-based cross-sectional study. Sleep and Breathing, 2011, 15, 63-69.	0.9	33
192	A community-based survey of varicella-zoster virus-specific immune responses in the elderly. Journal of Clinical Virology, 2012, 55, 46-50.	1.6	33
193	The combination of Helicobacter pylori- and cytotoxin-associated gene-A seropositivity in relation to the risk of myocardial infarction in middle-aged Japanese: The Japan Public Health Center-based study. Atherosclerosis, 2013, 230, 67-72.	0.4	33
194	The impact of C-reactive protein on risk of stroke, stroke subtypes, and ischemic heart disease in middle-aged Japanese: the Japan public health center-based study. Journal of Atherosclerosis and Thrombosis, 2012, 19, 756-66.	0.9	33
195	Prospective Study on Alcohol Intake and Risk of Subarachnoid Hemorrhage Among Japanese Men and Women. Alcoholism: Clinical and Experimental Research, 2000, 24, 386-389.	1.4	32
196	The relation of anger expression with blood pressure levels and hypertension in rural and urban Japanese communities. Journal of Hypertension, 2002, 20, 21-27.	0.3	32
197	Education, Social Roles, and the Risk of Cardiovascular Disease Among Middle-Aged Japanese Women. Stroke, 2008, 39, 2886-2890.	1.0	32
198	Diabetes and the risk of coronary heart disease in the general Japanese population: The Japan Public Health Center-based prospective (JPHC) study. Atherosclerosis, 2011, 216, 187-191.	0.4	32

#	Article	IF	Citations
199	Watching Television and Risk of Mortality From Pulmonary Embolism Among Japanese Men and Women. Circulation, 2016, 134, 355-357.	1.6	32
200	Dietary Inflammatory Index Is Associated with Risk of All-Cause and Cardiovascular Disease Mortality but Not with Cancer Mortality in Middle-Aged and Older Japanese Adults. Journal of Nutrition, 2019, 149, 1451-1459.	1.3	32
201	Effects of Coffee and Tea Consumption on Glucose Metabolism: A Systematic Review and Network Meta-Analysis. Nutrients, 2019, 11, 48.	1.7	32
202	Study protocol for the PURSUIT-HFPEF study: a Prospective, Multicenter, Observational Study of Patients with Heart Failure with Preserved Ejection Fraction. BMJ Open, 2020, 10, e038294.	0.8	32
203	Relationship Between Obesity and Incident Diabetes in Middle-Aged and Older Japanese Adults: The Ibaraki Prefectural Health Study. Mayo Clinic Proceedings, 2010, 85, 36-40.	1.4	31
204	Relation of Serum ^ ^alpha;- and ^ ^gamma;-Tocopherol Levels to Cardiovascular Disease-Related Mortality Among Japanese Men and Women. Journal of Epidemiology, 2012, 22, 402-410.	1.1	31
205	Dietary patterns and breast cancer risk in a prospective Japanese study. Breast Cancer, 2017, 24, 152-160.	1.3	31
206	Changes in the Employment Status and Risk of Stroke and Stroke Types. Stroke, 2017, 48, 1176-1182.	1.0	31
207	Prevalence and correlates of carotid atherosclerosis among elderly Japanese men. Atherosclerosis, 2004, 172, 353-359.	0.4	30
208	Association of lymphocyte sub-populations with clustered features of metabolic syndrome in middle-aged Japanese men. Atherosclerosis, 2004, 173, 295-300.	0.4	30
209	Caregiver Burden for Impaired Elderly Japanese with Prevalent Stroke and Dementia under Long-Term Care Insurance System. Cerebrovascular Diseases, 2008, 25, 234-240.	0.8	30
210	Associations of Psychological Distress with Metabolic Syndrome Among Japanese Urban Residents. Journal of Atherosclerosis and Thrombosis, 2011, 18, 396-402.	0.9	30
211	Associations of Anger, Anxiety, and Depressive Symptoms With Carotid Arterial Wall Thickness. Psychosomatic Medicine, 2012, 74, 517-525.	1.3	30
212	Serum α-linolenic and other ï‰-3 fatty acids, and risk of disabling dementia: Community-based nested case–control study. Clinical Nutrition, 2017, 36, 793-797.	2.3	30
213	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level. Hypertension, 2019, 73, 52-59.	1.3	30
214	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. Journal of Epidemiology, 2020, 30, 46-54.	1.1	30
215	Non-High-Density Lipoprotein Cholesterol and Risk of Stroke Subtypes and Coronary Heart Disease: The Japan Public Health Center-Based Prospective (JPHC) Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 363-374.	0.9	30
216	Serum uric acid and risk of stroke and its types: the Circulatory Risk in Communities Study (CIRCS). Hypertension Research, 2020, 43, 313-321.	1.5	30

#	Article	lF	CITATIONS
217	Prospective Study of Major and Minor ST-T Abnormalities and Risk of Stroke Among Japanese. Stroke, 2003, 34, e250-3.	1.0	29
218	Trends in Dietary Intake of Folate, Vitamins B6, and B12 among Japanese Adults in Two Rural Communities from 1974 through 2001. Journal of Epidemiology, 2005, 15, 29-37.	1.1	29
219	Education in relation to incidence of and mortality from cancer and cardiovascular disease in Japan. European Journal of Public Health, 2008, 18, 466-472.	0.1	29
220	Association between Non-High-Density Lipoprotein Cholesterol Concentrations and Mortality from Coronary Heart Disease Among Japanese Men and Women: The Ibaraki Prefectural Health Study. Journal of Atherosclerosis and Thrombosis, 2010, 17, 30-36.	0.9	29
221	Impact of obesity on incident hypertension independent of weight gain among nonhypertensive Japanese. Journal of Hypertension, 2012, 30, 1122-1128.	0.3	29
222	Long-Term Exposure to Particulate Matter in Relation to Mortality and Incidence of Cardiovascular Disease: The JPHC Study. Journal of Atherosclerosis and Thrombosis, 2013, 20, 296-309.	0.9	29
223	Adult Height and Body Mass Index in Relation to Risk of TotalÂStroke and its Subtypes: The Circulatory Risk in Communities Study. Journal of Stroke and Cerebrovascular Diseases, 2014, 23, 667-674.	0.7	29
224	Dietary intakes of fat and total mortality among Japanese populations with a low fat intake: the Japan Collaborative Cohort (JACC) Study. Nutrition and Metabolism, 2014, 11, 12.	1.3	29
225	Age trajectories of glycaemic traits in non-diabetic South Asian and white individuals: the Whitehall II cohort study. Diabetologia, 2015, 58, 534-542.	2.9	29
226	Relationship Between Dietary Vitamin D and Deaths From Stroke and Coronary Heart Disease. Stroke, 2018, 49, 454-457.	1.0	29
227	Relationships of Differential Leukocyte and Lymphocyte Subpopulations with Carotid Atherosclerosis in Elderly Men. Journal of Clinical Immunology, 2003, 23, 469-476.	2.0	28
228	Validity of the Self-administered Food Frequency Questionnaire Used in the 5-year Follow-up Survey for the JPHC Study to Assess Folate, Vitamin B6 and B12 Intake: Comparison with Dietary Records and Blood Level. Journal of Epidemiology, 2003, 13, 98-101.	1.1	28
229	Vascular Dysfunction in Patients with Chronic Arsenosis Can Be Reversed by Reduction of Arsenic Exposure. Environmental Health Perspectives, 2005, 113, 339-341.	2.8	28
230	Associations of Sleep-Disordered Breathing with Excessive Daytime Sleepiness and Blood Pressure in Japanese Women. Hypertension Research, 2008, 31, 501-506.	1.5	28
231	Age-Period-Cohort Analysis of Mortality due to Ischemic Heart Disease in Japan, 1955 to 2000. Circulation Journal, 2008, 72, 966-972.	0.7	28
232	Death by Suicide and Other Externally Caused Injuries After Stroke in Japan (1990–2010). Psychosomatic Medicine, 2014, 76, 452-459.	1.3	28
233	Diagnosed diabetes and premature death among middle-aged Japanese: results from a large-scale population-based cohort study in Japan (JPHC study). BMJ Open, 2015, 5, e007736-e007736.	0.8	28
234	Working Hours and Risk of Acute Myocardial Infarction and Stroke Among Middle-Aged Japanese Men ― The Japan Public Health Center-Based Prospective Study Cohort II ―. Circulation Journal, 2019, 83, 1072-1079.	0.7	28

#	Article	IF	Citations
235	Serum vitamin C concentration and hs-CRP level in middle-aged Japanese men and women. Atherosclerosis, 2010, 208, 496-500.	0.4	27
236	Self-Reported Snoring Frequency and Incidence of Cardiovascular Disease: The Circulatory Risk in Communities Study (CIRCS). Journal of Epidemiology, 2012, 22, 295-301.	1.1	27
237	The Impact of C-Reactive Protein on Risk of Stroke, Stroke Subtypes, and Ischemic Heart Disease in Middle-Aged Japanese: the Japan Public Health Center-Based Study. Journal of Atherosclerosis and Thrombosis, 2012, 19, .	0.9	27
238	Hemoglobin A1c Levels and the Risk of Cardiovascular Disease in People Without Known Diabetes. Medicine (United States), 2015, 94, e785.	0.4	27
239	Disease history and risk of comorbidity in women's life course: a comprehensive analysis of the Japan Nurses' Health Study baseline survey. BMJ Open, 2015, 5, e006360-e006360.	0.8	27
240	Modifiable Risk Factors for Cardiovascular Disease in Korea and Japan. Korean Circulation Journal, 2021, 51, 643.	0.7	27
241	Tuberculosis infection among homeless persons and caregivers in a high-tuberculosis-prevalence area in Japan: a cross-sectional study. BMC Infectious Diseases, 2011, 11, 22.	1.3	26
242	Prediagnostic Plasma Antibody Levels to Periodontopathic Bacteria and Risk of Coronary Heart Disease. International Heart Journal, 2012, 53, 209-214.	0.5	26
243	Geographically-based discrimination is a social determinant of mental health in a deprived or stigmatized area in Japan: A cross-sectional study. Social Science and Medicine, 2012, 75, 1015-1021.	1.8	26
244	Salt preference and mortality from stroke and coronary heart disease for Japanese men and women: The JACC study. Preventive Medicine, 2012, 54, 32-37.	1.6	26
245	Serum Albumin and High-Sensitivity C-reactive Protein are Independent Risk Factors of Chronic Kidney Disease in Middle-Aged Japanese Individuals: the Circulatory Risk in Communities Study. Journal of Atherosclerosis and Thrombosis, 2016, 23, 1089-1098.	0.9	26
246	Impact of Hypertension and Subclinical Organ Damage on the Incidence of Cardiovascular Disease Among Japanese Residents at the Population and Individual Levels ― The Circulatory Risk in Communities Study (CIRCS) ―. Circulation Journal, 2017, 81, 1022-1028.	0.7	26
247	Serum Fatty Acid and Risk of Coronary Artery Disease ― Circulatory Risk in Communities Study (CIRCS) ―. Circulation Journal, 2018, 82, 3013-3020.	0.7	26
248	Among the water-soluble vitamins, dietary intakes of vitamins C, B ₂ and folate are associated with the reduced risk of diabetes in Japanese women but not men. British Journal of Nutrition, 2019, 121, 1357-1364.	1.2	26
249	Low BMI and weight loss aggravate COPD mortality in men, findings from a large prospective cohort: the JACC study. Scientific Reports, 2021, 11, 1531.	1.6	26
250	Metabolic syndrome and urinary cGMP excretion in general population. Atherosclerosis, 2007, 190, 423-428.	0.4	25
251	The Association between Concentrations of Green Tea and Blood Glucose Levels. Journal of Clinical Biochemistry and Nutrition, 2009, 44, 41-45.	0.6	25
252	Incidence and survival trends for childhood cancer in Osaka, Japan, 1973–2001. Cancer Science, 2010, 101, 787-792.	1.7	25

#	Article	IF	CITATIONS
253	Nocturnal Intermittent Hypoxia and Metabolic Syndrome; the Effect of being Overweight: the CIRCS Study. Journal of Atherosclerosis and Thrombosis, 2010, 17, 369-377.	0.9	25
254	Salty Food Preference and Intake and Risk of Gastric Cancer: The JACC Study. Journal of Epidemiology, 2016, 26, 92-97.	1.1	25
255	Passive smoking and mortality from aortic dissection or aneurysm. Atherosclerosis, 2017, 263, 145-150.	0.4	25
256	Frequency of Seaweed Intake and Its Association with Cardiovascular Disease Mortality: The JACC Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 1340-1347.	0.9	25
257	Alcohol consumption and risks of hypertension and cardiovascular disease in Japanese men and women. Hypertension Research, 2020, 43, 477-481.	1.5	25
258	Leukocyte count is an independent predictor for risk of acute myocardial infarction in middle-aged Japanese men. Atherosclerosis, 2007, 195, 147-152.	0.4	24
259	High Sodium Intake Strengthens the Association between Angiotensinogen T174M Polymorphism and Blood Pressure Levels among Lean Men and Women: a Community-Based Study. Hypertension Research, 2004, 27, 53-60.	1.5	23
260	Low Serum LDL Cholesterol Levels Are Associated with Elevated Mortality from Liver Cancer in Japan: the Ibaraki Prefectural Health Study. Tohoku Journal of Experimental Medicine, 2013, 229, 203-211.	0.5	23
261	Association between Chlamydophila pneumoniae infection and risk of coronary heart disease for Japanese: The JPHC study. Atherosclerosis, 2014, 233, 338-342.	0.4	23
262	Effects of a traditional herbal medicine on peripheral blood flow in women experiencing peripheral coldness: a randomized controlled trial. BMC Complementary and Alternative Medicine, 2015, 15, 105.	3.7	23
263	Body Mass Index and Risks of Incident Ischemic Stroke Subtypes: The Japan Public Health Center-Based Prospective (JPHC) Study. Journal of Epidemiology, 2019, 29, 325-333.	1.1	23
264	Trends of Cardiovascular Risk Factors and Diseases in Japan: Implications for Primordial Prevention. Preventive Medicine, 1999, 29, S102-S105.	1.6	22
265	Relationship of smoking and smoking cessation with ankle-to-arm blood pressure index in elderly Japanese men. European Journal of Cardiovascular Prevention and Rehabilitation, 2006, 13, 243-248.	3.1	22
266	Non-fasting blood glucose and risk of incident coronary heart disease in middle-aged general population: The Circulatory Risk in Communities Study (CIRCS). Preventive Medicine, 2012, 55, 603-607.	1.6	22
267	Socioeconomic Status Inconsistency and Risk of Stroke Among Japanese Middle-Aged Women. Stroke, 2014, 45, 2592-2598.	1.0	22
268	VZV skin-test reaction, but not antibody, is an important predictive factor for postherpetic neuralgia. Journal of Dermatological Science, 2015, 79, 235-240.	1.0	22
269	Educational Levels and Risk of Suicide in Japan: The Japan Public Health Center Study (JPHC) Cohort I. Journal of Epidemiology, 2016, 26, 315-321.	1.1	22
270	Comparison between the triglycerides standardization of routine methods used in Japan and the chromotropic acid reference measurement procedure used by the CDC Lipid Standardization Programme. Annals of Clinical Biochemistry, 2016, 53, 632-639.	0.8	22

#	Article	IF	Citations
271	Work–family conflict and self-rated health among Japanese workers: How household income modifies associations. PLoS ONE, 2017, 12, e0169903.	1.1	22
272	Alpha-adducin G460W polymorphism, urinary sodium excretion, and blood pressure in community-based samples. American Journal of Hypertension, 2004, 17, 385-390.	1.0	21
273	The relationships between interest for and participation in health screening and risk of mortality: The Japan Collaborative Cohort Study. Preventive Medicine, 2005, 41, 767-771.	1.6	21
274	The Shozu Herpes Zoster (SHEZ) Study: Rationale, Design, and Description of a Prospective Cohort Study. Journal of Epidemiology, 2012, 22, 167-174.	1.1	21
275	Total and high molecular weight adiponectin levels and risk of cardiovascular disease in individuals with high blood glucose levels. Atherosclerosis, 2013, 229, 222-227.	0.4	21
276	Total cholesterol performance of Abell–Levy–Brodie–Kendall reference measurement procedure: Certification of Japanese in-vitro diagnostic assay manufacturers through CDC's Cholesterol Reference Method Laboratory Network. Clinica Chimica Acta, 2015, 445, 127-132.	0.5	21
277	Similarities and differences between coronary heart disease and stroke in the associations with cardiovascular risk factors: The Japan Collaborative Cohort Study. Atherosclerosis, 2017, 261, 124-130.	0.4	21
278	Associations of Perceived Mental Stress, Sense of Purpose in Life, and Negative Life Events With the Risk of Incident Herpes Zoster and Postherpetic Neuralgia. American Journal of Epidemiology, 2018, 187, 251-259.	1.6	21
279	Cardiovascular disease, a major global burden: Epidemiology of stroke and ischemic heart disease in Japan. Global Health & Medicine, 2021, 3, 358-364.	0.6	21
280	Impact of Caregiver Type for 3-Year-Old Children on Subsequent Between-Meal Eating Habits and Being Overweight From Childhood to Adulthood: A 20-Year Follow-up of the Ibaraki Children's Cohort (IBACHIL) Study. Journal of Epidemiology, 2015, 25, 600-607.	1.1	20
281	Prospective study of seaweed consumption and thyroid cancer incidence in women. European Journal of Cancer Prevention, 2016, 25, 239-245.	0.6	20
282	Marital Transition and Risk of Stroke. Stroke, 2016, 47, 991-998.	1.0	20
283	A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan. European Journal of Epidemiology, 2017, 32, 547-557.	2.5	20
284	Sleep duration and risk of breast cancer: The JACC Study. Breast Cancer Research and Treatment, 2019, 174, 219-225.	1.1	20
285	Prenatal tobacco smoking is associated with postpartum depression in Japanese pregnant women: The japan environment and children's study. Journal of Affective Disorders, 2020, 264, 76-81.	2.0	20
286	One-year community-based education program for hypercholesterolemia in middle-aged Japanese: a long-term outcome at 8-year follow-up. Atherosclerosis, 2002, 164, 195-202.	0.4	19
287	Screening for Sleep-Disordered Breathing at Workplaces. Industrial Health, 2005, 43, 53-57.	0.4	19
288	Plasma Fibrinogen Concentrations and Risk of Stroke and Its Subtypes Among Japanese Men and Women. Stroke, 2006, 37, 2488-2492.	1.0	19

#	Article	IF	Citations
289	Stratified Age-Period-Cohort Analysis of Stroke Mortality in Japan, 1960 to 2000. Journal of Stroke and Cerebrovascular Diseases, 2007, 16, 91-102.	0.7	19
290	LDL cholesterol performance of beta quantification reference measurement procedure. Clinica Chimica Acta, 2014, 431, 288-293.	0.5	19
291	Occupational status and job stress in relation to cardiovascular stress reactivity in Japanese workers. Preventive Medicine Reports, 2016, 4, 61-67.	0.8	19
292	Clinical and immunologic features of recurrent herpes zoster (HZ). Journal of the American Academy of Dermatology, 2016, 75, 950-956.e1.	0.6	19
293	Yellow Tongue Coating is Associated With Diabetes Mellitus Among Japanese Non-smoking Men and Women: The Toon Health Study. Journal of Epidemiology, 2018, 28, 287-291.	1.1	19
294	Oxalates, urinary stones and risk of cardiovascular diseases. Medical Hypotheses, 2020, 137, 109570.	0.8	19
295	Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. European Journal of Clinical Nutrition, 2021, 75, 954-968.	1.3	19
296	Circulating miR-21, miR-29a, and miR-126 are associated with premature death risk due to cancer and cardiovascular disease: the JACC Study. Scientific Reports, 2021, 11, 5298.	1.6	19
297	Television Viewing Time and Mortality From Stroke and Coronary Artery Disease Among Japanese Men and Women – The Japan Collaborative Cohort Study –. Circulation Journal, 2015, 79, 2389-2395.	0.7	18
298	Association Between Average Daily Television Viewing Time and Chronic Obstructive Pulmonary Disease-Related Mortality: Findings From the Japan Collaborative Cohort Study. Journal of Epidemiology, 2015, 25, 431-436.	1.1	18
299	The reasonable reliability of a self-administered food frequency questionnaire for an urban, Japanese, middle-aged population: the Suita study. Nutrition Research, 2015, 35, 14-22.	1.3	18
300	Plasma levels of n-3 fatty acids and risk of coronary heart disease among Japanese: The Japan Public Health Center-based (JPHC) study. Atherosclerosis, 2018, 272, 226-232.	0.4	18
301	Cruciferous vegetable intake and mortality in middle-aged adults: A prospective cohort study. Clinical Nutrition, 2019, 38, 631-643.	2.3	18
302	Associations of Carotid Intimaâ€Media Thickness and Plaque Heterogeneity With the Risks of Stroke Subtypes and Coronary Artery Disease in the Japanese General Population: The Circulatory Risk in Communities Study. Journal of the American Heart Association, 2020, 9, e017020.	1.6	18
303	Habitual tub bathing and risks of incident coronary heart disease and stroke. Heart, 2020, 106, 732-737.	1.2	18
304	Alcohol Consumption and Risk of Gastric Cancer: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2021, 31, 30-36.	1.1	18
305	A Case-Reference Study on Plasma Fibrinogen Concentrations and Coronary Atherosclerosis in Japanese. Journal of Epidemiology, 1996, 6, 81-86.	1.1	17
306	Plasma Fibrinogen and its Correlates in Urban Japanese Men. International Journal of Epidemiology, 1996, 25, 521-527.	0.9	17

#	Article	IF	Citations
307	A Prospective Cohort Study of Employment Status and Mortality from Circulatory Disorders among Japanese Workers. Journal of Occupational Health, 2005, 47, 510-517.	1.0	17
308	A Prospective Cohort Study of Perceived Noise Exposure at Work and Cerebrovascular Diseases among Male Workers in Japan. Journal of Occupational Health, 2007, 49, 382-388.	1.0	17
309	Alcohol Consumption Before Sleep Is Associated With Severity of Sleep-Disordered Breathing Among Professional Japanese Truck Drivers. Alcoholism: Clinical and Experimental Research, 2007, 31, 2053-2058.	1.4	17
310	Association of Body Mass Index and Mortality in Japanese Diabetic Men and Women Based on Self-Reports: The Japan Collaborative Cohort (JACC) Study. Journal of Epidemiology, 2015, 25, 553-558.	1.1	17
311	Passive smoking and chronic obstructive pulmonary disease mortality: findings from the Japan collaborative cohort study. International Journal of Public Health, 2017, 62, 489-494.	1.0	17
312	Dietary intakes of fat soluble vitamins as predictors of mortality from heart failure in a large prospective cohort study. Nutrition, 2018, 47, 50-55.	1.1	17
313	Fat-soluble vitamins from diet in relation to risk of type 2 diabetes mellitus in Japanese population. British Journal of Nutrition, 2019, 121, 647-653.	1.2	17
314	Maternal total energy, macronutrient and vitamin intakes during pregnancy associated with the offspring's birth size in the Japan Environment and Children's Study. British Journal of Nutrition, 2020, 124, 558-566.	1.2	17
315	Body Mass Index and Mortality From Aortic Aneurysm and Dissection. Journal of Atherosclerosis and Thrombosis, 2021, 28, 338-348.	0.9	17
316	Smoking cessation and COPD mortality among Japanese men and women: The JACC study. Preventive Medicine, 2012, 55, 639-643.	1.6	16
317	Association between dietary behavior and risk of hypertension among Japanese male workers. Hypertension Research, 2013, 36, 374-380.	1.5	16
318	Utility of the Triglyceride Level for Predicting Incident Diabetes Mellitus According to the Fasting Status and Body Mass Index Category: The Ibaraki Prefectural Health Study. Journal of Atherosclerosis and Thrombosis, 2014, 21, 1152-1169.	0.9	16
319	Trends in the proportions of stroke subtypes and coronary heart disease in the Japanese men and women from 1995 to 2009. Atherosclerosis, 2016, 248, 219-223.	0.4	16
320	Serum creatinine levels and risk of incident type 2 diabetes mellitus or dysglycemia in middle-aged Japanese men: a retrospective cohort study. BMJ Open Diabetes Research and Care, 2018, 6, e000492.	1.2	16
321	Association of Low Job Control with a Decrease in Memory(CD4+CD45RO+) T Lymphocytes in Japanese Middle-Aged Male Workers in an Electric Power Plant Industrial Health, 2002, 40, 142-148.	0.4	16
322	Facilitators and barriers of adopting healthy lifestyle in rural China: a qualitative analysis through social capital perspectives. Nagoya Journal of Medical Science, 2016, 78, 163-73.	0.6	16
323	HEMOSTATIC VARIABLES IN JAPANESE AND CAUCASIAN MEN. American Journal of Epidemiology, 1990, 132, 41-46.	1.6	15
324	Adiposity and risk of cardiovascular diseases in Japan: secular trend, individual level associations and causal pathway $\hat{a} \in \mathbb{C}^m$ implications for the prevention of cardiovascular diseases in societies with rapid economic development. EPMA Journal, 2011, 2, 65-73.	3.3	15

#	Article	IF	CITATIONS
325	Long-term exposure to elevated blood pressure and mortality from cardiovascular disease in a Japanese population: the Ibaraki Prefectural Health Study. Hypertension Research, 2011, 34, 139-144.	1.5	15
326	Cost-effectiveness and budget impact analyses of a long-term hypertension detection and control program for stroke prevention. Journal of Hypertension, 2012, 30, 1874-1879.	0.3	15
327	Neighborhood contextual factors for smoking among middle-aged Japanese: A multilevel analysis. Health and Place, 2015, 31, 17-23.	1.5	15
328	Diabetes Trend and Impact on Risk of Cardiovascular Disease in Middle-Aged Japanese People – The CIRCS Study –. Circulation Journal, 2016, 80, 2343-2348.	0.7	15
329	No modifying effect of education level on the association between lifestyle behaviors and cardiovascular mortality: the Japan Collaborative Cohort Study. Scientific Reports, 2017, 7, 39820.	1.6	15
330	Recurrent Pregnancy Loss and Cardiovascular Disease Mortality in Japanese Women: A Population-Based, Prospective Cohort Study. Journal of Stroke and Cerebrovascular Diseases, 2017, 26, 1047-1054.	0.7	15
331	Daily Total Physical Activity and Incident Cardiovascular Disease in Japanese Men and Women. Circulation, 2017, 135, 1471-1473.	1.6	15
332	Fifty-year Time Trends in Blood Pressures, Body Mass Index and their Relations in a Japanese Community: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2017, 24, 518-529.	0.9	15
333	Adult height and all-cause and cause-specific mortality in the Japan Public Health Center-based Prospective Study (JPHC). PLoS ONE, 2018, 13, e0197164.	1.1	15
334	Measurement Performance of Reagent Manufacturers by Centers for Disease Control and Prevention/Cholesterol Reference Method Laboratory Network Lipid Standardization Specified for Metabolic Syndrome-Focused Health Checkups Program in Japan. Journal of Atherosclerosis and Thrombosis, 2009, 16, 756-763.	0.9	15
335	Impact of modifiable healthy lifestyle adoption on lifetime gain from middle to older age. Age and Ageing, 2022, 51 , .	0.7	15
336	Validity of a self-administered food frequency questionnaire (FFQ) and its generalizability to the estimation of dietary folate intake in Japan. Nutrition Journal, 2005, 4, 26.	1.5	14
337	Sleep-Disordered Breathing and Blood Pressure Levels Among Shift and Day Workers. American Journal of Hypertension, 2006, 19, 346-351.	1.0	14
338	Prospective cohort study on television viewing time and incidence of lung cancer: findings from the Japan Collaborative Cohort Study. Cancer Causes and Control, 2013, 24, 1547-1553.	0.8	14
339	Relationship between the Ankle-Brachial Index and the Risk of Coronary Heart Disease and Stroke: The Circulatory Risk in Communities Study. Journal of Atherosclerosis and Thrombosis, 2014, 21, 1283-1289.	0.9	14
340	Descriptive Epidemiology of Diabetes Prevalence and HbA1c Distributions Based on a Self-Reported Questionnaire and a Health Checkup in the JPHC Diabetes Study. Journal of Epidemiology, 2014, 24, 460-468.	1.1	14
341	Association between markers of arterial stiffness and atrial fibrillation in the Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2017, 263, 244-248.	0.4	14
342	Retinal Vascular Changes and Prospective Risk of Disabling Dementia: the Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2017, 24, 687-695.	0.9	14

#	Article	IF	Citations
343	Associations of disaster-related and psychosocial factors with changes in smoking status after a disaster: a cross-sectional survey after the Great East Japan Earthquake. BMJ Open, 2018, 8, e018943.	0.8	14
344	Variations in the estimated intake of acrylamide from food in the Japanese population. Nutrition Journal, 2020, 19, 17.	1.5	14
345	Green Tea and Coffee Consumption and All-Cause Mortality Among Persons With and Without Stroke or Myocardial Infarction. Stroke, 2021, 52, 957-965.	1.0	14
346	Weight change during middle age and risk of stroke and coronary heart disease: The Japan Public Health Center–based Prospective Study. Atherosclerosis, 2021, 322, 67-73.	0.4	14
347	Serum anti-AP3D1 antibodies are risk factors for acute ischemic stroke related with atherosclerosis. Scientific Reports, 2021, 11, 13450.	1.6	14
348	Relationship between unhealthy sleep status and dry eye symptoms in a Japanese population: The JPHC-NEXT study. Ocular Surface, 2021, 21, 306-312.	2.2	14
349	Night Work, Rotating Shift Work, and the Risk of Cancer in Japanese Men and Women: The JACC Study. Journal of Epidemiology, 2021, 31, 585-592.	1.1	14
350	Relationships of Craniofacial Morphology and Body Mass Index with Sleep-Disordered Breathing in Japanese Men. Laryngoscope, 2004, 114, 1838-1842.	1,1	13
351	History of Blood Transfusion before 1990 Is a Risk Factor for Stroke and Cardiovascular Diseases: The Japan Collaborative Cohort Study (JACC Study). Cerebrovascular Diseases, 2005, 20, 164-171.	0.8	13
352	Parental History and Lifestyle Behaviors in Relation to Mortality From Stroke Among Japanese Men and Women: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2012, 22, 331-339.	1,1	13
353	Cellular growth factors in relation to mortality from cardiovascular disease in middle-aged Japanese: The JACC study. Atherosclerosis, 2012, 224, 154-160.	0.4	13
354	Body Mass Index and Weight Change During Adulthood Are Associated With Increased Mortality From Liver Cancer: The JACC Study. Journal of Epidemiology, 2013, 23, 219-226.	1.1	13
355	Influence of Municipal- and Individual-level Socioeconomic Conditions on Mortality in Japan. International Journal of Behavioral Medicine, 2014, 21, 737-749.	0.8	13
356	Association Between Weight Change Since 20 Years of Age With Mortality From Myocardial Infarction and Chronic Heart Failure in the Japan Collaborative Cohort (JACC) Study. Circulation Journal, 2014, 78, 649-655.	0.7	13
357	HDL cholesterol performance using an ultracentrifugation reference measurement procedure and the designated comparison method. Clinica Chimica Acta, 2015, 439, 185-190.	0.5	13
358	Occupational physical activity in relation to risk of cardiovascular mortality: The Japan Collaborative Cohort Study for Evaluation for Cancer Risk (JACC Study). Preventive Medicine, 2016, 89, 286-291.	1.6	13
359	Tobacco Control Measures to Reduce Socioeconomic Inequality in Smoking: The Necessity, Time-Course Perspective, and Future Implications. Journal of Epidemiology, 2018, 28, 170-175.	1.1	13
360	Sleep Shortage Is Associated With Postherpetic Neuralgia Development Through Hyperesthesia and Acute Pain Intensity: A Communityâ€Based Prospective Cohort Study. Pain Practice, 2019, 19, 476-483.	0.9	13

#	Article	IF	CITATIONS
361	Associations of Daily Walking Time With Pneumonia Mortality Among Elderly Individuals With or Without a Medical History of Myocardial Infarction or Stroke: Findings From the Japan Collaborative Cohort Study. Journal of Epidemiology, 2019, 29, 233-237.	1.1	13
362	Effects of brief self-exercise education on the management of chronic low back pain: A community-based, randomized, parallel-group pragmatic trial. Modern Rheumatology, 2021, 31, 890-898.	0.9	13
363	Manganese intake from foods and beverages is associated with a reduced risk of type 2 diabetes. Maturitas, 2021, 143, 127-131.	1.0	13
364	Serum anti-DIDO1, anti-CPSF2, and anti-FOXJ2 antibodies as predictive risk markers for acute ischemic stroke. BMC Medicine, 2021, 19, 131.	2.3	13
365	Relationships of Soluble E-Selectin and High-Sensitivity C-Reactive Protein with Carotid Atherosclerosis in Japanese Men. Journal of Atherosclerosis and Thrombosis, 2009, 16, 339-345.	0.9	13
366	Risk Classification for Metabolic Syndrome and the Incidence of Cardiovascular Disease in Japan With Low Prevalence of Obesity: A Pooled Analysis of 10 Prospective Cohort Studies. Journal of the American Heart Association, 2021, 10, e020760.	1.6	13
367	Dietary fiber intake and risk of incident disabling dementia: the Circulatory Risk in Communities Study. Nutritional Neuroscience, 2023, 26, 148-155.	1.5	13
368	Relationship between urinary cGMP excretion and serum total cholesterol levels in a general population. Atherosclerosis, 2005, 179, 379-386.	0.4	12
369	Validity of brief food frequency questionnaire for estimation of dietary intakes of folate, vitamins B6and B12, and their associations with plasma homocysteine concentrations. International Journal of Food Sciences and Nutrition, 2010, 61, 61-67.	1.3	12
370	Profile of Non-communicable Disease Risk Factors Among Young People in Palau. Journal of Epidemiology, 2015, 25, 392-397.	1.1	12
371	Dietary Antioxidant Micronutrients and All-Cause Mortality: The Japan Collaborative Cohort Study for Evaluation of Cancer Risk. Journal of Epidemiology, 2018, 28, 388-396.	1.1	12
372	Blood pressure levels and risk of cardiovascular disease mortality among Japanese men and women. Journal of Hypertension, 2019, 37, 1366-1371.	0.3	12
373	Relationships of diabetes and hyperglycaemia with intraocular pressure in a Japanese population: the JPHC-NEXT Eye Study. Scientific Reports, 2020, 10, 5355.	1.6	12
374	Defining foreign patients into â€~visitors' and â€~residents' in Japanese medical facilities: difficulties in the collection of adequate data. Journal of Epidemiology, 2021, , .	1.1	12
375	Polymorphism of the apolipoprotein B gene and blood lipid concentrations in Japanese and Caucasian population samples. Atherosclerosis, 1996, 126, 233-241.	0.4	11
376	Ankle-arm blood pressure index as a correlate of preclinical carotid atherosclerosis in elderly Japanese men. Atherosclerosis, 2006, 184, 420-424.	0.4	11
377	C-reactive protein levels and risk of disabling dementia with and without stroke in Japanese: The Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2014, 236, 438-443.	0.4	11
378	Menopausal Status in Relation to Cardiovascular Stress Reactivity in Healthy Japanese Participants. Psychosomatic Medicine, 2014, 76, 701-708.	1.3	11

#	Article	IF	Citations
379	Saturated Fat Intake and Cardiovascular Disease in Japanese Population. Journal of Atherosclerosis and Thrombosis, 2015, 22, 435-439.	0.9	11
380	Alcohol consumption and mortality from aortic disease among Japanese men: The Japan Collaborative Cohort study. Atherosclerosis, 2017, 266, 64-68.	0.4	11
381	Water intake from foods and beverages and risk of mortality from CVD: the Japan Collaborative Cohort (JACC) Study. Public Health Nutrition, 2018, 21, 3011-3017.	1.1	11
382	Association of Vegetable, Fruit, and Okinawan Vegetable Consumption With Incident Stroke and Coronary Heart Disease. Journal of Epidemiology, 2020, 30, 37-45.	1.1	11
383	Cross-Sectional Association Between Employment Status and Self-Rated Health Among Middle-Aged Japanese Women: The Influence of Socioeconomic Conditions and Work-Life Conflict. Journal of Epidemiology, 2020, 30, 396-403.	1.1	11
384	Time course of metabolic status in pregnant women: The Japan Environment and Children's Study. Journal of Diabetes Investigation, 2020, 11, 1318-1325.	1.1	11
385	Serum High-Sensitivity C-Reactive Protein Levels and the Risk of Atrial Fibrillation in Japanese Population: the Circulatory Risk in Communities Study. Journal of Atherosclerosis and Thrombosis, 2021, 28, 194-202.	0.9	11
386	Seaweed Intake and Risk of Cardiovascular Disease: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2021, 28, 1298-1306.	0.9	11
387	Japan Trial in High-Risk Individuals to Enhance Their Referral to Physicians (J-HARP)—A Nurse-Led, Community-Based Prevention Program of Lifestyle-Related Disease. Journal of Epidemiology, 2020, 30, 194-199.	1.1	11
388	Profile of non-communicable disease risk factors among adults in the Republic of Palau: findings of a national STEPS survey. Nagoya Journal of Medical Science, 2015, 77, 609-19.	0.6	11
389	Impact of Body Mass Index on Obesity-Related Cancer and Cardiovascular Disease Mortality; The Japan Collaborative Cohort Study. Journal of Atherosclerosis and Thrombosis, 2022, 29, 1547-1562.	0.9	11
390	Employment situation and risk of death among middle-aged Japanese women. Journal of Epidemiology and Community Health, 2015, 69, 1012-1017.	2.0	10
391	Multivitamin Use and Risk of Stroke Mortality. Stroke, 2015, 46, 1167-1172.	1.0	10
392	The relationship between sodium concentrations in spot urine and blood pressure increases: a prospective study of Japanese general population: the Circulatory Risk in Communities Study (CIRCS). BMC Cardiovascular Disorders, 2016, 16, 55.	0.7	10
393	Changes in ischaemic ECG abnormalities and subsequent risk of cardiovascular disease. Heart Asia, 2017, 9, 36-43.	1.1	10
394	Relationship between cellâ€mediated immunity to Varicella–Zoster virus and aging in subjects from the communityâ€based Shozu Herpes Zoster study. Journal of Medical Virology, 2017, 89, 313-317.	2.5	10
395	Association between average daily television viewing time and the incidence of ovarian cancer: findings from the Japan Collaborative Cohort Study. Cancer Causes and Control, 2018, 29, 213-219.	0.8	10
396	"lkigai― Subjective Wellbeing, as a Modifier of the Parity-Cardiovascular Mortality Association ― The Japan Collaborative Cohort Study ―. Circulation Journal, 2018, 82, 1302-1308.	0.7	10

#	Article	IF	Citations
397	Cardiovascular disease mortality in relation to physical activity during adolescence and adulthood in Japan: Does school-based sport club participation matter?. Preventive Medicine, 2018, 113, 102-108.	1.6	10
398	Fish intake and risk of mortality due to aortic dissection and aneurysm: A pooled analysis of the Japan cohort consortium. Clinical Nutrition, 2019, 38, 1678-1683.	2.3	10
399	Anger Expression and the Risk of Cardiovascular Disease Among Urban and Rural Japanese Residents: The Circulatory Risk in Communities Study. Psychosomatic Medicine, 2020, 82, 215-223.	1.3	10
400	Weight Change and Mortality from Cardiovascular Diseases: The Japan Collaborative Cohort Study. Journal of Atherosclerosis and Thrombosis, 2021, 28, 25-33.	0.9	10
401	Estimation of 10-Year Risk of Death from Coronary Heart Disease, Stroke, and Cardiovascular Disease in a Pooled Analysis of Japanese Cohorts: EPOCH-JAPAN. Journal of Atherosclerosis and Thrombosis, 2021, 28, 816-825.	0.9	10
402	Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population. PLoS ONE, 2020, 15, e0244007.	1.1	10
403	Prevalence and predictors of prediabetes and diabetes among adults in Palau: population-based national STEPS survey. Nagoya Journal of Medical Science, 2016, 78, 475-483.	0.6	10
404	Long-term exposure to fine particle matter and all-cause mortality and cause-specific mortality in Japan: the JPHC Study. BMC Public Health, 2022, 22, 466.	1.2	10
405	Urinary cyclic GMP excretion and blood pressure levels in a general population. Atherosclerosis, 2004, 172, 161-166.	0.4	9
406	Ten-year evaluation of homogeneous low-density lipoprotein cholesterol methods developed by Japanese manufacturers. Journal of Atherosclerosis and Thrombosis, 2010, 17, 1275-1281.	0.9	9
407	Associations between alcohol consumption and sleep-disordered breathing among Japanese women. Respiratory Medicine, 2011, 105, 796-800.	1.3	9
408	A prospective cohort study of neighborhood stress and ischemic heart disease in Japan: a multilevel analysis using the JACC study data. BMC Public Health, 2011, 11, 398.	1.2	9
409	An association between central aortic pressure and subclinical organ damage of the heart among a general Japanese cohort: Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2014, 232, 94-98.	0.4	9
410	Cohort Profile: The Fangshan Cohort Study of Cardiovascular Epidemiology in Beijing, China. Journal of Epidemiology, 2014, 24, 84-93.	1.1	9
411	Dehydroepiandrosterone-sulfate is associated with cardiovascular reactivity to stress in women. Psychoneuroendocrinology, 2016, 69, 116-122.	1.3	9
412	Longâ€Term Prognosis of Brugadaâ€Type ECG and ECG With Atypical STâ€Segment Elevation in the Right Precordial Leads Over 20ÂYears: Results From the Circulatory Risk in Communities Study (CIRCS). Journal of the American Heart Association, 2016, 5, .	1.6	9
413	High Myopia and Its Associated Factors in JPHC-NEXT Eye Study: A Cross-Sectional Observational Study. Journal of Clinical Medicine, 2019, 8, 1788.	1.0	9
414	Impact of intention and feeling toward being pregnant on postpartum depression: the Japan Environment and Children's Study (JECS). Archives of Women's Mental Health, 2020, 23, 131-137.	1.2	9

#	Article	IF	CITATIONS
415	The Association Between Habitual Sleep Duration and Mortality According to Sex and Age: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2021, 31, 109-118.	1.1	9
416	Intake of fish and long-chain n-3 polyunsaturated fatty acids and risk of diseases in a Japanese population: a narrative review. European Journal of Clinical Nutrition, 2021, 75, 902-920.	1.3	9
417	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. Journal of Cancer Survivorship, 2021, 15, 912-921.	1.5	9
418	Association between dietary inflammatory index and serum C-reactive protein concentrations in the Japan Collaborative Cohort Study. Nagoya Journal of Medical Science, 2020, 82, 237-249.	0.6	9
419	Association between Dietary Manganese Intake and Mortality from Cardiovascular Disease in Japanese Population: The Japan Collaborative Cohort Study. Journal of Atherosclerosis and Thrombosis, 2022, 29, 1432-1447.	0.9	9
420	Use of aspirin for prevention of cardiovascular disease—1981–1982 to 1985–1986: The Minnesota Heart Survey. American Heart Journal, 1988, 116, 827-830.	1.2	8
421	Lipoprotein(a) and its correlates in Japanese and U.S. population samples. Annals of Epidemiology, 1996, 6, 324-330.	0.9	8
422	Determinants of platelet aggregation in 50–70-year-old men from three Japanese communities. Atherosclerosis, 2002, 165, 327-334.	0.4	8
423	High Sodium Intake Strengthens the Association of ACE I/D Polymorphism with Blood Pressure in a Community. American Journal of Hypertension, 2007, 20, 751-757.	1.0	8
424	Projecting the probability of survival free from cancer and cardiovascular incidence through lifestyle modification in Japan. Preventive Medicine, 2009, 48, 128-133.	1.6	8
425	Prediction of Myocardial Infarction Using Coronary Risk Scores Among Japanese Male Workers: 3M Study. Journal of Atherosclerosis and Thrombosis, 2010, 17, 452-459.	0.9	8
426	Promoting predictive, preventive and personalized medicine in treatment of cardiovascular diseases. EPMA Journal, 2011, 2, 1-4.	3.3	8
427	Nonfasting Glucose and Incident Stroke and Its Types ― The Circulatory Risk in Communities Study (CIRCS) ―. Circulation Journal, 2018, 82, 1598-1604.	0.7	8
428	Association of cigarette smoking with radial augmentation index: the Circulatory Risk in Communities Study (CIRCS). Hypertension Research, 2018, 41, 1054-1062.	1.5	8
429	Television viewing time, walking time, and risk of type 2 diabetes in Japanese men and women: The Japan Collaborative Cohort Study. Preventive Medicine, 2019, 118, 220-225.	1.6	8
430	Factors Associated With Family Member's Spanking of 3.5-year-old Children in Japan. Journal of Epidemiology, 2020, 30, 464-473.	1.1	8
431	Relationships Between Reproductive History and Mortality From Cardiovascular Diseases Among Japanese Women: The Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. Journal of Epidemiology, 2020, 30, 509-515.	1.1	8
432	Diabetes and Mortality From Respiratory Diseases: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2020, 30, 457-463.	1.1	8

#	Article	IF	CITATIONS
433	Association between mode of delivery and postpartum depression: The Japan Environment and Children's Study (JECS): a prospective cohort study. Journal of Epidemiology, 2021, , .	1.1	8
434	Relationship between Endothelial Dysfunction and Prevalence of Chronic Kidney Disease: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2021, 28, 622-629.	0.9	8
435	Relation Between Body Mass Index and Dry Eye Disease: The Japan Public Health Center–Based Prospective Study for the Next Generation. Eye and Contact Lens, 2021, 47, 449-455.	0.8	8
436	Association of Maternal Total Cholesterol With SGA or LGA Birth at Term: the Japan Environment and Children's Study. Journal of Clinical Endocrinology and Metabolism, 2022, 107, e118-e129.	1.8	8
437	Association of dietary intakes of vitamin B12, vitamin B6, folate, and methionine with the risk of esophageal cancer: the Japan Public Health Center-based (JPHC) prospective study. BMC Cancer, 2021, 21, 982.	1.1	8
438	Television Viewing Time and Breast Cancer Incidence for Japanese Premenopausal and Postmenopausal Women: The JACC Study. Cancer Research and Treatment, 2019, 51, 1509-1517.	1.3	8
439	Relationship of Urinary cGMP Excretion with Aging and Menopausal Status in a General Population. Journal of Atherosclerosis and Thrombosis, 2009, 16, 457-462.	0.9	8
440	Secondhand smoke and the risk of incident cardiovascular disease among never-smoking women. Preventive Medicine, 2022, 162, 107145.	1.6	8
441	Is the association between healthy lifestyle behaviors and cardiovascular mortality modified by overweight status? The Japan Collaborative Cohort Study. Preventive Medicine, 2014, 62, 142-147.	1.6	7
442	Association of measles and mumps with cardiovascular disease: TheÂJapan Collaborative Cohort (JACC) study. Atherosclerosis, 2015, 241, 682-686.	0.4	7
443	Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. Journal of Epidemiology, 2017, 27, 331-337.	1.1	7
444	Fish Intake and Death From Pulmonary Embolisms Among Japanese Men and Women ― The Japan Collaborative Cohort (JACC) Study ―. Circulation Journal, 2018, 82, 2063-2070.	0.7	7
445	Dietary Intake of Energy and Nutrients from Breakfast and Risk of Stroke in The Japanese Population: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2019, 26, 145-153.	0.9	7
446	Green tea consumption and risk of hematologic neoplasms: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC Study). Cancer Causes and Control, 2019, 30, 1223-1230.	0.8	7
447	Effects of Low-Dose Therapist-Led Self-Exercise Education on the Management of Chronic Low Back Pain: Protocol for a Community-Based, Randomized, 6-Month Parallel-Group Study. Spine Surgery and Related Research, 2019, 3, 377-384.	0.4	7
448	Impact of hypertension stratified by diabetes on the lifetime risk of cardiovascular disease mortality in Japan: a pooled analysis of data from the Evidence for Cardiovascular Prevention from Observational Cohorts in Japan study. Hypertension Research, 2020, 43, 1437-1444.	1.5	7
449	Tobacco Price Increase and Successful Smoking Cessation for Two or More Years in Japan. Nicotine and Tobacco Research, 2021, 23, 716-723.	1.4	7
450	Myopia, corneal endothelial cell density and morphology in a Japanese population-based cross-sectional study: the JPHC-NEXT Eye Study. Scientific Reports, 2021, 11, 6366.	1.6	7

#	Article	IF	Citations
451	Smoking cessation, weight gain and risk of cardiovascular disease. Heart, 2022, 108, 375-381.	1.2	7
452	Associations of Body Mass Index, Weight Change, Physical Activity, and Sedentary Behavior With Endometrial Cancer Risk Among Japanese Women: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2021, 31, 621-627.	1.1	7
453	Dairy intake and the risk of pancreatic cancer: the Japan Collaborative Cohort Study (JACC Study) and meta-analysis of prospective cohort studies. British Journal of Nutrition, 2022, 128, 1147-1155.	1.2	7
454	Relations of mold, stove, and fragrance products on childhood wheezing and asthma: A prospective cohort study from the Japan Environment and Children's Study. Indoor Air, 2022, 32, .	2.0	7
455	Work and Family Conflicts, Depression, and "lkigaiâ€. A Mediation Analysis in a Cross-cultural Study Between Japanese and Egyptian Civil Workers. Journal of Epidemiology, 2023, 33, 360-366.	1.1	7
456	Effectiveness of the Combined Approach for Assessing Social Gradients in Stroke Risk Among Married Women in Japan. Journal of Epidemiology, 2012, 22, 324-330.	1.1	6
457	Self-Reported Eczema in Relation with Mortality from Cardiovascular Disease in Japanese: the Japan Collaborative Cohort Study. Journal of Atherosclerosis and Thrombosis, 2019, 26, 775-782.	0.9	6
458	Association of job category and occupational activity with breast cancer incidence in Japanese female workers: the JACC study. BMC Public Health, 2020, 20, 1106.	1.2	6
459	Milk Intake and Stroke Mortality in the Japan Collaborative Cohort Study—A Bayesian Survival Analysis. Nutrients, 2020, 12, 2743.	1.7	6
460	Longitudinal Trends in Blood Pressure Associated With the Frequency of Laughter: The Circulatory Risk in Communities Study (CIRCS), a Longitudinal Study of the Japanese General Population. Journal of Epidemiology, 2021, 31, 125-131.	1.1	6
461	Association Between Birth Weight and Risk of Pregnancy-Induced Hypertension and Gestational Diabetes in Japanese Women: JPHC-NEXT Study. Journal of Epidemiology, 2022, 32, 168-173.	1.1	6
462	Impact of Major Cardiovascular Risk Factors on the Incidence of Cardiovascular Disease among Overweight and Non-Overweight Individuals: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2022, 29, 422-437.	0.9	6
463	Engaging in musical activities and the risk of dementia in older adults: A longitudinal study from the Japan gerontological evaluation study. Geriatrics and Gerontology International, 2021, 21, 451-457.	0.7	6
464	Television Viewing Time and the Risk of Colorectal Cancer Mortality among Japanese Population: The JACC Study. Cancer Research and Treatment, 2021, 53, 497-505.	1.3	6
465	Combined patterns of participation in cervical, breast, and colorectal cancer screenings and factors for non-participation in each screening among women in Japan. Preventive Medicine, 2021, 150, 106627.	1.6	6
466	Chlamydia pneumoniae Infection was Associated with Risk of Mortality from Coronary Heart Disease in Japanese Women but not Men: the JACC Study. Journal of Atherosclerosis and Thrombosis, 2010, 17, 510-516.	0.9	6
467	Risk Stratification Based on Metabolic Syndrome as well as Non- Metabolic Risk Factors in the Assessment of Carotid Atherosclerosis. Journal of Atherosclerosis and Thrombosis, 2011, 18, 504-512.	0.9	6
468	Food frequency questionnaire reproducibility for middle-aged and elderly Japanese. Asia Pacific Journal of Clinical Nutrition, 2019, 28, 362-370.	0.3	6

#	Article	IF	CITATIONS
469	Secondhand Smoke Exposure During Childhood and Cancer Mortality in Adulthood Among Never Smokers. American Journal of Epidemiology, 2021, , .	1.6	6
470	Validity of the Food Frequency Questionnaireâ€"Estimated Intakes of Sodium, Potassium, and Sodium-to-Potassium Ratio for Screening at a Point of Absolute Intake among Middle-Aged and Older Japanese Adults. Nutrients, 2022, 14, 2594.	1.7	6
471	Trends of Blood Pressure Distributions in a Northeast Rural Japanese Population. Journal of Epidemiology, 1993, 3, 63-70.	1.1	5
472	Trends for Cardiovascular Risk Factors and Diseases in Japan. Journal of Epidemiology, 1996, 6, 183-188.	1.1	5
473	Possible effects of diets on serum lipids, fatty acids and blood pressure levels in male and female Japanese university students. Environmental Health and Preventive Medicine, 2005, 10, 42-47.	1.4	5
474	Aldosterone Synthase Gene T-344C Polymorphism, Sodium and Blood Pressure in a Free-Living Population: A Community-Based Study. Hypertension Research, 2007, 30, 497-502.	1.5	5
475	Alcohol consumption, alcohol-induced flushing and incidence of acute myocardial infarction among middle-aged men in Japan—Japan Public Health Center-based prospective study. Atherosclerosis, 2007, 194, 512-516.	0.4	5
476	Reproducibility of 24-hour dietary recall for vitamin intakes by middle-aged Japanese men and women. Journal of Nutrition, Health and Aging, 2010, 14, 196-200.	1.5	5
477	Social Epidemiology and Eastern Wisdom. Journal of Epidemiology, 2012, 22, 291-294.	1.1	5
478	Duration of cigarette smoking is a risk factor for oropharyngeal cancer mortality among Japanese men and women: the Ibaraki Prefectural Health Study (IPHS). Annals of Epidemiology, 2013, 23, 546-550.	0.9	5
479	Health Disparities in Ischaemic Heart Disease Mortality in Hungary From 1970 to 2010: An Age-Period-Cohort Analysis. Journal of Epidemiology, 2015, 25, 399-406.	1.1	5
480	Blood soluble Fas levels and mortality from cardiovascular disease in middle-aged Japanese: The JACC study. Atherosclerosis, 2017, 260, 97-101.	0.4	5
481	Validity and reliability of a self-administered food frequency questionnaire for the JPHC study: The assessment of amino acid intake. Journal of Epidemiology, 2017, 27, 242-247.	1.1	5
482	Evaluating the Effect on Mortality of a No-Tranexamic acid (TXA) Policy for Cardiovascular Surgery. Journal of Cardiothoracic and Vascular Anesthesia, 2018, 32, 1627-1634.	0.6	5
483	Joint impact of muscle mass and waist circumference on type 2 diabetes in Japanese middleâ€øged adults: The Circulatory Risk in Communities Study (<scp>CIRCS</scp>). Journal of Diabetes, 2020, 12, 677-685.	0.8	5
484	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. Preventive Medicine, 2021, 148, 106561.	1.6	5
485	Peanut Consumption and Risk of Stroke and Ischemic Heart Disease in Japanese Men and Women: The JPHC Study. Stroke, 2021, 52, 3543-3550.	1.0	5
486	Job stress factors measured by Brief Job Stress Questionnaire and sickness absence among Japanese workers: A longitudinal study. Fukushima Journal of Medical Sciences, 2020, 66, 88-96.	0.1	5

#	Article	IF	Citations
487	Hobby Engagement and Risk of Disabling Dementia. Journal of Epidemiology, 2023, 33, 456-463.	1.1	5
488	Pathologic Characteristics of Stroke and Myocardial Infarction in Japan. Journal of Epidemiology, 1992, 2, 137-147.	1.1	4
489	Dietary Reference Intakes for Japanese 2010: Fat. Journal of Nutritional Science and Vitaminology, 2012, 59, S44-S52.	0.2	4
490	Relationship between sleep duration and cause-specific mortality in diabetic men and women based on self-reports. Sleep and Biological Rhythms, 2015, 13, 85-93.	0.5	4
491	Impact and attribute of each obesity-related cardiovascular risk factor in combination with abdominal obesity on total health expenditures in adult Japanese National Health insurance beneficiaries: The Ibaraki Prefectural health study. Journal of Epidemiology, 2017, 27, 354-359.	1.1	4
492	Urinary 8-Hydroxy-2'-Deoxyguanosine Levels and Cardiovascular Disease Incidence in Japan. Journal of Atherosclerosis and Thrombosis, 2020, 27, 1086-1096.	0.9	4
493	Social support, social cohesion and pain during pregnancy: The Japan Environment and Children's Study. European Journal of Pain, 2021, 25, 872-885.	1.4	4
494	Determinants of alcohol consumption and marijuana use among young adults in the Republic of Palau. Environmental Health and Preventive Medicine, 2021, 26, 12.	1.4	4
495	Risk of Herpes Zoster in Relation to Body Mass Index Among Residents Aged ≥50 Years: The Shozu Herpes Zoster Study. Journal of Epidemiology, 2022, 32, 370-375.	1.1	4
496	Association between maternal employment status during pregnancy and risk of depressive symptomatology 1 month after childbirth: the Japan Environment and Children's Study. Journal of Epidemiology and Community Health, 2021, 75, 531-539.	2.0	4
497	Leukocyte Count and Risks of Stroke and Coronary Heart Disease: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2022, 29, 527-535.	0.9	4
498	Association of serum levels of antibodies against ALDOA and FH4 with transient ischemic attack and cerebral infarction. BMC Neurology, 2021, 21, 274.	0.8	4
499	Right bundle branch block and risk of cardiovascular mortality: the Ibaraki Prefectural Health Study. Heart and Vessels, 2021, , 1.	0.5	4
500	Body Mass Index and Serum γâ€glutamyltransferase Level as Risk Factors for Injuries Related to Professional Horse Racing: A Prospective Study. Journal of Occupational Health, 2009, 51, 323-331.	1.0	4
501	Overweight and Hypertension in Relation to Chronic Musculoskeletal Pain Among Community-Dwelling Adults: The Circulatory Risk in Communities Study (CIRCS). Journal of Epidemiology, 2020, 31, 566-572.	1.1	4
502	Prediction of Lifetime Risk of Cardiovascular Disease Deaths Stratified by Sex in the Japanese Population. Journal of the American Heart Association, 2021, 10, e021753.	1.6	4
503	Soy Intake and Risk of Type 2 Diabetes Among Japanese Men and Women: JACC Study. Frontiers in Nutrition, 2021, 8, 813742.	1.6	4
504	Retinal microvascular abnormalities and risks of incident stroke and its subtypes: The Circulatory Risk in Communities Study. Journal of Hypertension, 2022, 40, 732-740.	0.3	4

#	Article	IF	CITATIONS
505	Trends in the Incidence of Cardiovascular Diseases and their Risk Factors in a Rural Japanese Population. Journal of Epidemiology, 1992, 2, 123-135.	1.1	3
506	Differences in dietary habits, serum fatty acid compositions and other coronary risk characteristics between freshmen and fourth-year male university students. Environmental Health and Preventive Medicine, 2001, 6, 143-148.	1.4	3
507	Prehospital score for acute disease: a community-based observational study in Japan. BMC Emergency Medicine, 2007, 7, 17.	0.7	3
508	Interest in health screening as a predictor of long-term overall mortality: Multilevel analysis of a Japanese national cohort study. Preventive Medicine, 2011, 52, 78-83.	1.6	3
509	Message from the New Editor-in-Chief. Journal of Epidemiology, 2011, 21, 1.	1.1	3
510	Descriptive Epidemiology of Hypertension and Its Association With Obesity: Based on the WHO STEPwise Approach to Surveillance in Palau. Asia-Pacific Journal of Public Health, 2017, 29, 278-287.	0.4	3
511	Changes in the living arrangement and risk of stroke in Japan; does it matter who lives in the household? Who among the family matters?. PLoS ONE, 2017, 12, e0173860.	1.1	3
512	Prevalence and Correlates of Dyslipidemia Among Men and Women in Palau: Findings of the Palau STEPS Survey 2011–2013. Journal of Epidemiology, 2019, 29, 97-103.	1.1	3
513	Cardiovascular reactivity to acute stress associated with sickness absence among Japanese men and women: A prospective study. Brain and Behavior, 2020, 10, e01541.	1.0	3
514	Relationship between nerve fiber layer defect and the presence of epiretinal membrane in a Japanese population: The JPHC-NEXT Eye Study. Scientific Reports, 2020, 10, 779.	1.6	3
515	Moderate Levels of N-Terminal Pro-B-Type Natriuretic Peptide is Associated with Increased Risks of Total and Ischemic Strokes among Japanese: The Circulatory Risk in Communities Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 751-760.	0.9	3
516	Risk of postpartum depression and very early child mistreatment among mothers reporting higher autistic traits: Evidence from the Japan Environment and Children's Study. Journal of Affective Disorders, 2021, 280, 11-16.	2.0	3
517	Dietary intake of tocopherols and risk of incident disabling dementia. Scientific Reports, 2021, 11, 16429.	1.6	3
518	Associations Between Occupational Status, Support at Work, and Salivary Cortisol Levels. International Journal of Behavioral Medicine, 2022, 29, 299-307.	0.8	3
519	Long-term Response of <i>Helicobacter pylori</i> Antibody Titer After Eradication Treatment in Middle-aged Japanese: JPHC-NEXT Study. Journal of Epidemiology, 2023, 33, 1-7.	1.1	3
520	Effects of Intake of Fish Rich in n-3 Polyunsaturated Fatty Acid on Serum Lipids, Serum Fatty Acid Component and Hemostatic Factors Nihon EiyŕShokuryŕGakkai Shi = Nippon EiyŕShokuryŕGakkaishi = Journal of Japanese Society of Nutrition and Food Science, 2000, 53, 1-9.	0.2	3
521	Breastfeeding in Infancy in Relation to Subsequent Physical Size: A 20-year Follow-up of the Ibaraki Children's Cohort Study (IBACHIL). Journal of Epidemiology, 2023, 33, 63-67.	1.1	3
522	Association between Serum Uric Acid and Impaired Endothelial Function: The Circulatory Risk in Communities Study. Journal of Atherosclerosis and Thrombosis, 2022, 29, 1534-1546.	0.9	3

#	Article	IF	CITATIONS
523	Association of B Vitamins and Methionine Intake with the Risk of Gastric Cancer: The Japan Public Health Center–based Prospective Study. Cancer Prevention Research, 2022, 15, 101-110.	0.7	3
524	Factors associated with prolonged duration of viral clearance in non-severe SARS-CoV-2 patients in Osaka, Japan. Environmental Health and Preventive Medicine, 2021, 26, 115.	1.4	3
525	Occasions for laughter and dementia risk: Findings from a sixâ€year cohort study. Geriatrics and Gerontology International, 2022, 22, 392-398.	0.7	3
526	Effects of a laughter program on body weight and mental health among Japanese people with metabolic syndrome risk factors: a randomized controlled trial. BMC Geriatrics, 2022, 22, 361.	1.1	3
527	Smoking Cessation and Mortality from Aortic Dissection and Aneurysm: Findings from the Japan Collaborative Cohort (JACC) Study. Journal of Atherosclerosis and Thrombosis, 2023, 30, 348-363.	0.9	3
528	Validity of claims data for identifying cancer incidence in the Japan Public Health Centerâ€based Prospective Study for the Next Generation (<scp>JPHCâ€NEXT</scp>). Pharmacoepidemiology and Drug Safety, 0, , .	0.9	3
529	The reproducibility of 24-h dietary recall for estimating mineral intakes and their food sources among middle-aged Japanese men and women. International Journal of Food Sciences and Nutrition, 2009, 60, 30-40.	1.3	2
530	Reply to Dr. Rouhani et al Journal of Nutrition, 2011, 141, 1919.	1.3	2
531	Overview of the Role of Antioxidant Vitamins as Protection Against Cardiovascular Disease. , 2014, , 213-224.		2
532	Dietary Patterns and Risk of Esophageal Cancer Mortality: The Japan Collaborative Cohort Study. Nutrition and Cancer, 2016, 68, 1001-1009.	0.9	2
533	The Prospective Association Between Plasma Concentrations of Cellular Growth Factors and Risk of Heart Failure Mortality in Japanese Population. Journal of Epidemiology, 2019, 29, 104-109.	1.1	2
534	Association of tea consumption and the risk of gastric cancer in Japanese adults: the Japan Collaborative Cohort Study. BMJ Open, 2020, 10, e038243.	0.8	2
535	A method for measuring glycerol-blanked triglyceride concentrations by using gas chromatography-isotope dilution mass spectrometry. Annals of Clinical Biochemistry, 2020, 57, 253-261.	0.8	2
536	The association of conventionally medicated systolic and diastolic blood pressure level and mortality from cardiovascular disease: is the lower the better in high stroke population?. Clinical Research in Cardiology, 2020, 109, 944-948.	1.5	2
537	Salt taste perception and blood pressure levels in population-based samples: the Circulatory Risk in Communities Study (CIRCS). British Journal of Nutrition, 2021, 125, 203-211.	1.2	2
538	Impact of Perceived Social Support on the Association Between Anger Expression and the Risk of Stroke: The Circulatory Risk in Communities Study (CIRCS). Journal of Epidemiology, 2021, , .	1.1	2
539	Risk of stroke in cancer survivors using a propensity score-matched cohort analysis. Scientific Reports, 2021, 11, 5599.	1.6	2
540	Effects of Local Cooperation between Healthcare Facilities for Infection Control after Introduction of Additional Charge for Infection Control. Japanese Journal of Environmental Infections, 2016, 31, 24-31.	0.1	2

#	Article	IF	Citations
541	Green Tea, Coffee, and Diabetes. Annals of Internal Medicine, 2006, 145, 634.	2.0	2
542	Usefulness of Skinfold Thickness Measurements for Determining Body Fat Distribution and Disease Risk for Japanese Men and Women., 2012,, 2667-2678.		2
543	Urinary Stones and Risk of Coronary Heart Disease and Stroke: the Japan Public Health Center-Based Prospective Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 1208-1215.	0.9	2
544	Can Blood Pressure in the Elderly be Reduced? Findings From a Long-term Population Survey in Japan. The American Journal of Geriatric Cardiology, 1994, 3, 42-50.	0.7	2
545	Long-term weight loss as a predictor of mortality in haemodialysis patients. Journal of Epidemiology, 2022, , .	1.1	2
546	The association between midlife living arrangement and psychiatrist-diagnosed depression in later life: who among your family members reduces the risk of depression?. Translational Psychiatry, 2022, 12, 156.	2.4	2
547	Relationship between Health Counselor Characteristics and Counseling Impact on Individuals at High-Risk for Lifestyle-Related Disease: Sub-Analysis of the J-HARP Cluster-Randomized Controlled Trial. International Journal of Environmental Research and Public Health, 2022, 19, 6375.	1.2	2
548	Letter by Noda and Iso Regarding Article, "Low-Density Lipoprotein Cholesterol Concentrations and Death Due to Intraparenchymal Hemorrhage: The Ibaraki Prefectural Health Study― Circulation, 2009, 120, e280; author reply e281.	1.6	1
549	Response by Shirakawa and Iso to Letter Regarding Article, "Watching Television and Risk of Mortality From Pulmonary Embolism Among Japanese Men and Women: The JACC Study (Japan Collaborative) Tj ETQq1 1	0.71864314	∤rgBT/Over
550	Non-Fatal Injury in Thailand From 2005 to 2013: Incidence Trends and Links to Alcohol Consumption Patterns in the Thai Cohort Study. Journal of Epidemiology, 2016, 26, 471-480.	1.1	1
551	New Policy of the Journal of Epidemiology Regarding the Relationship With the Tobacco Industry. Journal of Epidemiology, 2018, 28, 1-2.	1.1	1
552	Successful Reduction of Blood Pressure and Stroke Risk in Japan. , 2019, , 47-55.		1
553	Glycated haemoglobin (HbA1c) as a reliable option for detecting diabetes among the urban poor population in Bangladesh. European Journal of Public Health, 2020, 30, 839-841.	0.1	1
554	Diabetes Mellitus Modifies the Association of Serum Triglycerides with Ischemic Cardiovascular Disease Mortality: The Ibaraki Prefectural Health Study (IPHS). Journal of Atherosclerosis and Thrombosis, 2021, , .	0.9	1
555	Association Between Interpregnancy Interval and Risk of Preterm Birth and Its Modification by Folate Intake: The Japan Environment and Children's Study. Journal of Epidemiology, 2023, 33, 113-119.	1.1	1
556	Validity of dietary isothiocyanate intake estimates from a food frequency questionnaire using 24 h urinary isothiocyanate excretion as an objective biomarker: the JPHC-NEXT protocol area. European Journal of Clinical Nutrition, 2021, , .	1.3	1
557	The apparent inverse association between dietary carotene intake and risk of cardiovascular mortality disappeared after adjustment for other cardioprotective dietary intakes: The Japan collaborative cohort study. Nutrition, Metabolism and Cardiovascular Diseases, 2021, 31, 3064-3075.	1.1	1
558	Having hobbies and the risk of cardiovascular disease incidence: A Japan public health center-based study. Atherosclerosis, 2021, 335, 1-7.	0.4	1

#	Article	IF	CITATIONS
559	Depressive Tendency and the Risk of Death from Pneumonia: The JACC Study. Internal Medicine, 2020, 59, 3123-3130.	0.3	1
560	Retrospective survey for clinical course and aggravating factors of adolescent atopic dermatitis in two years' cohort study on firstâ€year university students. Journal of Cutaneous Immunology and Allergy, 2022, 5, 47-55.	0.2	1
561	Walking time, sports activity, job type, and body posture during work in relation to incident colorectal cancer: the JACC prospective cohort study. Cancer Causes and Control, 2022, 33, 473-481.	0.8	1
562	Brief self-exercise education for adults with chronic knee pain: A randomized controlled trial. Modern Rheumatology, 2023, 33, 408-415.	0.9	1
563	Dairy intake and the risk of esophageal cancer: the JACC Study. Journal of Epidemiology, 2022, , .	1.1	1
564	Associations of dietary intakes of vitamins $B < sub > 1 < sub > and B < sub > 3 < sub > with risk of mortality from CVD among Japanese men and women: the Japan Collaborative Cohort study. British Journal of Nutrition, 2023, 129, 1213-1220.$	1.2	1
565	The Associations between Evacuation Status and Lifestyle-Related Diseases in Fukushima after the Great East Japan Earthquake: The Fukushima Health Management Survey. International Journal of Environmental Research and Public Health, 2022, 19, 5661.	1.2	1
566	Nonfasting serum triglycerides and risk of ischemic heart disease among Japanese men and women. International Congress Series, 2004, 1262, 200-202.	0.2	0
567	Associations between Cardiovascular Risk Factors and Carotid Atherosclerosis in Middle-aged Japanese Men with Multiple Risk Factors. Industrial Health, 2008, 46, 607-612.	0.4	0
568	Message from the New Editor-in-Chief. Journal of Epidemiology, 2012, 22, 1.	1.1	0
569	Impact of specialization in gynecology and obstetrics departments on pregnant women's choice of maternity institutions. Health Economics Review, 2013, 3, 31.	0.8	0
570	Response to Letter Regarding Article, "Socioeconomic Status Inconsistency and Risk of Stroke Among Japanese Middle-Aged Women― Stroke, 2014, 45, e307.	1.0	0
571	Alcohol and Sleep-Disordered Breathing. , 2015, , 349-352.		0
572	Response to eLetter. Heart, 2020, 106, 1365.2-1366.	1.2	0
573	Reader response: Low-density lipoprotein cholesterol and risk of intracerebral hemorrhage: A prospective study. Neurology, 2020, 95, 98.2-99.	1.5	0
574	Body mass index and mortality among middle-aged Japanese individuals with diagnosed diabetes: The Japan Public Health Center-based prospective study (JPHC study). Diabetes Research and Clinical Practice, 2020, 164, 108198.	1.1	0
575	Physical Activity and Risk of Mortality from Heart Failure among Japanese Population. Journal of Atherosclerosis and Thrombosis, 2021, , .	0.9	0
576	Non-alcoholic beverages intake and risk of cardiovascular disease among Japanese men and women: the JPHC study. British Journal of Nutrition, 2021, , 1-20.	1.2	0

#	Article	IF	CITATIONS
577	990Seaweed intake and risk of cardiovascular disease: the Japan Public Health Center–based Prospective (JPHC) Study. International Journal of Epidemiology, 2021, 50, .	0.9	О
578	Supper Timing and Cardiovascular Mortality: The Japan Collaborative Cohort Study. Nutrients, 2021, 13, 3389.	1.7	0
579	Unfinished business: lessons for future G20 meetings on a more inclusive understanding of universal health coverage. GHM Open, 2021, , .	0.1	0
580	Title is missing!. Japanese Journal of Electrocardiology, 2006, 26, 31-38.	0.0	0
581	The Prevalence and over 20 Years Long Term Prognosis of the Brugada-Type Electrocardiogram: From the Circulatory Risk in Communities Study (CIRCS). Journal of Arrhythmia, 2011, 27, OP67_3.	0.5	0
582	CHAPTER 42. Function and Effects of Pyridoxine (Vitamin B6): An Epidemiological Review of Evidence. Food and Nutritional Components in Focus, 2012, , 699-715.	0.1	0
583	Pedometer Measured Walking Habits and Serum HDL-cholesterol Level. The Journal of Japan Atherosclerosis Society, 1993, 21, 585-589.	0.0	0
584	Lifestyles and biomarkers as risk and beneficial factors of stroke in the Circulatory Risk in Communities Study. Nosotchu, 2015, 37, 367-373.	0.0	0
585	Global landscape of the COVID-19 vaccination policy: Ensuring equitable access to quality-assured vaccines. GHM Open, $2021, , .$	0.1	0
586	Disparity in metabolic risk factors of non-communicable diseases between Palauans and Filipinos living in Palau. Nagoya Journal of Medical Science, 2017, 79, 157-165.	0.6	0
587	The number of leisure-time activities and risk of functional disability among Japanese older population: the JAGES cohort. Preventive Medicine Reports, 2022, 26, 101741.	0.8	0
588	Serum High-Sensitivity Cardiac Troponin T as an Independent Predictor for Incident Coronary Heart Disease in the Japanese General Population: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2022, , .	0.9	0
589	The association of family history of herpes zoster and the risk of incident herpes zoster: the SHEZ Study. Environmental Health and Preventive Medicine, 2022, 27, 22-22.	1.4	0
590	Positive psychological factors and the risk of pneumonia-associated mortality: Japan Collaborative Cohort Study. Journal of Psychosomatic Research, 2022, , 110971.	1.2	0
591	Association of Physical Activity with Aortic Disease in Japanese Men and Women: The Japan Collaborative Cohort Study. Journal of Atherosclerosis and Thrombosis, 2022, , .	0.9	0
592	Alcohol Consumption and Long-Term Mortality in Men with or without a History of Myocardial Infarction. Journal of Atherosclerosis and Thrombosis, 2022, , .	0.9	0