

Hiroyasu Iso

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6817594/hiroyasu-iso-publications-by-citations.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

551
papers

18,485
citations

72
h-index

113
g-index

609
ext. papers

21,437
ext. citations

4.5
avg, IF

6.53
L-index

#	Paper	IF	Citations
551	Serum cholesterol levels and six-year mortality from stroke in 350,977 men screened for the multiple risk factor intervention trial. <i>New England Journal of Medicine</i> , 1989 , 320, 904-10	59.2	895
550	Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2007 , 11, 41-50	2.5	442
549	Intake of fish and n3 fatty acids and risk of coronary heart disease among Japanese: the Japan Public Health Center-Based (JPHC) Study Cohort I. <i>Circulation</i> , 2006 , 113, 195-202	16.7	422
548	Intake of fish and omega-3 fatty acids and risk of stroke in women. <i>JAMA - Journal of the American Medical Association</i> , 2001 , 285, 304-12	27.4	343
547	The relationship between green tea and total caffeine intake and risk for self-reported type 2 diabetes among Japanese adults. <i>Annals of Internal Medicine</i> , 2006 , 144, 554-62	8	316
546	What has made the population of Japan healthy?. <i>Lancet, The</i> , 2011 , 378, 1094-105	40	284
545	Association of sleep duration with mortality from cardiovascular disease and other causes for Japanese men and women: the JACC study. <i>Sleep</i> , 2009 , 32, 295-301	1.1	263
544	Serum triglycerides and risk of coronary heart disease among Japanese men and women. <i>American Journal of Epidemiology</i> , 2001 , 153, 490-9	3.8	246
543	Prospective study of calcium, potassium, and magnesium intake and risk of stroke in women. <i>Stroke</i> , 1999 , 30, 1772-9	6.7	239
542	Carotid intima-media thickness and plaque characteristics as a risk factor for stroke in Japanese elderly men. <i>Stroke</i> , 2004 , 35, 2788-94	6.7	223
541	Fish, omega-3 polyunsaturated fatty acids, and mortality from cardiovascular diseases in a nationwide community-based cohort of Japanese men and women the JACC (Japan Collaborative Cohort Study for Evaluation of Cancer Risk) Study. <i>Journal of the American College of Cardiology</i> , 2009 , 53, 222-31	15.1	213
540	Association of dietary intake of soy, beans, and isoflavones with risk of cerebral and myocardial infarctions in Japanese populations: the Japan Public Health Center-based (JPHC) study cohort I. <i>Circulation</i> , 2007 , 116, 2553-62	16.7	201
539	Perceived mental stress and mortality from cardiovascular disease among Japanese men and women: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk Sponsored by Monbusho (JACC Study). <i>Circulation</i> , 2002 , 106, 1229-36	16.7	197
538	Linoleic acid, other fatty acids, and the risk of stroke. <i>Stroke</i> , 2002 , 33, 2086-93	6.7	169
537	Cardiovascular disease epidemiology in Asia: an overview. <i>Circulation Journal</i> , 2013 , 77, 1646-52	2.9	163
536	The joint impact on being overweight of self reported behaviours of eating quickly and eating until full: cross sectional survey. <i>BMJ, The</i> , 2008 , 337, a2002	5.9	158
535	Metabolic syndrome and the risk of ischemic heart disease and stroke among Japanese men and women. <i>Stroke</i> , 2007 , 38, 1744-51	6.7	157

534	Alcohol consumption and risk of stroke among middle-aged men: the JPHC Study Cohort I. <i>Stroke</i> , 2004 , 35, 1124-9	6.7	153
533	A low prevalence of coronary heart disease among subjects with increased high-density lipoprotein cholesterol levels, including those with plasma cholesteryl ester transfer protein deficiency. <i>Preventive Medicine</i> , 1998 , 27, 659-67	4.3	148
532	Coffee, green tea, black tea and oolong tea consumption and risk of mortality from cardiovascular disease in Japanese men and women. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, 230-40	5.1	146
531	Adult mortality attributable to preventable risk factors for non-communicable diseases and injuries in Japan: a comparative risk assessment. <i>PLoS Medicine</i> , 2012 , 9, e1001160	11.6	144
530	Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. <i>American Journal of Clinical Nutrition</i> , 2008 , 88, 195-202	7	140
529	A prospective cohort study of shift work and risk of ischemic heart disease in Japanese male workers. <i>American Journal of Epidemiology</i> , 2006 , 164, 128-35	3.8	134
528	Prospective study of fat and protein intake and risk of intraparenchymal hemorrhage in women. <i>Circulation</i> , 2001 , 103, 856-63	16.7	133
527	Fruit and vegetable intake and risk of total cancer and cardiovascular disease: Japan Public Health Center-Based Prospective Study. <i>American Journal of Epidemiology</i> , 2008 , 167, 59-70	3.8	125
526	Daily total physical activity level and premature death in men and women: results from a large-scale population-based cohort study in Japan (JPHC study). <i>Annals of Epidemiology</i> , 2008 , 18, 522-30	6.4	123
525	Randomized controlled trial for an effect of green tea consumption on insulin resistance and inflammation markers. <i>Journal of Nutritional Science and Vitaminology</i> , 2005 , 51, 335-42	1.1	122
524	Trends in the incidence of coronary heart disease and stroke and their risk factors in Japan, 1964 to 2003: the Akita-Osaka study. <i>Journal of the American College of Cardiology</i> , 2008 , 52, 71-9	15.1	118
523	Walking and sports participation and mortality from coronary heart disease and stroke. <i>Journal of the American College of Cardiology</i> , 2005 , 46, 1761-7	15.1	118
522	Profile of the JACC study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 1, S4-8	3.4	118
521	Fruit, vegetable and bean intake and mortality from cardiovascular disease among Japanese men and women: the JACC Study. <i>British Journal of Nutrition</i> , 2009 , 102, 285-92	3.6	112
520	Low-density lipoprotein cholesterol concentrations and death due to intraparenchymal hemorrhage: the Ibaraki Prefectural Health Study. <i>Circulation</i> , 2009 , 119, 2136-45	16.7	112
519	Cigarette smoking and risk of stroke and its subtypes among middle-aged Japanese men and women: the JPHC Study Cohort I. <i>Stroke</i> , 2004 , 35, 1248-53	6.7	111
518	Serum total homocysteine concentrations and risk of stroke and its subtypes in Japanese. <i>Circulation</i> , 2004 , 109, 2766-72	16.7	111
517	Prospective study of depressive symptoms and risk of stroke among Japanese. <i>Stroke</i> , 2001 , 32, 903-8	6.7	111

516	Dietary intake of calcium in relation to mortality from cardiovascular disease: the JACC Study. <i>Stroke</i> , 2006 , 37, 20-6	6.7	110
515	Body mass index and mortality from cardiovascular disease among Japanese men and women: the JACC study. <i>Stroke</i> , 2005 , 36, 1377-82	6.7	109
514	The impact of green tea and coffee consumption on the reduced risk of stroke incidence in Japanese population: the Japan public health center-based study cohort. <i>Stroke</i> , 2013 , 44, 1369-74	6.7	106
513	Serum fatty acids and fish intake in rural Japanese, urban Japanese, Japanese American and Caucasian American men. <i>International Journal of Epidemiology</i> , 1989 , 18, 374-81	7.8	106
512	Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2009 , 18, 240-7	2	104
511	Lifestyle and cardiovascular disease in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011 , 18, 83-8	4	100
510	Smoking cessation and mortality from cardiovascular disease among Japanese men and women: the JACC Study. <i>American Journal of Epidemiology</i> , 2005 , 161, 170-9	3.8	100
509	Alcohol consumption and mortality from stroke and coronary heart disease among Japanese men and women: the Japan collaborative cohort study. <i>Stroke</i> , 2008 , 39, 2936-42	6.7	99
508	Dietary fiber intake is associated with reduced risk of mortality from cardiovascular disease among Japanese men and women. <i>Journal of Nutrition</i> , 2010 , 140, 1445-53	4.1	97
507	Cohort profile of the Japan Collaborative Cohort Study at final follow-up. <i>Journal of Epidemiology</i> , 2013 , 23, 227-32	3.4	95
506	Serum cholesterol levels in relation to the incidence of cancer: the JPHC study cohorts. <i>International Journal of Cancer</i> , 2009 , 125, 2679-86	7.5	94
505	Trends for blood pressure and its contribution to stroke incidence in the middle-aged Japanese population: the Circulatory Risk in Communities Study (CIRCS). <i>Stroke</i> , 2009 , 40, 1571-7	6.7	91
504	Associations of dietary magnesium intake with mortality from cardiovascular disease: the JACC study. <i>Atherosclerosis</i> , 2012 , 221, 587-95	3.1	90
503	Effects of habitual alcohol intake on ambulatory blood pressure, heart rate, and its variability among Japanese men. <i>Hypertension</i> , 2009 , 53, 13-9	8.5	90
502	Blood pressure and the risk of stroke, cardiovascular disease, and all-cause mortality among Japanese: the JPHC Study. <i>American Journal of Hypertension</i> , 2009 , 22, 273-80	2.3	90
501	Relationships of age at menarche and menopause, and reproductive year with mortality from cardiovascular disease in Japanese postmenopausal women: the JACC study. <i>Journal of Epidemiology</i> , 2006 , 16, 177-84	3.4	90
500	Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 456-64	7	87
499	Dietary calcium intake and risks of stroke, its subtypes, and coronary heart disease in Japanese: the JPHC Study Cohort I. <i>Stroke</i> , 2008 , 39, 2449-56	6.7	87

498	Alcohol intake and the risk of cardiovascular disease in middle-aged Japanese men. <i>Stroke</i> , 1995 , 26, 767-73	6.7	87
497	Egg consumption, serum total cholesterol concentrations and coronary heart disease incidence: Japan Public Health Center-based prospective study. <i>British Journal of Nutrition</i> , 2006 , 96, 921-8	3.6	86
496	Obstructive sleep apnea and incident type 2 diabetes. <i>Sleep Medicine</i> , 2016 , 25, 156-161	4.6	86
495	Fat and protein intakes and risk of intraparenchymal hemorrhage among middle-aged Japanese. <i>American Journal of Epidemiology</i> , 2003 , 157, 32-9	3.8	83
494	Trends in the incidence of coronary heart disease and stroke and the prevalence of cardiovascular risk factors among Japanese men from 1963 to 1994. <i>American Journal of Medicine</i> , 2002 , 112, 104-9	2.4	82
493	The impact of the metabolic syndrome and its components on the incidence of ischemic heart disease and stroke: the Japan public health center-based study. <i>Hypertension Research</i> , 2009 , 32, 289-98	4.7	81
492	Accuracy of death certificate diagnosis of intracranial hemorrhage and nonhemorrhagic stroke. The Minnesota Heart Survey. <i>American Journal of Epidemiology</i> , 1990 , 132, 993-8	3.8	81
491	Serum and dietary magnesium and risk of ischemic stroke: the Atherosclerosis Risk in Communities Study. <i>American Journal of Epidemiology</i> , 2009 , 169, 1437-44	3.8	79
490	Message from the New Editor-in-Chief. <i>Journal of Epidemiology</i> , 2012 , 22, 1	3.4	78
489	Alcohol consumption and mortality among middle-aged and elderly Japanese men and women. <i>Annals of Epidemiology</i> , 2005 , 15, 590-7	6.4	77
488	Application of computer tomography-oriented criteria for stroke subtype classification in a prospective study. <i>Annals of Epidemiology</i> , 2000 , 10, 81-7	6.4	76
487	Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. <i>Clinical Nutrition</i> , 2013 , 32, 300-8	5.9	74
486	Serum total cholesterol levels and risk of mortality from stroke and coronary heart disease in Japanese: the JACC study. <i>Atherosclerosis</i> , 2007 , 194, 415-20	3.1	74
485	Dietary folate and vitamin b6 and B12 intake in relation to mortality from cardiovascular diseases: Japan collaborative cohort study. <i>Stroke</i> , 2010 , 41, 1285-9	6.7	73
484	Changes in coronary heart disease risk among Japanese. <i>Circulation</i> , 2008 , 118, 2725-9	16.7	73
483	Impact of the revision of a nutrient database on the validity of a self-administered food frequency questionnaire (FFQ). <i>Journal of Epidemiology</i> , 2006 , 16, 107-16	3.4	72
482	Dietary patterns and all-cause, cancer, and cardiovascular disease mortality in Japanese men and women: The Japan public health center-based prospective study. <i>PLoS ONE</i> , 2017 , 12, e0174848	3.7	71
481	Serum total and non-high-density lipoprotein cholesterol and the risk prediction of cardiovascular events - the JALS-ECC -. <i>Circulation Journal</i> , 2010 , 74, 1346-56	2.9	69

480	Healthy lifestyle behaviours and cardiovascular mortality among Japanese men and women: the Japan collaborative cohort study. <i>European Heart Journal</i> , 2012 , 33, 467-77	9.5	68
479	Mild retinopathy is a risk factor for cardiovascular mortality in Japanese with and without hypertension: the Ibaraki Prefectural Health Study. <i>Circulation</i> , 2011 , 124, 2502-11	16.7	65
478	Low-density lipoprotein cholesterol and risk of coronary heart disease among Japanese men and women: the Circulatory Risk in Communities Study (CIRCS). <i>Preventive Medicine</i> , 2011 , 52, 381-6	4.3	64
477	Dietary intake of saturated fatty acids and mortality from cardiovascular disease in Japanese: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 759-65	7	64
476	Relation between serum total cholesterol level and cardiovascular disease stratified by sex and age group: a pooled analysis of 65 594 individuals from 10 cohort studies in Japan. <i>Journal of the American Heart Association</i> , 2012 , 1, e001974	6	64
475	Metabolic syndrome and all-cause and cardiovascular disease mortality: Japan Public Health Center-based Prospective (JPHC) Study. <i>Circulation Journal</i> , 2009 , 73, 878-84	2.9	63
474	The effects of smoking and smoking cessation on mortality from cardiovascular disease among Japanese: pooled analysis of three large-scale cohort studies in Japan. <i>Tobacco Control</i> , 2010 , 19, 50-7	5.3	62
473	Diabetes mellitus and risk of stroke and its subtypes among Japanese: the Japan public health center study. <i>Stroke</i> , 2011 , 42, 2611-4	6.7	62
472	Risk and protective factors related to mortality from pneumonia among middleaged and elderly community residents: the JACC Study. <i>Journal of Epidemiology</i> , 2007 , 17, 194-202	3.4	62
471	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCH-JAPAN study. <i>Journal of Clinical Lipidology</i> , 2018 , 12, 674-684.e5	4.9	61
470	Social support and stroke and coronary heart disease: the JPHC study cohorts II. <i>Stroke</i> , 2008 , 39, 768-756.7	5.7	59
469	Underweight as a predictor of diabetes in older adults: a large cohort study. <i>Diabetes Care</i> , 2008 , 31, 583-4	14.6	59
468	Proportions of stroke subtypes among men and women > or =40 years of age in an urban Japanese city in 1992, 1997, and 2002. <i>Stroke</i> , 2006 , 37, 1374-8	6.7	59
467	Soft drink intake in relation to incident ischemic heart disease, stroke, and stroke subtypes in Japanese men and women: the Japan Public Health Centre-based study cohort I. <i>American Journal of Clinical Nutrition</i> , 2012 , 96, 1390-7	7	57
466	Perceived level of life enjoyment and risks of cardiovascular disease incidence and mortality: the Japan public health center-based study. <i>Circulation</i> , 2009 , 120, 956-63	16.7	57
465	Risk factors for fatal subarachnoid hemorrhage: the Japan Collaborative Cohort Study. <i>Stroke</i> , 2003 , 34, 2781-7	6.7	57
464	Dietary intakes of antioxidant vitamins and mortality from cardiovascular disease: the Japan Collaborative Cohort Study (JACC) study. <i>Stroke</i> , 2011 , 42, 1665-72	6.7	55
463	Smoking raises the risk of total and ischemic strokes in hypertensive men. <i>Hypertension Research</i> , 2003 , 26, 209-17	4.7	55

462	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>Annals of Epidemiology</i> , 2015 , 25, 512-518.e3	6.4	54
461	C-reactive protein levels and risk of mortality from cardiovascular disease in Japanese: the JACC Study. <i>Atherosclerosis</i> , 2009 , 207, 291-7	3.1	53
460	The criteria for metabolic syndrome and the national health screening and education system in Japan. <i>Epidemiology and Health</i> , 2017 , 39, e2017003	5.6	52
459	Social support and cancer incidence and mortality: the JPHC study cohort II. <i>Cancer Causes and Control</i> , 2013 , 24, 847-60	2.8	52
458	Relationship between sleep-disordered breathing and blood pressure levels in community-based samples of Japanese men. <i>Hypertension Research</i> , 2004 , 27, 479-84	4.7	52
457	Usual alcohol consumption and arterial oxygen desaturation during sleep. <i>JAMA - Journal of the American Medical Association</i> , 2004 , 292, 923-5	27.4	52
456	Fasting and non-fasting triglycerides and risk of ischemic cardiovascular disease in Japanese men and women: the Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2014 , 237, 361-8	3.1	51
455	Community-based education classes for hypertension control. A 1.5-year randomized controlled trial. <i>Hypertension</i> , 1996 , 27, 968-74	8.5	51
454	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2015 , 101, 1029-37	7	50
453	Body mass index and risk of stroke and myocardial infarction in a relatively lean population: meta-analysis of 16 Japanese cohorts using individual data. <i>Circulation: Cardiovascular Quality and Outcomes</i> , 2010 , 3, 498-505	5.8	48
452	Rice intake is associated with reduced risk of mortality from cardiovascular disease in Japanese men but not women. <i>Journal of Nutrition</i> , 2011 , 141, 595-602	4.1	48
451	Effects of a long-term hypertension control program on stroke incidence and prevalence in a rural community in northeastern Japan. <i>Stroke</i> , 1998 , 29, 1510-8	6.7	48
450	Serum total cholesterol and mortality in a Japanese population. <i>Journal of Clinical Epidemiology</i> , 1994 , 47, 961-9	5.7	48
449	Associations between dietary intakes of iron, copper and zinc with risk of type 2 diabetes mellitus: A large population-based prospective cohort study. <i>Clinical Nutrition</i> , 2018 , 37, 667-674	5.9	47
448	Dietary intake of saturated fatty acids and incident stroke and coronary heart disease in Japanese communities: the JPHC Study. <i>European Heart Journal</i> , 2013 , 34, 1225-32	9.5	47
447	Intake of folate, vitamin B6 and vitamin B12 and the risk of CHD: the Japan Public Health Center-Based Prospective Study Cohort I. <i>Journal of the American College of Nutrition</i> , 2008 , 27, 127-36	3.5	47
446	Blood pressure-lowering effect of Shinrin-yoku (Forest bathing): a systematic review and meta-analysis. <i>BMC Complementary and Alternative Medicine</i> , 2017 , 17, 409	4.7	46
445	Association of Breakfast Intake With Incident Stroke and Coronary Heart Disease: The Japan Public Health Center-Based Study. <i>Stroke</i> , 2016 , 47, 477-81	6.7	46

444	Social network, social support, and risk of incident stroke: Atherosclerosis Risk in Communities study. <i>Stroke</i> , 2014 , 45, 2868-73	6.7	46
443	Cigarette smoking and risk of coronary heart disease incidence among middle-aged Japanese men and women: the JPHC Study Cohort I. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 207-13		46
442	Genetic Predisposition to Ischemic Stroke: A Polygenic Risk Score. <i>Stroke</i> , 2017 , 48, 253-258	6.7	45
441	Relations between protein intake and blood pressure in Japanese men and women: the Circulatory Risk in Communities Study (CIRCS). <i>American Journal of Clinical Nutrition</i> , 2009 , 90, 377-84	7	45
440	A low level of C-reactive protein in Japanese adults and its association with cardiovascular risk factors: the Japan NCVC-Collaborative Inflammation Cohort (JNIC) study. <i>Atherosclerosis</i> , 2007 , 194, 238-44	3.1	45
439	Frequency of food intake and estimated nutrient intake among men and women: the JACC Study. <i>Journal of Epidemiology</i> , 2005 , 15 Suppl 1, S24-42	3.4	45
438	Serum total homocysteine concentrations and risk of mortality from stroke and coronary heart disease in Japanese: The JACC study. <i>Atherosclerosis</i> , 2008 , 198, 412-8	3.1	44
437	Body fat distribution and the risk of hypertension and diabetes among Japanese men and women. <i>Hypertension Research</i> , 2008 , 31, 851-7	4.7	43
436	Effects of serum B vitamins on elevated plasma homocysteine levels associated with the mutation of methylenetetrahydrofolate reductase gene in Japanese. <i>Atherosclerosis</i> , 2002 , 164, 321-8	3.1	43
435	Effect of coffee consumption on all-cause and total cancer mortality: findings from the JACC study. <i>European Journal of Epidemiology</i> , 2011 , 26, 285-93	12.1	42
434	Age- and gender-specific BMI in terms of the lowest mortality in Japanese general population. <i>Obesity</i> , 2008 , 16, 2348-55	8	42
433	Prognostic value of plasma high-sensitivity C-reactive protein levels in Japanese patients with stable coronary artery disease: the Japan NCVC-Collaborative Inflammation Cohort (JNIC) Study. <i>Atherosclerosis</i> , 2009 , 207, 272-6	3.1	41
432	Associations of all-cause mortality with census-based neighbourhood deprivation and population density in Japan: a multilevel survival analysis. <i>PLoS ONE</i> , 2014 , 9, e97802	3.7	41
431	High serum total cholesterol levels is a risk factor of ischemic stroke for general Japanese population: the JPHC study. <i>Atherosclerosis</i> , 2012 , 221, 565-9	3.1	40
430	Nutrition and disease in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). <i>Asian Pacific Journal of Cancer Prevention</i> , 2007 , 8 Suppl, 35-80	1.7	40
429	Daily Total Physical Activity and Incident Stroke: The Japan Public Health Center-Based Prospective Study. <i>Stroke</i> , 2017 , 48, 1730-1736	6.7	39
428	Association between non-high-density lipoprotein cholesterol levels and the incidence of coronary heart disease among Japanese: the Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2011 , 18, 454-63	4	39
427	Self-reported stroke and myocardial infarction had adequate sensitivity in a population-based prospective study JPHC (Japan Public Health Center)-based Prospective Study. <i>Journal of Clinical Epidemiology</i> , 2009 , 62, 667-73	5.7	39

426	Cigarette smoking and risk of disabling dementia in a Japanese rural community: a nested case-control study. <i>Cerebrovascular Diseases</i> , 2008 , 25, 324-31	3.2	39
425	Metabolic Syndrome and the Risk of Ischemic Heart Disease and Stroke among Middle-Aged Japanese. <i>Hypertension Research</i> , 2008 , 31, 1887-94	4.7	39
424	Age-specific relationship between blood pressure and the risk of total and cardiovascular mortality in Japanese men and women. <i>Hypertension Research</i> , 2005 , 28, 901-9	4.7	39
423	Incidences of Herpes Zoster and Postherpetic Neuralgia in Japanese Adults Aged 50 Years and Older From a Community-based Prospective Cohort Study: The SHEZ Study. <i>Journal of Epidemiology</i> , 2015 , 25, 617-25	3.4	38
422	Anemia and reduced kidney function as risk factors for new onset of atrial fibrillation (from the Ibaraki prefectural health study). <i>American Journal of Cardiology</i> , 2015 , 115, 328-33	3	38
421	Influenza (H1N1) 2009 outbreak and school closure, Osaka Prefecture, Japan. <i>Emerging Infectious Diseases</i> , 2009 , 15, 1685	10.2	38
420	Prognostic impact of supraventricular premature complexes in community-based health checkups: the Ibaraki Prefectural Health Study. <i>European Heart Journal</i> , 2015 , 36, 170-8	9.5	37
419	Carotid artery wall thickness and risk of stroke subtypes: the atherosclerosis risk in communities study. <i>Stroke</i> , 2011 , 42, 397-403	6.7	37
418	Adult height and the risk of cardiovascular disease among middle aged men and women in Japan. <i>European Journal of Epidemiology</i> , 2011 , 26, 13-21	12.1	37
417	Magnesium intake and risk of self-reported type 2 diabetes among Japanese. <i>Journal of the American College of Nutrition</i> , 2010 , 29, 99-106	3.5	37
416	Effects of a laughter and exercise program on physiological and psychological health among community-dwelling elderly in Japan: randomized controlled trial. <i>Geriatrics and Gerontology International</i> , 2013 , 13, 152-60	2.9	36
415	Chronic kidney disease and drinking status in relation to risks of stroke and its subtypes: the Circulatory Risk in Communities Study (CIRCS). <i>Stroke</i> , 2011 , 42, 2531-7	6.7	36
414	Relationship between cold temperature and cardiovascular mortality, with assessment of effect modification by individual characteristics: Ibaraki Prefectural Health Study. <i>Circulation Journal</i> , 2013 , 77, 1854-61	2.9	35
413	Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. <i>Journal of Nutrition</i> , 2017 , 147, 1749-1756	4.756	33
412	High-density lipoprotein subclasses and risk of stroke and its subtypes in Japanese population: the Circulatory Risk in Communities Study. <i>Stroke</i> , 2013 , 44, 327-33	6.7	33
411	Type A behaviour and risk of coronary heart disease: the JPHC Study. <i>International Journal of Epidemiology</i> , 2008 , 37, 1395-405	7.8	33
410	Angiotensinogen T174M and M235T variants, sodium intake and hypertension among non-drinking, lean Japanese men and women. <i>Journal of Hypertension</i> , 2000 , 18, 1197-206	1.9	33
409	The dose-response relationship between body mass index and the risk of incident stage B chronic kidney disease in a general Japanese population: the Ibaraki prefectural health study (IPHS). <i>Journal of Epidemiology</i> , 2014 , 24, 444-51	3.4	32

408	Milk drinking and mortality: findings from the Japan collaborative cohort study. <i>Journal of Epidemiology</i> , 2015 , 25, 66-73	3.4	32
407	Development of a point-based prediction model for the incidence of total stroke: Japan public health center study. <i>Stroke</i> , 2013 , 44, 1295-302	6.7	32
406	Physician-pharmacist cooperation program for blood pressure control in patients with hypertension: a randomized-controlled trial. <i>American Journal of Hypertension</i> , 2010 , 23, 1144-52	2.3	32
405	Associations between metabolic syndrome and mortality from cardiovascular disease in Japanese general population, findings on overweight and non-overweight individuals. Ibaraki Prefectural Health Study. <i>Circulation Journal</i> , 2009 , 73, 1635-42	2.9	32
404	Trends for coronary heart disease and its risk factors in Japan: epidemiologic and pathologic studies. <i>Japanese Circulation Journal</i> , 1990 , 54, 428-35		32
403	Associations between copper and zinc intakes from diet and mortality from cardiovascular disease in a large population-based prospective cohort study. <i>Journal of Nutritional Biochemistry</i> , 2018 , 56, 126-132	6.3	31
402	Combined effect of blood pressure and total cholesterol levels on long-term risks of subtypes of cardiovascular death: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan. <i>Hypertension</i> , 2015 , 65, 517-24	8.5	31
401	gamma-Glutamyltranspeptidase and incident stroke among Japanese men and women: the Circulatory Risk in Communities Study (CIRCS). <i>Stroke</i> , 2010 , 41, 385-8	6.7	31
400	Epidemiology of cerebrovascular disease: stroke epidemic in Japan. <i>Journal of Epidemiology</i> , 1996 , 6, S43-7	3.4	31
399	The relation of body fat distribution and body mass with haemoglobin A1c, blood pressure and blood lipids in urban Japanese men. <i>International Journal of Epidemiology</i> , 1991 , 20, 88-94	7.8	31
398	Antigens of Tissue Plasminogen Activator and Plasminogen Activator Inhibitor 1: Correlates in Nonsmoking Japanese and Caucasian Men and Women. <i>Thrombosis and Haemostasis</i> , 1993 , 70, 475-480	7	31
397	Rice consumption is not associated with risk of cardiovascular disease morbidity or mortality in Japanese men and women: a large population-based, prospective cohort study. <i>American Journal of Clinical Nutrition</i> , 2014 , 100, 199-207	7	30
396	Alcohol consumption patterns in Thailand and their relationship with non-communicable disease. <i>BMC Public Health</i> , 2015 , 15, 1297	4.1	30
395	Incidence of metabolic syndrome according to combinations of lifestyle factors among middle-aged Japanese male workers. <i>Preventive Medicine</i> , 2010 , 51, 118-22	4.3	30
394	Prospective Study on Alcohol Intake and Risk of Subarachnoid Hemorrhage Among Japanese Men and Women. <i>Alcoholism: Clinical and Experimental Research</i> , 2000 , 24, 386-389	3.7	30
393	Heavy alcohol consumption and risk of atrial fibrillation. The Circulatory Risk in Communities Study (CIRCS). <i>Circulation Journal</i> , 2014 , 78, 955-61	2.9	29
392	Population profile and residential environment of an urban poor community in Dhaka, Bangladesh. <i>Environmental Health and Preventive Medicine</i> , 2017 , 22, 1	4.2	29
391	Impact of neighborhood socioeconomic conditions on the risk of stroke in Japan. <i>Journal of Epidemiology</i> , 2015 , 25, 254-60	3.4	29

390	C-reactive protein levels and risk of stroke and its subtype in Japanese: The Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2011 , 217, 187-93	3.1	29
389	Risk factors for snoring among Japanese men and women: a community-based cross-sectional study. <i>Sleep and Breathing</i> , 2011 , 15, 63-9	3.1	29
388	Education, social roles, and the risk of cardiovascular disease among middle-aged Japanese women: the JPHC Study Cohort I. <i>Stroke</i> , 2008 , 39, 2886-90	6.7	29
387	Ankle-arm blood pressure index and cardiovascular risk factors in elderly Japanese men. <i>Hypertension Research</i> , 2003 , 26, 377-82	4.7	29
386	Calcium intake and blood pressure in seven Japanese populations. <i>American Journal of Epidemiology</i> , 1991 , 133, 776-83	3.8	29
385	The trend of coronary heart disease and its risk factors based on epidemiological investigations. <i>Japanese Circulation Journal</i> , 1987 , 51, 319-24		29
384	Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. <i>BMJ, The</i> , 2020 , 368, m34	5.9	28
383	Coping strategies and risk of cardiovascular disease incidence and mortality: the Japan Public Health Center-based prospective Study. <i>European Heart Journal</i> , 2016 , 37, 890-899	9.5	28
382	Pulse pressure is an independent risk factor for stroke among middle-aged Japanese with normal systolic blood pressure: the JPHC study. <i>Journal of Hypertension</i> , 2011 , 29, 319-24	1.9	28
381	Smoking cessation: predictive factors among middle-aged Japanese. <i>Nicotine and Tobacco Research</i> , 2010 , 12, 1050-4	4.9	28
380	Alcohol consumption, social support, and risk of stroke and coronary heart disease among Japanese men: the JPHC Study. <i>Alcoholism: Clinical and Experimental Research</i> , 2009 , 33, 1025-32	3.7	28
379	Associations of anger, anxiety, and depressive symptoms with carotid arterial wall thickness: the multi-ethnic study of atherosclerosis. <i>Psychosomatic Medicine</i> , 2012 , 74, 517-25	3.7	28
378	Association of lymphocyte sub-populations with clustered features of metabolic syndrome in middle-aged Japanese men. <i>Atherosclerosis</i> , 2004 , 173, 295-300	3.1	28
377	Impact of Alcohol Intake and Drinking Patterns on Mortality From All Causes and Major Causes of Death in a Japanese Population. <i>Journal of Epidemiology</i> , 2018 , 28, 140-148	3.4	27
376	Development of a Risk Equation for the Incidence of Coronary Artery Disease and Ischemic Stroke for Middle-Aged Japanese - Japan Public Health Center-Based Prospective Study. <i>Circulation Journal</i> , 2016 , 80, 1386-95	2.9	27
375	Alkaline phosphatase and risk of stroke among Japanese: the Circulatory Risk in Communities Study (CIRCS). <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2013 , 22, 1046-55	2.8	27
374	The combination of Helicobacter pylori- and cytotoxin-associated gene-A seropositivity in relation to the risk of myocardial infarction in middle-aged Japanese: The Japan Public Health Center-based study. <i>Atherosclerosis</i> , 2013 , 230, 67-72	3.1	27
373	Impact of obesity on incident hypertension independent of weight gain among nonhypertensive Japanese: the Ibaraki Prefectural Health Study (IPHS). <i>Journal of Hypertension</i> , 2012 , 30, 1122-8	1.9	27

372	Relationships between sleep-disordered breathing and blood pressure and excessive daytime sleepiness among truck drivers. <i>Hypertension Research</i> , 2006 , 29, 605-10	4.7	27
371	The impact of C-reactive protein on risk of stroke, stroke subtypes, and ischemic heart disease in middle-aged Japanese: the Japan public health center-based study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19, 756-66	4	27
370	Association of high-density lipoprotein cholesterol concentration with different types of stroke and coronary heart disease: The Japan Public Health Center-based prospective (JPHC) study. <i>Atherosclerosis</i> , 2017 , 265, 147-154	3.1	26
369	Relation of serum folic acid and folic acid levels to cardiovascular disease-related mortality among Japanese men and women. <i>Journal of Epidemiology</i> , 2012 , 22, 402-10	3.4	26
368	Associations of dietary iron intake with mortality from cardiovascular disease: the JACC study. <i>Journal of Epidemiology</i> , 2012 , 22, 484-93	3.4	26
367	Diabetes and the risk of coronary heart disease in the general Japanese population: the Japan Public Health Center-based prospective (JPHC) study. <i>Atherosclerosis</i> , 2011 , 216, 187-91	3.1	26
366	Association between non-high-density lipoprotein cholesterol concentrations and mortality from coronary heart disease among Japanese men and women: the Ibaraki Prefectural Health Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010 , 17, 30-6	4	26
365	Relationship between obesity and incident diabetes in middle-aged and older Japanese adults: the Ibaraki Prefectural Health Study. <i>Mayo Clinic Proceedings</i> , 2010 , 85, 36-40	6.4	26
364	Caregiver burden for impaired elderly Japanese with prevalent stroke and dementia under long-term care insurance system. <i>Cerebrovascular Diseases</i> , 2008 , 25, 234-40	3.2	26
363	Validity of the self-administered food frequency questionnaire used in the 5-year follow-up survey for the JPHC Study to assess folate, vitamin B6 and B12 intake: comparison with dietary records and blood level. <i>Journal of Epidemiology</i> , 2003 , 13, S98-101	3.4	26
362	Relationships of differential leukocyte and lymphocyte subpopulations with carotid atherosclerosis in elderly men. <i>Journal of Clinical Immunology</i> , 2003 , 23, 469-76	5.7	26
361	Trends in dietary intake of folate, vitamins B6, and B12 among Japanese adults in two rural communities from 1974 through 2001. <i>Journal of Epidemiology</i> , 2005 , 15, 29-37	3.4	26
360	The relation of anger expression with blood pressure levels and hypertension in rural and urban Japanese communities. <i>Journal of Hypertension</i> , 2002 , 20, 21-7	1.9	26
359	The Japanese food score and risk of all-cause, CVD and cancer mortality: the Japan Collaborative Cohort Study. <i>British Journal of Nutrition</i> , 2018 , 120, 464-471	3.6	26
358	Effects of Coffee and Tea Consumption on Glucose Metabolism: A Systematic Review and Network Meta-Analysis. <i>Nutrients</i> , 2018 , 11,	6.7	26
357	Adult height and body mass index in relation to risk of total stroke and its subtypes: the circulatory risk in communities study. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014 , 23, 667-74	2.8	25
356	An inverse correlation of VZV skin-test reaction, but not antibody, with severity of herpes zoster skin symptoms and zoster-associated pain. <i>Journal of Dermatological Science</i> , 2013 , 69, 243-9	4.3	25
355	A community-based survey of varicella-zoster virus-specific immune responses in the elderly. <i>Journal of Clinical Virology</i> , 2012 , 55, 46-50	14.5	25

354	Long-term exposure to particulate matter in relation to mortality and incidence of cardiovascular disease: the JPHC Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 296-309	4	25
353	Associations of psychological distress with metabolic syndrome among Japanese urban residents. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011 , 18, 396-402	4	25
352	Prevalence and correlates of carotid atherosclerosis among elderly Japanese men. <i>Atherosclerosis</i> , 2004 , 172, 353-9	3.1	25
351	Watching Television and Risk of Mortality From Pulmonary Embolism Among Japanese Men and Women: The JACC Study (Japan Collaborative Cohort). <i>Circulation</i> , 2016 , 134, 355-7	16.7	24
350	Prevalence of non-communicable disease risk factors among poor shantytown residents in Dhaka, Bangladesh: a community-based cross-sectional survey. <i>BMJ Open</i> , 2017 , 7, e014710	3	24
349	Education in relation to incidence of and mortality from cancer and cardiovascular disease in Japan. <i>European Journal of Public Health</i> , 2008 , 18, 466-72	2.1	24
348	Prospective study of major and minor ST-T abnormalities and risk of stroke among Japanese. <i>Stroke</i> , 2003 , 34, e250-3	6.7	24
347	Association between mortality and incidence rates of coronary heart disease and stroke: The Japan Public Health Center-based prospective (JPHC) study. <i>International Journal of Cardiology</i> , 2016 , 222, 281-286	3.2	24
346	The Association of Having a Late Dinner or Bedtime Snack and Skipping Breakfast with Overweight in Japanese Women. <i>Journal of Obesity</i> , 2019 , 2019, 2439571	3.7	23
345	Identification of dementia using standard clinical assessments by primary care physicians in Japan. <i>Geriatrics and Gerontology International</i> , 2018 , 18, 738-744	2.9	23
344	Bowel Movement Frequency, Laxative Use, and Mortality From Coronary Heart Disease and Stroke Among Japanese Men and Women: The Japan Collaborative Cohort (JACC) Study. <i>Journal of Epidemiology</i> , 2016 , 26, 242-8	3.4	23
343	Dietary intakes of fat and total mortality among Japanese populations with a low fat intake: the Japan Collaborative Cohort (JACC) Study. <i>Nutrition and Metabolism</i> , 2014 , 11, 12	4.6	23
342	Geographically-based discrimination is a social determinant of mental health in a deprived or stigmatized area in Japan: a cross-sectional study. <i>Social Science and Medicine</i> , 2012 , 75, 1015-21	5.1	23
341	Nocturnal intermittent hypoxia and metabolic syndrome; the effect of being overweight: the CIRCS study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010 , 17, 369-77	4	23
340	Associations of sleep-disordered breathing with excessive daytime sleepiness and blood pressure in Japanese women. <i>Hypertension Research</i> , 2008 , 31, 501-6	4.7	23
339	Metabolic syndrome and urinary cGMP excretion in general population. <i>Atherosclerosis</i> , 2007 , 190, 423-8	3.1	23
338	Polymorphisms of the Beta Fibrinogen Gene and Plasma Fibrinogen Concentration in Caucasian and Japanese Population Samples. <i>Thrombosis and Haemostasis</i> , 1995 , 73, 106-111	7	23
337	Serum Linolenic and other Ω 3 fatty acids, and risk of disabling dementia: Community-based nested case-control study. <i>Clinical Nutrition</i> , 2017 , 36, 793-797	5.9	22

336	Death by suicide and other externally caused injuries after stroke in Japan (1990-2010): the Japan Public Health Center-based prospective study. <i>Psychosomatic Medicine</i> , 2014 , 76, 452-9	3.7	22
335	Prediagnostic plasma antibody levels to periodontopathic bacteria and risk of coronary heart disease. <i>International Heart Journal</i> , 2012 , 53, 209-14	1.8	22
334	A Japanese health success story: trends in cardiovascular diseases, their risk factors, and the contribution of public health and personalized approaches. <i>EPMA Journal</i> , 2011 , 2, 49-57	8.8	22
333	Incidence and survival trends for childhood cancer in Osaka, Japan, 1973-2001. <i>Cancer Science</i> , 2010 , 101, 787-92	6.9	22
332	Serum vitamin C concentration and hs-CRP level in middle-aged Japanese men and women. <i>Atherosclerosis</i> , 2010 , 208, 496-500	3.1	22
331	The Association between Concentrations of Green Tea and Blood Glucose Levels. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2009 , 44, 41-5	3.1	22
330	Self-reported snoring frequency and incidence of cardiovascular disease: the Circulatory Risk in Communities Study (CIRCS). <i>Journal of Epidemiology</i> , 2012 , 22, 295-301	3.4	22
329	High sodium intake strengthens the association between angiotensinogen T174M polymorphism and blood pressure levels among lean men and women: a community-based study. <i>Hypertension Research</i> , 2004 , 27, 53-60	4.7	22
328	Vascular dysfunction in patients with chronic arsenosis can be reversed by reduction of arsenic exposure. <i>Environmental Health Perspectives</i> , 2005 , 113, 339-41	8.4	22
327	Seaweed intake and risk of cardiovascular disease: the Japan Public Health Center-based Prospective (JPHC) Study. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1449-1455	7	21
326	Age trajectories of glycaemic traits in non-diabetic South Asian and white individuals: the Whitehall II cohort study. <i>Diabetologia</i> , 2015 , 58, 534-42	10.3	21
325	Association between Chlamydia pneumoniae infection and risk of coronary heart disease for Japanese: the JPHC study. <i>Atherosclerosis</i> , 2014 , 233, 338-342	3.1	21
324	Age-Period-Cohort analysis of mortality due to ischemic heart disease in Japan, 1955 to 2000. <i>Circulation Journal</i> , 2008 , 72, 966-72	2.9	21
323	Dietary patterns and breast cancer risk in a prospective Japanese study. <i>Breast Cancer</i> , 2017 , 24, 152-160	3.4	20
322	Non-communicable disease risk factor profile among public employees in a regional city in northern Ethiopia. <i>Scientific Reports</i> , 2018 , 8, 9298	4.9	20
321	Hemoglobin a1c levels and the risk of cardiovascular disease in people without known diabetes: a population-based cohort study in Japan. <i>Medicine (United States)</i> , 2015 , 94, e785	1.8	20
320	Salt preference and mortality from stroke and coronary heart disease for Japanese men and women: the JACC study. <i>Preventive Medicine</i> , 2012 , 54, 32-7	4.3	20
319	Total and high molecular weight adiponectin levels and risk of cardiovascular disease in individuals with high blood glucose levels. <i>Atherosclerosis</i> , 2013 , 229, 222-7	3.1	20

318	Diagnosed diabetes and premature death among middle-aged Japanese: results from a large-scale population-based cohort study in Japan (JPHC study). <i>BMJ Open</i> , 2015 , 5, e007736	3	20
317	Tuberculosis infection among homeless persons and caregivers in a high-tuberculosis-prevalence area in Japan: a cross-sectional study. <i>BMC Infectious Diseases</i> , 2011 , 11, 22	4	20
316	Serum Albumin and High-Sensitivity C-reactive Protein are Independent Risk Factors of Chronic Kidney Disease in Middle-Aged Japanese Individuals: the Circulatory Risk in Communities Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2016 , 23, 1089-98	4	20
315	Changes in the Employment Status and Risk of Stroke and Stroke Types. <i>Stroke</i> , 2017 , 48, 1176-1182	6.7	19
314	Serum coenzyme Q10 and risk of disabling dementia: the Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2014 , 237, 400-3	3.1	19
313	The Circulatory Risk in Communities Study (CIRCS): A Long-Term Epidemiological Study for Lifestyle-Related Disease Among Japanese Men and Women Living in Communities. <i>Journal of Epidemiology</i> , 2019 , 29, 83-91	3.4	19
312	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level: EPOCH-JAPAN (Evidence for Cardiovascular Prevention From Observational Cohorts in Japan). <i>Hypertension</i> , 2019 , 73, 52-59	8.5	19
311	Disease history and risk of comorbidity in women's life course: a comprehensive analysis of the Japan Nurses' Health Study baseline survey. <i>BMJ Open</i> , 2015 , 5, e006360	3	18
310	The Shozu Herpes Zoster (SHEZ) study: rationale, design, and description of a prospective cohort study. <i>Journal of Epidemiology</i> , 2012 , 22, 167-74	3.4	18
309	Low serum LDL cholesterol levels are associated with elevated mortality from liver cancer in Japan: the Ibaraki Prefectural health study. <i>Tohoku Journal of Experimental Medicine</i> , 2013 , 229, 203-11	2.4	18
308	Relationship of smoking and smoking cessation with ankle-to-arm blood pressure index in elderly Japanese men. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 243-8		18
307	Alpha-adducin G460W polymorphism, urinary sodium excretion, and blood pressure in community-based samples. <i>American Journal of Hypertension</i> , 2004 , 17, 385-90	2.3	18
306	Working Hours and Risk of Acute Myocardial Infarction and Stroke Among Middle-Aged Japanese Men - The Japan Public Health Center-Based Prospective Study Cohort II. <i>Circulation Journal</i> , 2019 , 83, 1072-1079	2.9	17
305	Effects of a traditional herbal medicine on peripheral blood flow in women experiencing peripheral coldness: a randomized controlled trial. <i>BMC Complementary and Alternative Medicine</i> , 2015 , 15, 105	4.7	17
304	VZV skin-test reaction, but not antibody, is an important predictive factor for postherpetic neuralgia. <i>Journal of Dermatological Science</i> , 2015 , 79, 235-40	4.3	17
303	Relationship Between Dietary Vitamin D and Deaths From Stroke and Coronary Heart Disease: The Japan Collaborative Cohort Study. <i>Stroke</i> , 2018 , 49, 454-457	6.7	17
302	Salty Food Preference and Intake and Risk of Gastric Cancer: The JACC Study. <i>Journal of Epidemiology</i> , 2016 , 26, 92-7	3.4	17
301	Socioeconomic status inconsistency and risk of stroke among Japanese middle-aged women. <i>Stroke</i> , 2014 , 45, 2592-8	6.7	17

300	Leukocyte count is an independent predictor for risk of acute myocardial infarction in middle-aged Japanese men. <i>Atherosclerosis</i> , 2007 , 195, 147-52	3.1	17
299	Prospective study of seaweed consumption and thyroid cancer incidence in women: the Japan collaborative cohort study. <i>European Journal of Cancer Prevention</i> , 2016 , 25, 239-45	2	17
298	Passive smoking and mortality from aortic dissection or aneurysm. <i>Atherosclerosis</i> , 2017 , 263, 145-150	3.1	16
297	Serum uric acid and risk of stroke and its types: the Circulatory Risk in Communities Study (CIRCS). <i>Hypertension Research</i> , 2020 , 43, 313-321	4.7	16
296	Impact of Hypertension and Subclinical Organ Damage on the Incidence of Cardiovascular Disease Among Japanese Residents at the Population and Individual Levels - The Circulatory Risk in Communities Study (CIRCS). <i>Circulation Journal</i> , 2017 , 81, 1022-1028	2.9	16
295	Plasma fibrinogen concentrations and risk of stroke and its subtypes among Japanese men and women. <i>Stroke</i> , 2006 , 37, 2488-92	6.7	16
294	Screening for sleep-disordered breathing at workplaces. <i>Industrial Health</i> , 2005 , 43, 53-7	2.5	16
293	One-year community-based education program for hypercholesterolemia in middle-aged Japanese: a long-term outcome at 8-year follow-up. <i>Atherosclerosis</i> , 2002 , 164, 195-202	3.1	16
292	Trends of cardiovascular risk factors and diseases in Japan: implications for primordial prevention. <i>Preventive Medicine</i> , 1999 , 29, S102-5	4.3	16
291	Frequency of Seaweed Intake and Its Association with Cardiovascular Disease Mortality: The JACC Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 1340-1347	4	15
290	Comparison between the triglycerides standardization of routine methods used in Japan and the chromotropic acid reference measurement procedure used by the CDC Lipid Standardization Programme. <i>Annals of Clinical Biochemistry</i> , 2016 , 53, 632-639	2.2	15
289	LDL cholesterol performance of beta quantification reference measurement procedure. <i>Clinica Chimica Acta</i> , 2014 , 431, 288-93	6.2	15
288	The relationships between interest for and participation in health screening and risk of mortality: the Japan Collaborative Cohort Study. <i>Preventive Medicine</i> , 2005 , 41, 767-71	4.3	15
287	A Prospective Cohort Study of Employment Status and Mortality from Circulatory Disorders among Japanese Workers. <i>Journal of Occupational Health</i> , 2005 , 47, 510-7	2.3	15
286	Work-family conflict and self-rated health among Japanese workers: How household income modifies associations. <i>PLoS ONE</i> , 2017 , 12, e0169903	3.7	15
285	Association of low job control with a decrease in memory (CD4+ CD45RO+) T lymphocytes in Japanese middle-aged male workers in an electric power plant. <i>Industrial Health</i> , 2002 , 40, 142-8	2.5	15
284	Passive smoking and chronic obstructive pulmonary disease mortality: findings from the Japan collaborative cohort study. <i>International Journal of Public Health</i> , 2017 , 62, 489-494	4	14
283	Similarities and differences between coronary heart disease and stroke in the associations with cardiovascular risk factors: The Japan Collaborative Cohort Study. <i>Atherosclerosis</i> , 2017 , 261, 124-130	3.1	14

282	Dietary Inflammatory Index Is Associated with Risk of All-Cause and Cardiovascular Disease Mortality but Not with Cancer Mortality in Middle-Aged and Older Japanese Adults. <i>Journal of Nutrition</i> , 2019 , 149, 1451-1459	4.1	14
281	Oxalates, urinary stones and risk of cardiovascular diseases. <i>Medical Hypotheses</i> , 2020 , 137, 109570	3.8	14
280	A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan. <i>European Journal of Epidemiology</i> , 2017 , 32, 547-557	12.1	14
279	Impact of Caregiver Type for 3-Year-Old Children on Subsequent Between-Meal Eating Habits and Being Overweight From Childhood to Adulthood: A 20-Year Follow-up of the Ibaraki Children Cohort (IBACHIL) Study. <i>Journal of Epidemiology</i> , 2015 , 25, 600-7	3.4	14
278	The reasonable reliability of a self-administered food frequency questionnaire for an urban, Japanese, middle-aged population: the Suita study. <i>Nutrition Research</i> , 2015 , 35, 14-22	4	14
277	Non-fasting blood glucose and risk of incident coronary heart disease in middle-aged general population: the Circulatory Risk in Communities Study (CIRCS). <i>Preventive Medicine</i> , 2012 , 55, 603-7	4.3	14
276	Plasma fibrinogen and its correlates in urban Japanese men. <i>International Journal of Epidemiology</i> , 1996 , 25, 521-7	7.8	14
275	Stratified age-period-cohort analysis of stroke mortality in Japan, 1960 to 2000. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2007 , 16, 91-102	2.8	14
274	Hemostatic variables in Japanese and Caucasian men. Tissue plasminogen activator, antithrombin III, and protein C and their relations to coronary risk factors. <i>American Journal of Epidemiology</i> , 1990 , 132, 41-6	3.8	14
273	Study protocol for the PURSUIT-HFpEF study: a Prospective, Multicenter, Observational Study of Patients with Heart Failure with Preserved Ejection Fraction. <i>BMJ Open</i> , 2020 , 10, e038294	3	14
272	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. <i>Journal of Epidemiology</i> , 2020 , 30, 46-54	3.4	14
271	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 1027-1035	7	13
270	Dietary magnesium intake and risk of incident coronary heart disease in men: A prospective cohort study. <i>Clinical Nutrition</i> , 2018 , 37, 1602-1608	5.9	13
269	Plasma levels of n-3 fatty acids and risk of coronary heart disease among Japanese: The Japan Public Health Center-based (JPHC) study. <i>Atherosclerosis</i> , 2018 , 272, 226-232	3.1	13
268	Occupational physical activity in relation to risk of cardiovascular mortality: The Japan Collaborative Cohort Study for Evaluation for Cancer Risk (JACC Study). <i>Preventive Medicine</i> , 2016 , 89, 286-291	4.3	13
267	Utility of the triglyceride level for predicting incident diabetes mellitus according to the fasting status and body mass index category: the Ibaraki Prefectural Health Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 1152-69	4	13
266	Parental history and lifestyle behaviors in relation to mortality from stroke among Japanese men and women: the Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2012 , 22, 331-9	3.4	13
265	Adiposity and risk of cardiovascular diseases in Japan: secular trend, individual level associations and causal pathway - implications for the prevention of cardiovascular diseases in societies with rapid economic development. <i>EPMA Journal</i> , 2011 , 2, 65-73	8.8	13

264	Alcohol consumption before sleep is associated with severity of sleep-disordered breathing among professional Japanese truck drivers. <i>Alcoholism: Clinical and Experimental Research</i> , 2007 , 31, 2053-8	3.7	13
263	History of blood transfusion before 1990 is a risk factor for stroke and cardiovascular diseases: the Japan collaborative cohort study (JACC study). <i>Cerebrovascular Diseases</i> , 2005 , 20, 164-71	3.2	13
262	Measurement performance of reagent manufacturers by Centers for Disease Control and Prevention/Cholesterol Reference Method Laboratory Network lipid standardization specified for metabolic syndrome-focused health checkups program in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2009 , 16, 756-63	4	13
261	Occupational status and job stress in relation to cardiovascular stress reactivity in Japanese workers. <i>Preventive Medicine Reports</i> , 2016 , 4, 61-7	2.6	13
260	Clinical and immunologic features of recurrent herpes zoster (HZ). <i>Journal of the American Academy of Dermatology</i> , 2016 , 75, 950-956.e1	4.5	13
259	Marital Transition and Risk of Stroke: How Living Arrangement and Employment Status Modify Associations. <i>Stroke</i> , 2016 , 47, 991-8	6.7	12
258	Association between weight change since 20 years of age with mortality from myocardial infarction and chronic heart failure in the Japan collaborative cohort (JACC) study. <i>Circulation Journal</i> , 2014 , 78, 649-55	2.9	12
257	Prospective cohort study on television viewing time and incidence of lung cancer: findings from the Japan Collaborative Cohort Study. <i>Cancer Causes and Control</i> , 2013 , 24, 1547-53	2.8	12
256	Cellular growth factors in relation to mortality from cardiovascular disease in middle-aged Japanese: the JACC study. <i>Atherosclerosis</i> , 2012 , 224, 154-60	3.1	12
255	Validity of brief food frequency questionnaire for estimation of dietary intakes of folate, vitamins B6 and B12, and their associations with plasma homocysteine concentrations. <i>International Journal of Food Sciences and Nutrition</i> , 2010 , 61, 61-7	3.7	12
254	Validity of a self-administered food frequency questionnaire (FFQ) and its generalizability to the estimation of dietary folate intake in Japan. <i>Nutrition Journal</i> , 2005 , 4, 26	4.3	12
253	Sleep-disordered breathing and blood pressure levels among shift and day workers. <i>American Journal of Hypertension</i> , 2006 , 19, 346-51; discussion 352	2.3	12
252	Body Mass Index and Risks of Incident Ischemic Stroke Subtypes: The Japan Public Health Center-Based Prospective (JPHC) Study. <i>Journal of Epidemiology</i> , 2019 , 29, 325-333	3.4	12
251	Total cholesterol performance of Abell-Levy-Brodie-Kendall reference measurement procedure: Certification of Japanese in-vitro diagnostic assay manufacturers through CDC/Cholesterol Reference Method Laboratory Network. <i>Clinica Chimica Acta</i> , 2015 , 445, 127-32	6.2	11
250	Dietary intakes of fat soluble vitamins as predictors of mortality from heart failure in a large prospective cohort study. <i>Nutrition</i> , 2018 , 47, 50-55	4.8	11
249	Diabetes Trend and Impact on Risk of Cardiovascular Disease in Middle-Aged Japanese People - The CIRCS Study. <i>Circulation Journal</i> , 2016 , 80, 2343-2348	2.9	11
248	Association between markers of arterial stiffness and atrial fibrillation in the Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2017 , 263, 244-248	3.1	11
247	Television viewing time and mortality from stroke and coronary artery disease among Japanese men and women -- the Japan Collaborative Cohort Study. <i>Circulation Journal</i> , 2015 , 79, 2389-95	2.9	11

246	Association Between Average Daily Television Viewing Time and Chronic Obstructive Pulmonary Disease-Related Mortality: Findings From the Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2015 , 25, 431-6	3.4	11
245	Association of Body Mass Index and Mortality in Japanese Diabetic Men and Women Based on Self-Reports: The Japan Collaborative Cohort (JACC) Study. <i>Journal of Epidemiology</i> , 2015 , 25, 553-8	3.4	11
244	Saturated fat intake and cardiovascular disease in Japanese population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015 , 22, 435-9	4	11
243	Relationship between the ankle-brachial index and the risk of coronary heart disease and stroke: the circulatory risk in communities study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 1283-9	4	11
242	The Impact of C-Reactive Protein on Risk of Stroke, Stroke Subtypes, and Ischemic Heart Disease in Middle-Aged Japanese: the Japan Public Health Center-Based Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19,	4	11
241	Association between dietary behavior and risk of hypertension among Japanese male workers. <i>Hypertension Research</i> , 2013 , 36, 374-80	4.7	11
240	Cost-effectiveness and budget impact analyses of a long-term hypertension detection and control program for stroke prevention. <i>Journal of Hypertension</i> , 2012 , 30, 1874-9	1.9	11
239	A prospective cohort study of perceived noise exposure at work and cerebrovascular diseases among male workers in Japan. <i>Journal of Occupational Health</i> , 2007 , 49, 382-8	2.3	11
238	Relationship between urinary cGMP excretion and serum total cholesterol levels in a general population. <i>Atherosclerosis</i> , 2005 , 179, 379-86	3.1	11
237	Ankle-arm blood pressure index as a correlate of preclinical carotid atherosclerosis in elderly Japanese men. <i>Atherosclerosis</i> , 2006 , 184, 420-4	3.1	11
236	Relationships of craniofacial morphology and body mass index with sleep-disordered breathing in Japanese men. <i>Laryngoscope</i> , 2004 , 114, 1838-42	3.6	11
235	Polymorphism of the apolipoprotein B gene and blood lipid concentrations in Japanese and Caucasian population samples. <i>Atherosclerosis</i> , 1996 , 126, 233-41	3.1	11
234	Facilitators and barriers of adopting healthy lifestyle in rural China: a qualitative analysis through social capital perspectives. <i>Nagoya Journal of Medical Science</i> , 2016 , 78, 163-73	0.7	11
233	Relationships of soluble E-selectin and high-sensitivity C-reactive protein with carotid atherosclerosis in Japanese men. <i>Journal of Atherosclerosis and Thrombosis</i> , 2009 , 16, 339-45	4	11
232	Cruciferous vegetable intake and mortality in middle-aged adults: A prospective cohort study. <i>Clinical Nutrition</i> , 2019 , 38, 631-643	5.9	11
231	Non-High-Density Lipoprotein Cholesterol and Risk of Stroke Subtypes and Coronary Heart Disease: The Japan Public Health Center-Based Prospective (JPHC) Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 363-374	4	11
230	Alcohol Consumption and Risk of Gastric Cancer: The Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2021 , 31, 30-36	3.4	11
229	Impact of seaweed intake on health. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 877-889	5.2	11

228	Circulating miR-21, miR-29a, and miR-126 are associated with premature death risk due to cancer and cardiovascular disease: the JACC Study. <i>Scientific Reports</i> , 2021 , 11, 5298	4.9	11
227	Yellow Tongue Coating is Associated With Diabetes Mellitus Among Japanese Non-smoking Men and Women: The Toon Health Study. <i>Journal of Epidemiology</i> , 2018 , 28, 287-291	3.4	11
226	Daily Total Physical Activity and Incident Cardiovascular Disease in Japanese Men and Women: Japan Public Health Center-Based Prospective Study. <i>Circulation</i> , 2017 , 135, 1471-1473	16.7	10
225	Among the water-soluble vitamins, dietary intakes of vitamins C, B2 and folate are associated with the reduced risk of diabetes in Japanese women but not men. <i>British Journal of Nutrition</i> , 2019 , 121, 1357-1364	3.6	10
224	Neighborhood contextual factors for smoking among middle-aged Japanese: a multilevel analysis. <i>Health and Place</i> , 2015 , 31, 17-23	4.6	10
223	HDL cholesterol performance using an ultracentrifugation reference measurement procedure and the designated comparison method. <i>Clinica Chimica Acta</i> , 2015 , 439, 185-90	6.2	10
222	Educational Levels and Risk of Suicide in Japan: The Japan Public Health Center Study (JPHC) Cohort I. <i>Journal of Epidemiology</i> , 2016 , 26, 315-21	3.4	10
221	Smoking cessation and COPD mortality among Japanese men and women: the JACC study. <i>Preventive Medicine</i> , 2012 , 55, 639-43	4.3	10
220	Serum Fatty Acid and Risk of Coronary Artery Disease - Circulatory Risk in Communities Study (CIRCS). <i>Circulation Journal</i> , 2018 , 82, 3013-3020	2.9	10
219	Adult height and all-cause and cause-specific mortality in the Japan Public Health Center-based Prospective Study (JPHC). <i>PLoS ONE</i> , 2018 , 13, e0197164	3.7	10
218	Multivitamin use and risk of stroke mortality: the Japan collaborative cohort study. <i>Stroke</i> , 2015 , 46, 1167-72	6.7	9
217	Habitual tub bathing and risks of incident coronary heart disease and stroke. <i>Heart</i> , 2020 , 106, 732-737	5.1	9
216	Serum creatinine levels and risk of incident type 2 diabetes mellitus or dysglycemia in middle-aged Japanese men: a retrospective cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2018 ,	4.5	9
215	Associations of Perceived Mental Stress, Sense of Purpose in Life, and Negative Life Events With the Risk of Incident Herpes Zoster and Postherpetic Neuralgia: The SHEZ Study. <i>American Journal of Epidemiology</i> , 2018 , 187, 251-259	3.8	9
214	Descriptive epidemiology of diabetes prevalence and HbA1c distributions based on a self-reported questionnaire and a health checkup in the JPHC diabetes study. <i>Journal of Epidemiology</i> , 2014 , 24, 460-8 ^{3.4}	3.4	9
213	Employment situation and risk of death among middle-aged Japanese women. <i>Journal of Epidemiology and Community Health</i> , 2015 , 69, 1012-7	5.1	9
212	C-reactive protein levels and risk of disabling dementia with and without stroke in Japanese: the Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2014 , 236, 438-43	3.1	9
211	An association between central aortic pressure and subclinical organ damage of the heart among a general Japanese cohort: Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2014 , 232, 94-8 ^{3.1}	3.1	9

210	Body mass index and weight change during adulthood are associated with increased mortality from liver cancer: the JACC Study. <i>Journal of Epidemiology</i> , 2013 , 23, 219-26	3.4	9
209	Long-term exposure to elevated blood pressure and mortality from cardiovascular disease in a Japanese population: the Ibaraki Prefectural Health Study. <i>Hypertension Research</i> , 2011 , 34, 139-44	4.7	9
208	Sleep duration and risk of breast cancer: The JACC Study. <i>Breast Cancer Research and Treatment</i> , 2019 , 174, 219-225	4.4	9
207	Tobacco Control Measures to Reduce Socioeconomic Inequality in Smoking: The Necessity, Time-Course Perspective, and Future Implications. <i>Journal of Epidemiology</i> , 2018 , 28, 170-175	3.4	9
206	Associations of disaster-related and psychosocial factors with changes in smoking status after a disaster: a cross-sectional survey after the Great East Japan Earthquake. <i>BMJ Open</i> , 2018 , 8, e018943	3	9
205	Alcohol consumption and risks of hypertension and cardiovascular disease in Japanese men and women. <i>Hypertension Research</i> , 2020 , 43, 477-481	4.7	8
204	Retinal Vascular Changes and Prospective Risk of Disabling Dementia: the Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2017 , 24, 687-695	4	8
203	Association between average daily television viewing time and the incidence of ovarian cancer: findings from the Japan Collaborative Cohort Study. <i>Cancer Causes and Control</i> , 2018 , 29, 213-219	2.8	8
202	Influence of municipal- and individual-level socioeconomic conditions on mortality in Japan. <i>International Journal of Behavioral Medicine</i> , 2014 , 21, 737-49	2.6	8
201	Menopausal status in relation to cardiovascular stress reactivity in healthy Japanese participants. <i>Psychosomatic Medicine</i> , 2014 , 76, 701-8	3.7	8
200	Promoting predictive, preventive and personalized medicine in treatment of cardiovascular diseases. <i>EPMA Journal</i> , 2011 , 2, 1-4	8.8	8
199	Prediction of myocardial infarction using coronary risk scores among Japanese male workers: 3M Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010 , 17, 452-9	4	8
198	High sodium intake strengthens the association of ACE I/D polymorphism with blood pressure in a community. <i>American Journal of Hypertension</i> , 2007 , 20, 751-7	2.3	8
197	Determinants of platelet aggregation in 50-70-year-old men from three Japanese communities. <i>Atherosclerosis</i> , 2002 , 165, 327-34	3.1	8
196	Use of aspirin for prevention of cardiovascular disease--1981-82 to 1985-86: the Minnesota Heart Survey. <i>American Heart Journal</i> , 1988 , 116, 827-30	4.9	8
195	Profile of non-communicable disease risk factors among adults in the Republic of Palau: findings of a national STEPS survey. <i>Nagoya Journal of Medical Science</i> , 2015 , 77, 609-19	0.7	8
194	Prenatal tobacco smoking is associated with postpartum depression in Japanese pregnant women: The Japan environment and children's study. <i>Journal of Affective Disorders</i> , 2020 , 264, 76-81	6.6	8
193	Trends in the proportions of stroke subtypes and coronary heart disease in the Japanese men and women from 1995 to 2009. <i>Atherosclerosis</i> , 2016 , 248, 219-23	3.1	8

192	Long-Term Prognosis of Brugada-Type ECG and ECG With Atypical ST-Segment Elevation in the Right Precordial Leads Over 20 Years: Results From the Circulatory Risk in Communities Study (CIRCS). <i>Journal of the American Heart Association</i> , 2016 , 5,	6	8
191	No modifying effect of education level on the association between lifestyle behaviors and cardiovascular mortality: the Japan Collaborative Cohort Study. <i>Scientific Reports</i> , 2017 , 7, 39820	4.9	7
190	Alcohol consumption and mortality from aortic disease among Japanese men: The Japan Collaborative Cohort study. <i>Atherosclerosis</i> , 2017 , 266, 64-68	3.1	7
189	Sleep Shortage Is Associated With Postherpetic Neuralgia Development Through Hyperesthesia and Acute Pain Intensity: A Community-Based Prospective Cohort Study. <i>Pain Practice</i> , 2019 , 19, 476-483 ³		7
188	Variations in the estimated intake of acrylamide from food in the Japanese population. <i>Nutrition Journal</i> , 2020 , 19, 17	4.3	7
187	The relationship between sodium concentrations in spot urine and blood pressure increases: a prospective study of Japanese general population: the Circulatory Risk in Communities Study (CIRCS). <i>BMC Cardiovascular Disorders</i> , 2016 , 16, 55	2.3	7
186	Water intake from foods and beverages and risk of mortality from CVD: the Japan Collaborative Cohort (JACC) Study. <i>Public Health Nutrition</i> , 2018 , 21, 3011-3017	3.3	7
185	Cohort profile: the Fangshan Cohort Study of cardiovascular epidemiology in Beijing, China. <i>Journal of Epidemiology</i> , 2014 , 24, 84-93	3.4	7
184	Relationship between cell-mediated immunity to Varicella-Zoster virus and aging in subjects from the community-based Shozu Herpes Zoster study. <i>Journal of Medical Virology</i> , 2017 , 89, 313-317	19.7	7
183	Profile of Non-communicable Disease Risk Factors Among Young People in Palau. <i>Journal of Epidemiology</i> , 2015 , 25, 392-7	3.4	7
182	Is the association between healthy lifestyle behaviors and cardiovascular mortality modified by overweight status? The Japan Collaborative Cohort Study. <i>Preventive Medicine</i> , 2014 , 62, 142-7	4.3	7
181	Ten-year evaluation of homogeneous low-density lipoprotein cholesterol methods developed by Japanese manufacturers. Application of the Centers for Disease Control and Prevention/Cholesterol Reference Method Laboratory Network lipid standardization protocol. <i>Journal of Clinical Lipidology</i> , 2010 , 4, 127-31	4	7
180	Projecting the probability of survival free from cancer and cardiovascular incidence through lifestyle modification in Japan. <i>Preventive Medicine</i> , 2009 , 48, 128-33	4.3	7
179	Lipoprotein(a) and its correlates in Japanese and U.S. population samples. <i>Annals of Epidemiology</i> , 1996 , 6, 324-30	6.4	7
178	Prevalence and predictors of prediabetes and diabetes among adults in Palau: population-based national STEPS survey. <i>Nagoya Journal of Medical Science</i> , 2016 , 78, 475-483	0.7	7
177	Impact of intention and feeling toward being pregnant on postpartum depression: the Japan Environment and Children's Study (JECS). <i>Archives of Women's Mental Health</i> , 2020 , 23, 131-137	5	7
176	Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 954-968	5.2	7
175	Association of cigarette smoking with radial augmentation index: the Circulatory Risk in Communities Study (CIRCS). <i>Hypertension Research</i> , 2018 , 41, 1054-1062	4.7	7

174	Dietary Antioxidant Micronutrients and All-Cause Mortality: The Japan Collaborative Cohort Study for Evaluation of Cancer Risk. <i>Journal of Epidemiology</i> , 2018 , 28, 388-396	3.4	7
173	Recurrent Pregnancy Loss and Cardiovascular Disease Mortality in Japanese Women: A Population-Based, Prospective Cohort Study. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017 , 26, 1047-1054	2.8	6
172	Maternal total energy, macronutrient and vitamin intakes during pregnancy associated with the offspring's birth size in the Japan Environment and Children's Study. <i>British Journal of Nutrition</i> , 2020 , 124, 558-566	3.6	6
171	Fifty-year Time Trends in Blood Pressures, Body Mass Index and their Relations in a Japanese Community: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2017 , 24, 518-529	4	6
170	Associations between alcohol consumption and sleep-disordered breathing among Japanese women. <i>Respiratory Medicine</i> , 2011 , 105, 796-800	4.6	6
169	Effectiveness of the combined approach for assessing social gradients in stroke risk among married women in Japan. <i>Journal of Epidemiology</i> , 2012 , 22, 324-30	3.4	6
168	Urinary cyclic GMP excretion and blood pressure levels in a general population. <i>Atherosclerosis</i> , 2004 , 172, 161-6	3.1	6
167	A case-reference study on plasma fibrinogen concentrations and coronary atherosclerosis in Japanese. <i>Journal of Epidemiology</i> , 1996 , 6, 81-6	3.4	6
166	Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population. <i>PLoS ONE</i> , 2020 , 15, e0244007	3.7	6
165	Risk stratification based on metabolic syndrome as well as non- metabolic risk factors in the assessment of carotid atherosclerosis. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011 , 18, 504-12	4	6
164	Relationship of urinary cGMP excretion with aging and menopausal status in a general population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2009 , 16, 457-62	4	6
163	Relationship between unhealthy sleep status and dry eye symptoms in a Japanese population: The JPHC-NEXT study. <i>Ocular Surface</i> , 2021 , 21, 306-312	6.5	6
162	Dehydroepiandrosterone-sulfate is associated with cardiovascular reactivity to stress in women. <i>Psychoneuroendocrinology</i> , 2016 , 69, 116-22	5	6
161	Blood pressure levels and risk of cardiovascular disease mortality among Japanese men and women: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC Study). <i>Journal of Hypertension</i> , 2019 , 37, 1366-1371	1.9	6
160	Fat-soluble vitamins from diet in relation to risk of type 2 diabetes mellitus in Japanese population. <i>British Journal of Nutrition</i> , 2019 , 121, 647-653	3.6	6
159	Associations of Daily Walking Time With Pneumonia Mortality Among Elderly Individuals With or Without a Medical History of Myocardial Infarction or Stroke: Findings From the Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2019 , 29, 233-237	3.4	6
158	Nonfasting Glucose and Incident Stroke and Its Types - The Circulatory Risk in Communities Study (CIRCS). <i>Circulation Journal</i> , 2018 , 82, 1598-1604	2.9	6
157	Relationships of diabetes and hyperglycaemia with intraocular pressure in a Japanese population: the JPHC-NEXT Eye Study. <i>Scientific Reports</i> , 2020 , 10, 5355	4.9	5

156	"Ikigai", Subjective Wellbeing, as a Modifier of the Parity-Cardiovascular Mortality Association - The Japan Collaborative Cohort Study. <i>Circulation Journal</i> , 2018 , 82, 1302-1308	2.9	5
155	Fish Intake and Death From Pulmonary Embolisms Among Japanese Men and Women - The Japan Collaborative Cohort (JACC) Study. <i>Circulation Journal</i> , 2018 , 82, 2063-2070	2.9	5
154	Fish intake and risk of mortality due to aortic dissection and aneurysm: A pooled analysis of the Japan cohort consortium. <i>Clinical Nutrition</i> , 2019 , 38, 1678-1683	5.9	5
153	Duration of cigarette smoking is a risk factor for oropharyngeal cancer mortality among Japanese men and women: the Ibaraki Prefectural Health Study (IPHS). <i>Annals of Epidemiology</i> , 2013 , 23, 546-50	6.4	5
152	Social epidemiology and Eastern Wisdom. <i>Journal of Epidemiology</i> , 2012 , 22, 291-4	3.4	5
151	A prospective cohort study of neighborhood stress and ischemic heart disease in Japan: a multilevel analysis using the JACC study data. <i>BMC Public Health</i> , 2011 , 11, 398	4.1	5
150	Alcohol consumption, alcohol-induced flushing and incidence of acute myocardial infarction among middle-aged men in Japan—Japan Public Health Center-based prospective study. <i>Atherosclerosis</i> , 2007 , 194, 512-6	3.1	5
149	Television Viewing Time and Breast Cancer Incidence for Japanese Premenopausal and Postmenopausal Women: The JACC Study. <i>Cancer Research and Treatment</i> , 2019 , 51, 1509-1517	5.2	5
148	Television viewing time, walking time, and risk of type 2 diabetes in Japanese men and women: The Japan Collaborative Cohort Study. <i>Preventive Medicine</i> , 2019 , 118, 220-225	4.3	5
147	Weight Change and Mortality from Cardiovascular Diseases: The Japan Collaborative Cohort Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 25-33	4	5
146	Low BMI and weight loss aggravate COPD mortality in men, findings from a large prospective cohort: the JACC study. <i>Scientific Reports</i> , 2021 , 11, 1531	4.9	5
145	Changes in ischaemic ECG abnormalities and subsequent risk of cardiovascular disease. <i>Heart Asia</i> , 2017 , 9, 36-43	1.9	4
144	Blood soluble Fas levels and mortality from cardiovascular disease in middle-aged Japanese: The JACC study. <i>Atherosclerosis</i> , 2017 , 260, 97-101	3.1	4
143	Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. <i>Journal of Epidemiology</i> , 2017 , 27, 331-337	3.4	4
142	Joint impact of muscle mass and waist circumference on type 2 diabetes in Japanese middle-aged adults: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Diabetes</i> , 2020 , 12, 677-685	3.8	4
141	Health Disparities in Ischaemic Heart Disease Mortality in Hungary From 1970 to 2010: An Age-Period-Cohort Analysis. <i>Journal of Epidemiology</i> , 2015 , 25, 399-406	3.4	4
140	Dietary Reference Intakes for Japanese 2010: Fat. <i>Journal of Nutritional Science and Vitaminology</i> , 2012 , 59, S44-S52	1.1	4
139	Reproducibility of 24-hour dietary recall for vitamin intakes by middle-aged Japanese men and women. <i>Journal of Nutrition, Health and Aging</i> , 2010 , 14, 196-200	5.2	4

138	Trends of Blood Pressure Distributions in a Northeast Rural Japanese Population. <i>Journal of Epidemiology</i> , 1993 , 3, 63-70	3.4	4
137	Association between dietary inflammatory index and serum C-reactive protein concentrations in the Japan Collaborative Cohort Study. <i>Nagoya Journal of Medical Science</i> , 2020 , 82, 237-249	0.7	4
136	Japan Trial in High-Risk Individuals to Enhance Their Referral to Physicians (J-HARP)-A Nurse-Led, Community-Based Prevention Program of Lifestyle-Related Disease. <i>Journal of Epidemiology</i> , 2020 , 30, 194-199	3.4	4
135	Chlamydia pneumoniae infection was associated with risk of mortality from coronary heart disease in Japanese women but not men: the JACC Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2010 , 17, 510-6	4	4
134	Effects of brief self-exercise education on the management of chronic low back pain: A community-based, randomized, parallel-group pragmatic trial. <i>Modern Rheumatology</i> , 2021 , 31, 890-898	3.3	4
133	Weight change during middle age and risk of stroke and coronary heart disease: The Japan Public Health Center-based Prospective Study. <i>Atherosclerosis</i> , 2021 , 322, 67-73	3.1	4
132	Association of Vegetable, Fruit, and Okinawan Vegetable Consumption With Incident Stroke and Coronary Heart Disease. <i>Journal of Epidemiology</i> , 2020 , 30, 37-45	3.4	4
131	Modifiable Risk Factors for Cardiovascular Disease in Korea and Japan. <i>Korean Circulation Journal</i> , 2021 , 51, 643-655	2.2	4
130	Cardiovascular disease, a major global burden: Epidemiology of stroke and ischemic heart disease in Japan.. <i>Global Health & Medicine</i> , 2021 , 3, 358-364	2.4	4
129	Estimation of 10-Year Risk of Death from Coronary Heart Disease, Stroke, and Cardiovascular Disease in a Pooled Analysis of Japanese Cohorts: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 816-825	4	4
128	Green tea consumption and risk of hematologic neoplasms: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC Study). <i>Cancer Causes and Control</i> , 2019 , 30, 1223-1230	2.8	3
127	Association of measles and mumps with cardiovascular disease: The Japan Collaborative Cohort (JACC) study. <i>Atherosclerosis</i> , 2015 , 241, 682-6	3.1	3
126	Relationship between sleep duration and cause-specific mortality in diabetic men and women based on self-reports. <i>Sleep and Biological Rhythms</i> , 2015 , 13, 85-93	1.3	3
125	Impact of hypertension stratified by diabetes on the lifetime risk of cardiovascular disease mortality in Japan: a pooled analysis of data from the Evidence for Cardiovascular Prevention from Observational Cohorts in Japan study. <i>Hypertension Research</i> , 2020 , 43, 1437-1444	4.7	3
124	Message from the New Editor-in-Chief. <i>Journal of Epidemiology</i> , 2011 , 21, 1	3.4	3
123	Aldosterone synthase gene T-344C polymorphism, sodium and blood pressure in a free-living population: a community-based study. <i>Hypertension Research</i> , 2007 , 30, 497-502	4.7	3
122	Possible effects of diets on serum lipids, fatty acids and blood pressure levels in male and female Japanese university students. <i>Environmental Health and Preventive Medicine</i> , 2005 , 10, 42-7	4.2	3
121	Trends in the Incidence of Cardiovascular Diseases and their Risk Factors in a Rural Japanese Population. <i>Journal of Epidemiology</i> , 1992 , 2, 123-135	3.4	3

120	Body mass index and serum gamma-glutamyltransferase level as risk factors for injuries related to professional horse racing: a prospective study. <i>Journal of Occupational Health</i> , 2009 , 51, 323-31	2-3	3
119	Night Work, Rotating Shift Work, and the Risk of Cancer in Japanese Men and Women: The JACC Study. <i>Journal of Epidemiology</i> , 2021 , 31, 585-592	3-4	3
118	Food frequency questionnaire reproducibility for middle-aged and elderly Japanese. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2019 , 28, 362-370	1	3
117	Factors Associated With Family Member Spanking of 3.5-year-old Children in Japan. <i>Journal of Epidemiology</i> , 2020 , 30, 464-473	3-4	3
116	Associations of Carotid Intima-Media Thickness and Plaque Heterogeneity With the Risks of Stroke Subtypes and Coronary Artery Disease in the Japanese General Population: The Circulatory Risk in Communities Study. <i>Journal of the American Heart Association</i> , 2020 , 9, e017020	6	3
115	Television Viewing Time and the Risk of Colorectal Cancer Mortality among Japanese Population: The JACC Study. <i>Cancer Research and Treatment</i> , 2021 , 53, 497-505	5-2	3
114	High Myopia and Its Associated Factors in JPHC-NEXT Eye Study: A Cross-Sectional Observational Study. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5-1	3
113	Effects of Low-Dose Therapist-Led Self-Exercise Education on the Management of Chronic Low Back Pain: Protocol for a Community-Based, Randomized, 6-Month Parallel-Group Study. <i>Spine Surgery and Related Research</i> , 2019 , 3, 377-384	1-7	3
112	The Association Between Habitual Sleep Duration and Mortality According to Sex and Age: The Japan Public Health Center-based Prospective Study. <i>Journal of Epidemiology</i> , 2021 , 31, 109-118	3-4	3
111	Longitudinal Trends in Blood Pressure Associated With the Frequency of Laughter: The Circulatory Risk in Communities Study (CIRCS), a Longitudinal Study of the Japanese General Population. <i>Journal of Epidemiology</i> , 2021 , 31, 125-131	3-4	3
110	Manganese intake from foods and beverages is associated with a reduced risk of type 2 diabetes. <i>Maturitas</i> , 2021 , 143, 127-131	5	3
109	Serum High-Sensitivity C-Reactive Protein Levels and the Risk of Atrial Fibrillation in Japanese Population: the Circulatory Risk in Communities Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 194-202	4	3
108	Impact and attribute of each obesity-related cardiovascular risk factor in combination with abdominal obesity on total health expenditures in adult Japanese National Health insurance beneficiaries: The Ibaraki Prefectural health study. <i>Journal of Epidemiology</i> , 2017 , 27, 354-359	3-4	2
107	Descriptive Epidemiology of Hypertension and Its Association With Obesity: Based on the WHO STEPwise Approach to Surveillance in Palau. <i>Asia-Pacific Journal of Public Health</i> , 2017 , 29, 278-287	2	2
106	Validity and reliability of a self-administered food frequency questionnaire for the JPHC study: The assessment of amino acid intake. <i>Journal of Epidemiology</i> , 2017 , 27, 242-247	3-4	2
105	Cardiovascular reactivity to acute stress associated with sickness absence among Japanese men and women: A prospective study. <i>Brain and Behavior</i> , 2020 , 10, e01541	3-4	2
104	Diabetes and Mortality From Respiratory Diseases: The Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2020 , 30, 457-463	3-4	2
103	Time course of metabolic status in pregnant women: The Japan Environment and Children Study. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1318-1325	3-9	2

102	The association of conventionally medicated systolic and diastolic blood pressure level and mortality from cardiovascular disease: is the lower the better in high stroke population?. <i>Clinical Research in Cardiology</i> , 2020 , 109, 944-948	6.1	2
101	Changes in the living arrangement and risk of stroke in Japan; does it matter who lives in the household? Who among the family matters?. <i>PLoS ONE</i> , 2017 , 12, e0173860	3.7	2
100	Dietary Intake of Energy and Nutrients from Breakfast and Risk of Stroke in The Japanese Population: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 145-153	4	2
99	Interest in health screening as a predictor of long-term overall mortality: multilevel analysis of a Japanese national cohort study. <i>Preventive Medicine</i> , 2011 , 52, 78-83	4.3	2
98	Reply to Dr. Rouhani et al.. <i>Journal of Nutrition</i> , 2011 , 141, 1919-1919	4.1	2
97	Prehospital score for acute disease: a community-based observational study in Japan. <i>BMC Emergency Medicine</i> , 2007 , 7, 17	2.4	2
96	Trends for Cardiovascular Risk Factors and Diseases in Japan. <i>Journal of Epidemiology</i> , 1996 , 6, 183-188	3.4	2
95	Soy Intake and Risk of Type 2 Diabetes Among Japanese Men and Women: JACC Study.. <i>Frontiers in Nutrition</i> , 2021 , 8, 813742	6.2	2
94	Associations of Body Mass Index, Weight Change, Physical Activity, and Sedentary Behavior With Endometrial Cancer Risk Among Japanese Women: The Japan Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2021 , 31, 621-627	3.4	2
93	Effects of Local Cooperation between Healthcare Facilities for Infection Control after Introduction of Additional Charge for Infection Control. <i>Japanese Journal of Environmental Infections</i> , 2016 , 31, 24-31	0.1	2
92	Effects of Intake of Fish Rich in n-3 Polyunsaturated Fatty Acid on Serum Lipids, Serum Fatty Acid Component and Hemostatic Factors.. <i>Nihon Eiyōshokuryō Gakkai Shi = Nippon Eiyōshokuryō Gakkaishi = Journal of Japanese Society of Nutrition and Food Science</i> , 2000 , 53, 1-9	0.2	2
91	Usefulness of Skinfold Thickness Measurements for Determining Body Fat Distribution and Disease Risk for Japanese Men and Women 2012 , 2667-2678		2
90	Relationships Between Reproductive History and Mortality From Cardiovascular Diseases Among Japanese Women: The Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. <i>Journal of Epidemiology</i> , 2020 , 30, 509-515	3.4	2
89	Association of job category and occupational activity with breast cancer incidence in Japanese female workers: the JACC study. <i>BMC Public Health</i> , 2020 , 20, 1106	4.1	2
88	Milk Intake and Stroke Mortality in the Japan Collaborative Cohort Study-A Bayesian Survival Analysis. <i>Nutrients</i> , 2020 , 12,	6.7	2
87	Green Tea and Coffee Consumption and All-Cause Mortality Among Persons With and Without Stroke or Myocardial Infarction. <i>Stroke</i> , 2021 , 52, 957-965	6.7	2
86	Myopia, corneal endothelial cell density and morphology in a Japanese population-based cross-sectional study: the JPHC-NEXT Eye Study. <i>Scientific Reports</i> , 2021 , 11, 6366	4.9	2
85	Relationship between Endothelial Dysfunction and Prevalence of Chronic Kidney Disease: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 622-629	4	2

84	Serum anti-DIDO1, anti-CPSF2, and anti-FOXJ2 antibodies as predictive risk markers for acute ischemic stroke. <i>BMC Medicine</i> , 2021 , 19, 131	11.4	2
83	Smoking cessation, weight gain and risk of cardiovascular disease. <i>Heart</i> , 2021 ,	5.1	2
82	Cross-Sectional Association Between Employment Status and Self-Rated Health Among Middle-Aged Japanese Women: The Influence of Socioeconomic Conditions and Work-Life Conflict. <i>Journal of Epidemiology</i> , 2020 , 30, 396-403	3.4	2
81	Social support, social cohesion and pain during pregnancy: The Japan Environment and Children's Study. <i>European Journal of Pain</i> , 2021 , 25, 872-885	3.7	2
80	Association between mode of delivery and postpartum depression: The Japan Environment and Children's Study (JECS): a prospective cohort study. <i>Journal of Epidemiology</i> , 2021 ,	3.4	2
79	Evaluating the Effect on Mortality of a No-Tranexamic acid (TXA) Policy for Cardiovascular Surgery. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> , 2018 , 32, 1627-1634	2.1	2
78	Association of serum levels of antibodies against ALDOA and FH4 with transient ischemic attack and cerebral infarction. <i>BMC Neurology</i> , 2021 , 21, 274	3.1	2
77	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. <i>Preventive Medicine</i> , 2021 , 148, 106561	4.3	2
76	Right bundle branch block and risk of cardiovascular mortality: the Ibaraki Prefectural Health Study. <i>Heart and Vessels</i> , 2021 , 1	2.1	2
75	Can Blood Pressure in the Elderly be Reduced? Findings From a Long-term Population Survey in Japan. <i>The American Journal of Geriatric Cardiology</i> , 1994 , 3, 42-50		2
74	Impact of modifiable healthy lifestyle adoption on lifetime gain from middle to older age.. <i>Age and Ageing</i> , 2022 , 51,	3	2
73	Self-Reported Eczema in Relation with Mortality from Cardiovascular Disease in Japanese: the Japan Collaborative Cohort Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019 , 26, 775-782	4	1
72	A method for measuring glycerol-blanked triglyceride concentrations by using gas chromatography-isotope dilution mass spectrometry. <i>Annals of Clinical Biochemistry</i> , 2020 , 57, 253-261	2.2	1
71	Urinary 8-Hydroxy-2-Deoxyguanosine Levels and Cardiovascular Disease Incidence in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 1086-1096	4	1
70	Relationship between nerve fiber layer defect and the presence of epiretinal membrane in a Japanese population: The JPHC-NEXT Eye Study. <i>Scientific Reports</i> , 2020 , 10, 779	4.9	1
69	Moderate Levels of N-Terminal Pro-B-Type Natriuretic Peptide is Associated with Increased Risks of Total and Ischemic Strokes among Japanese: The Circulatory Risk in Communities Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 751-760	4	1
68	Differences in dietary habits, serum fatty acid compositions and other coronary risk characteristics between freshmen and fourth-year male university students. <i>Environmental Health and Preventive Medicine</i> , 2001 , 6, 143-8	4.2	1
67	Dietary fiber intake and risk of incident disabling dementia: the Circulatory Risk in Communities Study.. <i>Nutritional Neuroscience</i> , 2022 , 1-8	3.6	1

66	Impact of Body Mass Index on Obesity-Related Cancer and Cardiovascular Disease Mortality; The Japan Collaborative Cohort Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 ,	4	1
65	Overweight and Hypertension in Relation to Chronic Musculoskeletal Pain Among Community-Dwelling Adults: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Epidemiology</i> , 2021 , 31, 566-572	3.4	1
64	Green Tea, Coffee, and Diabetes. <i>Annals of Internal Medicine</i> , 2006 , 145, 634	8	1
63	Breastfeeding in Infancy in Relation to Subsequent Physical Size: A 20-year Follow-up of the Ibaraki Children Cohort Study (IBACHIL). <i>Journal of Epidemiology</i> , 2021 ,	3.4	1
62	Engaging in musical activities and the risk of dementia in older adults: A longitudinal study from the Japan gerontological evaluation study. <i>Geriatrics and Gerontology International</i> , 2021 , 21, 451-457	2.9	1
61	Body Mass Index and Mortality From Aortic Aneurysm and Dissection. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021 , 28, 338-348	4	1
60	Serum anti-AP3D1 antibodies are risk factors for acute ischemic stroke related with atherosclerosis. <i>Scientific Reports</i> , 2021 , 11, 13450	4.9	1
59	Dietary Patterns and Risk of Esophageal Cancer Mortality: The Japan Collaborative Cohort Study. <i>Nutrition and Cancer</i> , 2016 , 68, 1001-9	2.8	1
58	Response by Shirakawa and Iso to Letter Regarding Article, "Watching Television and Risk of Mortality From Pulmonary Embolism Among Japanese Men and Women: The JACC Study (Japan Collaborative Cohort)". <i>Circulation</i> , 2016 , 134, e501-e502	16.7	1
57	Prevalence and Correlates of Dyslipidemia Among Men and Women in Palau: Findings of the Palau STEPS Survey 2011-2013. <i>Journal of Epidemiology</i> , 2019 , 29, 97-103	3.4	1
56	The Prospective Association Between Plasma Concentrations of Cellular Growth Factors and Risk of Heart Failure Mortality in Japanese Population. <i>Journal of Epidemiology</i> , 2019 , 29, 104-109	3.4	1
55	Glycated haemoglobin (HbA1c) as a reliable option for detecting diabetes among the urban poor population in Bangladesh. <i>European Journal of Public Health</i> , 2020 , 30, 839-841	2.1	1
54	Intake of fish and long-chain n-3 polyunsaturated fatty acids and risk of diseases in a Japanese population: a narrative review. <i>European Journal of Clinical Nutrition</i> , 2021 , 75, 902-920	5.2	1
53	Tobacco Price Increase and Successful Smoking Cessation for Two or More Years in Japan. <i>Nicotine and Tobacco Research</i> , 2021 , 23, 716-723	4.9	1
52	Risk of Herpes Zoster in Relation to Body Mass Index Among Residents Aged ≥50 Years: The Shozu Herpes Zoster Study. <i>Journal of Epidemiology</i> , 2021 ,	3.4	1
51	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. <i>Journal of Cancer Survivorship</i> , 2021 , 15, 912-921	5.1	1
50	Cardiovascular disease mortality in relation to physical activity during adolescence and adulthood in Japan: Does school-based sport club participation matter?. <i>Preventive Medicine</i> , 2018 , 113, 102-108	4.3	1
49	Association of maternal total cholesterol with SGA or LGA birth at term: The Japan Environment and Children Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 ,	5.6	1

48	Dietary intake of tocopherols and risk of incident disabling dementia. <i>Scientific Reports</i> , 2021 , 11, 16429	4.9	1
47	Associations Between Occupational Status, Support at Work, and Salivary Cortisol Levels. <i>International Journal of Behavioral Medicine</i> , 2021 , 1	2.6	1
46	Association of dietary intakes of vitamin B12, vitamin B6, folate, and methionine with the risk of esophageal cancer: the Japan Public Health Center-based (JPHC) prospective study. <i>BMC Cancer</i> , 2021 , 21, 982	4.8	1
45	Peanut Consumption and Risk of Stroke and Ischemic Heart Disease in Japanese Men and Women: The JPHC Study. <i>Stroke</i> , 2021 , 52, 3543-3550	6.7	1
44	Non-Fatal Injury in Thailand From 2005 to 2013: Incidence Trends and Links to Alcohol Consumption Patterns in the Thai Cohort Study. <i>Journal of Epidemiology</i> , 2016 , 26, 471-80	3.4	0
43	Overview of the Role of Antioxidant Vitamins as Protection Against Cardiovascular Disease 2014 , 213-224		0
42	Letter by Noda and Iso regarding article, "Low-density lipoprotein cholesterol concentrations and death due to intraparenchymal hemorrhage: the Ibaraki Prefectural Health Study". <i>Circulation</i> , 2009 , 120, e280; author reply e281	16.7	0
41	The reproducibility of 24-h dietary recall for estimating mineral intakes and their food sources among middle-aged Japanese men and women. <i>International Journal of Food Sciences and Nutrition</i> , 2009 , 60 Suppl 7, 30-40	3.7	0
40	Factors associated with prolonged duration of viral clearance in non-severe SARS-CoV-2 patients in Osaka, Japan. <i>Environmental Health and Preventive Medicine</i> , 2021 , 26, 115	4.2	0
39	Job stress factors measured by Brief Job Stress Questionnaire and sickness absence among Japanese workers: A longitudinal study. <i>Fukushima Journal of Medical Sciences</i> , 2020 , 66, 88-96	0.9	0
38	Urinary Stones and Risk of Coronary Heart Disease and Stroke: the Japan Public Health Center-Based Prospective Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020 , 27, 1208-1215	4	0
37	Dairy intake and the risk of pancreatic cancer: the Japan Collaborative Cohort Study (JACC Study) and meta-analysis of prospective cohort studies. <i>British Journal of Nutrition</i> , 2021 , 1-9	3.6	0
36	Prediction of Lifetime Risk of Cardiovascular Disease Deaths Stratified by Sex in the Japanese Population. <i>Journal of the American Heart Association</i> , 2021 , 10, e021753	6	0
35	Risk Classification for Metabolic Syndrome and the Incidence of Cardiovascular Disease in Japan With Low Prevalence of Obesity: A Pooled Analysis of 10 Prospective Cohort Studies. <i>Journal of the American Heart Association</i> , 2021 , 10, e020760	6	0
34	Anger Expression and the Risk of Cardiovascular Disease Among Urban and Rural Japanese Residents: The Circulatory Risk in Communities Study. <i>Psychosomatic Medicine</i> , 2020 , 82, 215-223	3.7	0
33	Relation Between Body Mass Index and Dry Eye Disease: The Japan Public Health Center-Based Prospective Study for the Next Generation. <i>Eye and Contact Lens</i> , 2021 , 47, 449-455	3.2	0
32	Successful Reduction of Blood Pressure and Stroke Risk in Japan 2019 , 47-55		0
31	Risk of postpartum depression and very early child mistreatment among mothers reporting higher autistic traits: Evidence from the Japan Environment and Children's Study. <i>Journal of Affective Disorders</i> , 2021 , 280, 11-16	6.6	0

12	Depressive Tendency and the Risk of Death from Pneumonia: The JACC Study. <i>Internal Medicine</i> , 2020 , 59, 3123-3130	1.1
11	Pedometer Measured Walking Habits and Serum HDL-cholesterol Level. <i>The Journal of Japan Atherosclerosis Society</i> , 1993 , 21, 585-589	
10	Lifestyles and biomarkers as risk and beneficial factors of stroke in the Circulatory Risk in Communities Study. <i>Nosotchu</i> , 2015 , 37, 367-373	0.1
9	Association of tea consumption and the risk of gastric cancer in Japanese adults: the Japan Collaborative Cohort Study. <i>BMJ Open</i> , 2020 , 10, e038243	3
8	Response to eLetter. <i>Heart</i> , 2020 , 106, 1365-1366	5.1
7	Reader response: Low-density lipoprotein cholesterol and risk of intracerebral hemorrhage: A prospective study. <i>Neurology</i> , 2020 , 95, 98-99	6.5
6	Salt taste perception and blood pressure levels in population-based samples: the Circulatory Risk in Communities Study (CIRCS). <i>British Journal of Nutrition</i> , 2021 , 125, 203-211	3.6
5	Non-alcoholic beverages intake and risk of CVD among Japanese men and women: the Japan Public Health Center study. <i>British Journal of Nutrition</i> , 2021 , 1-8	3.6
4	The apparent inverse association between dietary carotene intake and risk of cardiovascular mortality disappeared after adjustment for other cardioprotective dietary intakes: The Japan collaborative cohort study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 3064-3075	4.5
3	The number of leisure-time activities and risk of functional disability among Japanese older population: the JAGES cohort.. <i>Preventive Medicine Reports</i> , 2022 , 26, 101741	2.6
2	Effects of a laughter program on body weight and mental health among Japanese people with metabolic syndrome risk factors: a randomized controlled trial.. <i>BMC Geriatrics</i> , 2022 , 22, 361	4.1
1	The association of family history of herpes zoster and the risk of incident herpes zoster: the SHEZ Study. <i>Environmental Health and Preventive Medicine</i> , 2022 , 27, 22-22	4.2