Laura Gray

List of Publications by Citations

Source: https://exaly.com/author-pdf/6838242/laura-gray-publications-by-citations.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

150
papers

4,839
citations

h-index

67
g-index

173
ext. papers

5,999
ext. citations

5 avg, IF

L-index

#	Paper	IF	Citations
150	Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. <i>Diabetologia</i> , 2012 , 55, 2895-905	10.3	1100
149	The effects of high-intensity interval training on glucose regulation and insulin resistance: a meta-analysis. <i>Obesity Reviews</i> , 2015 , 16, 942-61	10.6	289
148	Ethnicity and clinical outcomes in COVID-19: A systematic review and meta-analysis. <i>EClinicalMedicine</i> , 2020 , 29, 100630	11.3	235
147	Associations of objectively measured sedentary behaviour and physical activity with markers of cardiometabolic health. <i>Diabetologia</i> , 2013 , 56, 1012-20	10.3	226
146	Effectiveness of a diabetes education and self management programme (DESMOND) for people with newly diagnosed type 2 diabetes mellitus: three year follow-up of a cluster randomised controlled trial in primary care. <i>BMJ, The</i> , 2012 , 344, e2333	5.9	214
145	Progression rates from HbA1c 6.0-6.4% and other prediabetes definitions to type 2 diabetes: a meta-analysis. <i>Diabetologia</i> , 2013 , 56, 1489-93	10.3	152
144	The Leicester Risk Assessment score for detecting undiagnosed Type 2 diabetes and impaired glucose regulation for use in a multiethnic UK setting. <i>Diabetic Medicine</i> , 2010 , 27, 887-95	3.5	122
143	Prevalence and Incidence of Hypoglycaemia in 532,542 People with Type 2 Diabetes on Oral Therapies and Insulin: A Systematic Review and Meta-Analysis of Population Based Studies. <i>PLoS ONE</i> , 2015 , 10, e0126427	3.7	119
142	Sedentary time and markers of chronic low-grade inflammation in a high risk population. <i>PLoS ONE</i> , 2013 , 8, e78350	3.7	109
141	The Effect of Glucagon-Like Peptide 1 Receptor Agonists on Weight Loss in Type 2 Diabetes: A Systematic Review and Mixed Treatment Comparison Meta-Analysis. <i>PLoS ONE</i> , 2015 , 10, e0126769	3.7	102
140	Effectiveness of the Stand More AT (SMArT) Work intervention: cluster randomised controlled trial. <i>BMJ, The,</i> 2018 , 363, k3870	5.9	76
139	Cardiovascular disease risk factors in chronic kidney disease: A systematic review and meta-analysis. <i>PLoS ONE</i> , 2018 , 13, e0192895	3.7	75
138	Defining obesity cut-off points for migrant South Asians. <i>PLoS ONE</i> , 2011 , 6, e26464	3.7	72
137	Rationale and design of the ADDITION-Leicester study, a systematic screening programme and randomised controlled trial of multi-factorial cardiovascular risk intervention in people with type 2 diabetes mellitus detected by screening. <i>Trials</i> , 2010 , 11, 16	2.8	68
136	Socio-demographic heterogeneity in the prevalence of COVID-19 during lockdown is associated with ethnicity and household size: Results from an observational cohort study. <i>EClinicalMedicine</i> , 2020 , 25, 100466	11.3	65
135	The association between neighbourhood greenspace and type 2 diabetes in a large cross-sectional study. <i>BMJ Open</i> , 2014 , 4, e006076	3	61
134	Detection of impaired glucose regulation and/or type 2 diabetes mellitus, using primary care electronic data, in a multiethnic UK community setting. <i>Diabetologia</i> , 2012 , 55, 959-66	10.3	60

(2014-2016)

133	and lifestyle intervention for prevention: the LetB Prevent Diabetes cluster randomised controlled trial. <i>Preventive Medicine</i> , 2016 , 84, 48-56	4.3	55	
132	Guidelines for managing diabetes in Ramadan. <i>Diabetic Medicine</i> , 2016 , 33, 1315-29	3.5	52	
131	Levels of physical activity and relationship with markers of diabetes and cardiovascular disease risk in 5474 white European and South Asian adults screened for type 2 diabetes. <i>Preventive Medicine</i> , 2010 , 51, 290-4	4.3	50	
130	The potential impact of using glycated haemoglobin as the preferred diagnostic tool for detecting Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2010 , 27, 762-9	3.5	49	
129	The potential impact and optimal cut-points of using glycated haemoglobin, HbA1c, to detect people with impaired glucose regulation in a UK multi-ethnic cohort. <i>Diabetes Research and Clinical Practice</i> , 2010 , 90, 100-8	7.4	48	
128	Walking Away from Type 2 diabetes: a cluster randomized controlled trial. <i>Diabetic Medicine</i> , 2017 , 34, 698-707	3.5	46	
127	Screening for diabetes using an oral glucose tolerance test within a western multi-ethnic population identifies modifiable cardiovascular risk: the ADDITION-Leicester study. <i>Diabetologia</i> , 2011 , 54, 2237-46	10.3	46	
126	Walking away from type 2 diabetes: trial protocol of a cluster randomised controlled trial evaluating a structured education programme in those at high risk of developing type 2 diabetes. <i>BMC Family Practice</i> , 2012 , 13, 46	2.6	44	
125	Stand More AT Work (SMArT Work): using the behaviour change wheel to develop an intervention to reduce sitting time in the workplace. <i>BMC Public Health</i> , 2018 , 18, 319	4.1	42	
124	A randomized controlled trial comparing the GLP-1 receptor agonist liraglutide to a sulphonylurea as add on to metformin in patients with established type 2 diabetes during Ramadan: the Treat 4 Ramadan Trial. <i>Diabetes, Obesity and Metabolism</i> , 2014 , 16, 527-36	6.7	41	
123	Patterns of Multimorbidity in Middle-Aged and Older Adults: An Analysis of the UK Biobank Data. <i>Mayo Clinic Proceedings</i> , 2018 , 93, 857-866	6.4	39	
122	Frailty Factors and Outcomes in Vascular Surgery Patients: A Systematic Review and Meta-analysis. <i>Annals of Surgery</i> , 2020 , 272, 266-276	7.8	37	
121	Ethnic differences in SARS-CoV-2 vaccine hesitancy in United Kingdom healthcare workers: Results from the UK-REACH prospective nationwide cohort study. <i>Lancet Regional Health - Europe, The</i> , 2021 , 9, 100180		37	
120	Fruit and vegetable intake and the association with glucose parameters: a cross-sectional analysis of the Letß Prevent Diabetes Study. <i>European Journal of Clinical Nutrition</i> , 2013 , 67, 12-7	5.2	34	
119	Multifactorial intervention in individuals with type 2 diabetes and microalbuminuria: the Microalbuminuria Education and Medication Optimisation (MEMO) study. <i>Diabetes Research and Clinical Practice</i> , 2011 , 93, 328-36	7.4	34	
118	Is the number of fast-food outlets in the neighbourhood related to screen-detected type 2 diabetes mellitus and associated risk factors?. <i>Public Health Nutrition</i> , 2015 , 18, 1698-705	3.3	33	
117	The effect of exercise training on intrahepatic triglyceride and hepatic insulin sensitivity: a systematic review and meta-analysis. <i>Obesity Reviews</i> , 2018 , 19, 1446-1459	10.6	32	
116	Body mass index and waist circumference cut-points in multi-ethnic populations from the UK and India: the ADDITION-Leicester, Jaipur heart watch and New Delhi cross-sectional studies. <i>PLoS ONE</i> , 2014 , 9, e90813	3.7	31	

115	Risk assessment tools for detecting those with pre-diabetes: a systematic review. <i>Diabetes Research and Clinical Practice</i> , 2014 , 105, 1-13	7.4	30
114	A systematic review and mixed treatment comparison of pharmacological interventions for the treatment of obesity. <i>Obesity Reviews</i> , 2012 , 13, 483-98	10.6	30
113	Letß prevent diabetes: study protocol for a cluster randomised controlled trial of an educational intervention in a multi-ethnic UK population with screen detected impaired glucose regulation. <i>Cardiovascular Diabetology</i> , 2012 , 11, 56	8.7	30
112	Implementation of the automated Leicester Practice Risk Score in two diabetes prevention trials provides a high yield of people with abnormal glucose tolerance. <i>Diabetologia</i> , 2012 , 55, 3238-44	10.3	30
111	Aspirin and cardiovascular primary prevention in non-endstage chronic kidney disease: A meta-analysis. <i>Atherosclerosis</i> , 2016 , 251, 177-182	3.1	27
110	Efficacy and tolerability of sodium-glucose co-transporter-2 inhibitors and glucagon-like peptide-1 receptor agonists: A systematic review and network meta-analysis. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 1035-1046	6.7	26
109	The Kidney Failure Risk Equation for prediction of end stage renal disease in UK primary care: An external validation and clinical impact projection cohort study. <i>PLoS Medicine</i> , 2019 , 16, e1002955	11.6	26
108	Effects of glucose-lowering and multifactorial interventions on cardiovascular and mortality outcomes: a meta-analysis of randomized control trials. <i>Diabetic Medicine</i> , 2016 , 33, 280-9	3.5	26
107	Type 2 diabetes mellitus and obesity in young adults: the extreme phenotype with early cardiovascular dysfunction. <i>Diabetic Medicine</i> , 2014 , 31, 794-8	3.5	25
106	The association between air pollution and type 2 diabetes in a large cross-sectional study in Leicester: The CHAMPIONS Study. <i>Environment International</i> , 2017 , 104, 41-47	12.9	23
105	Analysis of the Modified Rankin Scale in Randomised Controlled Trials of Acute Ischaemic Stroke: A Systematic Review. <i>Stroke Research and Treatment</i> , 2016 , 2016, 9482876	1.7	23
104	Safety and effectiveness of non-insulin glucose-lowering agents in the treatment of people with type 2 diabetes who observe Ramadan: a systematic review and meta-analysis. <i>Diabetes, Obesity and Metabolism</i> , 2015 , 17, 639-48	6.7	21
103	Development of a lifestyle intervention using the MRC framework for diabetes prevention in people with impaired glucose regulation. <i>Journal of Public Health</i> , 2016 , 38, 493-501	3.5	21
102	Cardiovascular efficacy and safety of sodium-glucose co-transporter-2 inhibitors and glucagon-like peptide-1 receptor agonists: a systematic review and network meta-analysis. <i>Diabetic Medicine</i> , 2019 , 36, 444-452	3.5	20
101	Type 2 diabetes in younger adults: clinical characteristics, diabetes-related complications and management of risk factors. <i>Primary Care Diabetes</i> , 2011 , 5, 57-62	2.4	19
100	Clinical characteristics of polycystic ovary syndrome: investigating differences in White and South Asian women. <i>Clinical Endocrinology</i> , 2015 , 83, 542-9	3.4	18
99	Multimorbidity and lifestyle factors among adults with intellectual disabilities: a cross-sectional analysis of a UK cohort. <i>Journal of Intellectual Disability Research</i> , 2019 , 63, 255-265	3.2	18
98	Cost-effectiveness of a pragmatic structured education intervention for the prevention of type 2 diabetes: economic evaluation of data from the LetB Prevent Diabetes cluster-randomised controlled trial. <i>BMJ Open</i> , 2017 , 7, e013592	3	17

(2017-2015)

97	Systematic Review and Meta-Analysis of Response Rates and Diagnostic Yield of Screening for Type 2 Diabetes and Those at High Risk of Diabetes. <i>PLoS ONE</i> , 2015 , 10, e0135702	3.7	17
96	The cost-effectiveness of testing strategies for type 2 diabetes: a modelling study. <i>Health Technology Assessment</i> , 2015 , 19, 1-80	4.4	17
95	Long-term mortality after acute myocardial infarction among individuals with and without diabetes: A systematic review and meta-analysis of studies in the post-reperfusion era. <i>Diabetes, Obesity and Metabolism,</i> 2017 , 19, 364-374	6.7	16
94	PRomotion Of Physical activity through structured Education with differing Levels of ongoing Support for people at high risk of type 2 diabetes (PROPELS): study protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 289	2.8	16
93	Screening for type 2 diabetes in a multiethnic setting using known risk factors to identify those at high risk: a cross-sectional study. <i>Vascular Health and Risk Management</i> , 2010 , 6, 837-42	4.4	16
92	Engagement, Retention, and Progression to Type 2 Diabetes: A Retrospective Analysis of the Cluster-Randomised "Letß Prevent Diabetes" Trial. <i>PLoS Medicine</i> , 2016 , 13, e1002078	11.6	16
91	An imbalance in cluster sizes does not lead to notable loss of power in cross-sectional, stepped-wedge cluster randomised trials with a continuous outcome. <i>Trials</i> , 2017 , 18, 109	2.8	15
90	Incident Type 2 diabetes and the effect of early regression to normoglycaemia in a population with impaired glucose regulation. <i>Diabetic Medicine</i> , 2017 , 34, 396-404	3.5	13
89	Incidence and severity of hypoglycaemia in type 2 diabetes by treatment regimen: A UK multisite 12-month prospective observational study. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 1585-1595	6.7	13
88	Increasing capacity to deliver diabetes self-management education: results of the DESMOND lay educator non-randomized controlled equivalence trial. <i>Diabetic Medicine</i> , 2014 , 31, 1431-8	3.5	13
87	Self-knowledge of HbA1c in people with Type 2 Diabetes Mellitus and its association with glycaemic control. <i>Primary Care Diabetes</i> , 2017 , 11, 414-420	2.4	13
86	Impact of Depression and Anxiety on Change to Physical Activity Following a Pragmatic Diabetes Prevention Program Within Primary Care: Pooled Analysis From Two Randomized Controlled Trials. <i>Diabetes Care</i> , 2019 , 42, 1847-1853	14.6	12
85	Unequal cluster sizes in stepped-wedge cluster randomised trials: a systematic review. <i>BMJ Open</i> , 2017 , 7, e017151	3	12
84	The development and validation of the Portuguese risk score for detecting type 2 diabetes and impaired fasting glucose. <i>Primary Care Diabetes</i> , 2013 , 7, 11-8	2.4	11
83	The effect of antiobesity drugs on waist circumference: a mixed treatment comparison. <i>Diabetes, Obesity and Metabolism</i> , 2014 , 16, 237-47	6.7	11
82	Rationale and design of the randomised controlled trial to assess the impact of liraglutide on cardiac function and structure in young adults with type 2 diabetes (the LYDIA study). <i>Cardiovascular Diabetology</i> , 2016 , 15, 102	8.7	11
81	A three arm cluster randomised controlled trial to test the effectiveness and cost-effectiveness of the SMART Work & Life intervention for reducing daily sitting time in office workers: study protocol. <i>BMC Public Health</i> , 2018 , 18, 1120	4.1	11
8o	Type 2 diabetes and glucose intolerance in a population with intellectual disabilities: the STOP diabetes cross-sectional screening study. <i>Journal of Intellectual Disability Research</i> , 2017 , 61, 668-681	3.2	10

79	Associations Between Anthropometric Measurements and Cardiometabolic Risk Factors in White European and South Asian Adults in the United Kingdom. <i>Mayo Clinic Proceedings</i> , 2017 , 92, 925-933	6.4	10
78	Screening for glucose intolerance and development of a lifestyle education programme for prevention of type 2 diabetes in a population with intellectual disabilities: the STOP Diabetes research project. <i>Programme Grants for Applied Research</i> , 2017 , 5, 1-316	1.5	10
77	A community faith centre based screening and educational intervention to reduce the risk of type 2 diabetes: A feasibility study. <i>Diabetes Research and Clinical Practice</i> , 2016 , 120, 73-80	7.4	8
76	Change in Sedentary Time, Physical Activity, Bodyweight, and HbA1c in High-Risk Adults. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 1120-1125	1.2	8
75	The United Kingdom Research study into Ethnicity And COVID-19 outcomes in Healthcare workers (UK-REACH): protocol for a prospective longitudinal cohort study of healthcare and ancillary workers in UK healthcare settings. <i>BMJ Open</i> , 2021 , 11, e050647	3	8
74	A randomized, open-label, active comparator trial assessing the effects of 26 weeks of liraglutide or sitagliptin on cardiovascular function in young obese adults with type 2 diabetes. <i>Diabetes, Obesity and Metabolism,</i> 2020 , 22, 1187-1196	6.7	7
73	The Role of the Oral Healthcare Team in Identification of Type 2 Diabetes Mellitus: A Systematic Review. <i>Current Oral Health Reports</i> , 2020 , 7, 87-97	1.2	7
72	A Cost and Cost-Benefit Analysis of the Stand More AT Work (SMArT Work) Intervention. <i>International Journal of Environmental Research and Public Health</i> , 2020 , 17,	4.6	7
71	Associations of Physical Activity Intensities with Markers of Insulin Sensitivity. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 2451-2458	1.2	7
70	External national validation of the Leicester Self-Assessment score for Type 2 diabetes using data from the English Longitudinal Study of Ageing. <i>Diabetic Medicine</i> , 2017 , 34, 1575-1583	3.5	7
69	The use of feasibility studies for stepped-wedge cluster randomised trials: protocol for a review of impact and scope. <i>BMJ Open</i> , 2017 , 7, e017290	3	7
68	Prevalence of Cognitive Impairment in Individuals with Vascular Surgical Pathology: a Systematic Review and Meta-Analysis. <i>European Journal of Vascular and Endovascular Surgery</i> , 2021 , 61, 664-674	2.3	7
67	Assessment of response rates and yields for Two opportunistic Tools for Early detection of Non-diabetic hyperglycaemia and Diabetes (ATTEND). A randomised controlled trial and cost-effectiveness analysis. <i>Diabetes Research and Clinical Practice</i> , 2016 , 118, 12-20	7.4	7
66	Baby Steps - a structured group education programme with accompanying mobile web application designed to promote physical activity in women with a history of gestational diabetes: study protocol for a randomised controlled trial. <i>Trials</i> , 2018 , 19, 682	2.8	7
65	Contrasting cardiovascular risk profiles and prescribed cardio-protective therapies in newly-diagnosed type 2 diabetes identified through screening and standard practice. <i>Diabetes Research and Clinical Practice</i> , 2011 , 91, 280-5	7.4	6
64	Identifying stroke therapeutics from preclinical models: A protocol for a novel application of network meta-analysis. <i>F1000Research</i> , 2019 , 8, 11	3.6	6
63	Associations of Clinical Frailty with Severity of Limb Threat and Outcomes in Chronic Limb-threatening Ischaemia. <i>Annals of Vascular Surgery</i> , 2021 , 76, 406-416	1.7	6
62	Neuroprotection by anaesthetics in rodent models of traumatic brain injury: a systematic review and network meta-analysis. <i>British Journal of Anaesthesia</i> , 2018 , 121, 1272-1281	5.4	6

61	Healthcare workersPviews on mandatory SARS-CoV-2 vaccination in the UK: A cross-sectional, mixed-methods analysis from the UK-REACH study <i>EClinicalMedicine</i> , 2022 , 101346	11.3	6
60	The current use of feasibility studies in the assessment of feasibility for stepped-wedge cluster randomised trials: a systematic review. <i>BMC Medical Research Methodology</i> , 2019 , 19, 12	4.7	5
59	Epidemiology of major lower limb amputation using routinely collected electronic health data in the UK: a systematic review protocol. <i>BMJ Open</i> , 2020 , 10, e037053	3	5
58	A five-year retrospective study shows increasing rates of antimicrobial drug resistance in Cabo Verde for both Staphylococcus aureus and Escherichia coli. <i>Journal of Global Antimicrobial Resistance</i> , 2020 , 22, 483-487	3.4	5
57	Diagnostic performance of using one- or two-HbA1c cut-point strategies to detect undiagnosed type 2 diabetes and impaired glucose regulation within a multi-ethnic population. <i>Diabetes and Vascular Disease Research</i> , 2013 , 10, 84-92	3.3	5
56	A community-based primary prevention programme for type 2 diabetes mellitus integrating identification and lifestyle intervention for prevention: a cluster randomised controlled trial. <i>Programme Grants for Applied Research</i> , 2017 , 5, 1-290	1.5	5
55	Movement through Active Personalised engagement (MAP) - a self-management programme designed to promote physical activity in people with multimorbidity: study protocol for a randomised controlled trial. <i>Trials</i> , 2018 , 19, 576	2.8	5
54	Predicting hospital stay, mortality and readmission in people admitted for hypoglycaemia: prognostic models derivation and validation. <i>Diabetologia</i> , 2017 , 60, 1007-1015	10.3	4
53	Reducing sitting at work: process evaluation of the SMArT Work (Stand More At Work) intervention. <i>Trials</i> , 2020 , 21, 403	2.8	4
52	Use of dental practices for the identification of adults with undiagnosed type 2 diabetes mellitus or non-diabetic hyperglycaemia: a systematic review. <i>Diabetic Medicine</i> , 2020 , 37, 1443-1453	3.5	4
51	Persistent hesitancy for SARS-CoV-2 vaccines among healthcare workers in the United Kingdom: analysis of longitudinal data from the UK-REACH cohort study <i>Lancet Regional Health - Europe, The</i> , 2022 , 13, 100299		4
50	The Ready to Reduce Risk (3R) Study for a Group Educational Intervention With Telephone and Text Messaging Support to Improve Medication Adherence for the Primary Prevention of Cardiovascular Disease: Protocol for a Randomized Controlled Trial. <i>JMIR Research Protocols</i> , 2018 , 7, e11289	2	4
49	Ethnic differences in SARS-CoV-2 vaccine hesitancy in United Kingdom healthcare workers: Results from the UK-REACH prospective nationwide cohort study		4
48	Ethnic disparities in COVID-19: increased risk of infection or severe disease?. Lancet, The, 2021 , 398, 38	9-490	4
47	Efficacy and safety of sodium-glucose cotransporter 2 inhibitors (SGLT-2is) and glucagon-like peptide-1 receptor agonists (GLP-1RAs) in patients with type 2 diabetes: a systematic review and network meta-analysis study protocol. <i>BMJ Open</i> , 2018 , 8, e023206	3	4
46	Access to personal protective equipment in healthcare workers during the COVID-19 pandemic in the United Kingdom: results from a nationwide cohort study (UK-REACH)		4
45	Evaluation and refinement of the PRESTARt tool for identifying 12-14 year olds at high lifetime risk of developing type 2 diabetes compared to a clinicians assessment of risk: a cross-sectional study. BMC Endocrine Disorders, 2019 , 19, 79	3.3	3
44	External validation of two diabetes risk scores in a young UK South Asian population. <i>Diabetes Research and Clinical Practice</i> , 2014 , 104, 451-8	7.4	3

Cluster randomised controlled trial to investigate the effectiveness and cost-effectiveness of a 43 Structured Health Intervention For Truckers (the SHIFT study): a study protocol. BMJ Open, **2019**, 9, e03 $^{\circ}$ 175 Ethnicity and clinical outcomes in COVID-19: A Systematic Review and Meta-analysis 42 Missed Opportunities for Timely Recognition of Chronic Limb Threatening Ischaemia in Patients Undergoing a Major Amputation: A Population Based Cohort Study Using the UKB Clinical Practice 41 2.3 3 Research Datalink. European Journal of Vascular and Endovascular Surgery, 2020, 60, 703-710 Effectiveness of the Ready to Reduce Risk (3R) complex intervention for the primary prevention of 40 11.4 cardiovascular disease: a pragmatic randomised controlled trial. BMC Medicine, 2020, 18, 198 COVID-19 and the new variant strain in England - What are the implications for those from ethnic 39 11.3 3 minority groups?. EClinicalMedicine, 2021, 33, 100805 United Kingdom Research study into Ethnicity And COVID-19 outcomes in Healthcare workers (UK-REACH): a retrospective cohort study using linked routinely collected data, study protocol. BMJ 38 Open, **2021**, 11, e046392 Diabetes in adults with intellectual disability: prevalence and associated demographic, lifestyle, 3.2 2 37 independence and health factors. Journal of Intellectual Disability Research, 2020, 64, 287-295 36 LetB Prevent Diabetes: from idea to implementation. Practical Diabetes, 2017, 34, 55-57 0.7 Association between undiagnosed hypertension and microalbuminuria in South Asians without 2.6 2 35 known diabetes. Journal of Human Hypertension, 2015, 29, 185-9 South Asian individuals at high risk of type 2 diabetes have lower plasma vitamin C levels than 34 2.7 white Europeans. Journal of Nutritional Science, 2013, 2, e21 The Microalbuminuria Education Medication and Optimisation (MEMO) study: 4 years follow-up of 2 33 multifactorial intervention in high-risk individuals with type 2 diabetes. Diabetic Medicine, 2020, 37, 286-297 Comorbidities and outcomes in South Asian individuals with chronic kidney disease: an 4.3 observational primary care cohort. Nephrology Dialysis Transplantation, 2021, Sleep duration and sleep efficiency in UK long-distance heavy goods vehicle drivers. Occupational 31 2.1 2 and Environmental Medicine, 2021, Risk factors associated with SARS-CoV-2 infection in a multiethnic cohort of United Kingdom 2 30 healthcare workers (UK-REACH): A cross-sectional analysis. PLoS Medicine, 2022, 19, e1004015 The impact of an intervention to increase uptake to structured self-management education for people with type 2 diabetes mellitus in primary care (the embedding package), compared to usual 2.6 1 29 care, on glycaemic control: study protocol for a mixed methods study incorporating a wait-list Predictors of SARS-CoV-2 infection in a multi-ethnic cohort of United Kingdom healthcare workers: 28 a prospective nationwide cohort study (UK-REACH) The General Practitioner Prompt Study to Reduce Cardiovascular and Renal Complications in Patients With Type 2 Diabetes and Renal Complications: Protocol and Baseline Characteristics for a 27 1 Cluster Randomized Controlled Trial. JMIR Research Protocols, 2018, 7, e152 Use of Dental Practices for the Identification of Adults With Undiagnosed Type 2 Diabetes Mellitus 26 or Nondiabetic Hyperglycemia: Protocol for a Systematic Review. JMIR Research Protocols, 2018, 7, e11843

Screening for Type 2 Diabetes 2014, 119-126 25 1 Increasing uptake of structured self-management education programmes for type 2 diabetes in a 24 1.9 primary care setting: a feasibility study. Pilot and Feasibility Studies, 2020, 6, 71 Prediction of type 2 diabetes risk in people with non-diabetic hyperglycaemia: model derivation 23 3 1 and validation using UK primary care data. BMJ Open, 2020, 10, e037937 Does the Introduction of a Vascular Limb Salvage Service Improve One Year Amputation Outcomes for Patients with Chronic Limb-Threatening Ischaemia?. European Journal of Vascular and 2.3 Endovascular Surgery, 2021, 61, 612-619 Promoting physical activity in a multi-ethnic population at high risk of diabetes: the 48-month 21 11.4 1 PROPELS randomised controlled trial. BMC Medicine. 2021. 19, 130 Leg ischaemia management collaboration (LIMb): study protocol for a prospective cohort study at a 20 single UK centre. BMJ Open, 2019, 9, e031257 Development of an Interactive Lifestyle Programme for Adolescents at Risk of Developing Type 2 2.8 19 1 Diabetes: PRE-STARt. Children, 2021, 8, The United Kingdom Research study into Ethnicity And COVID-19 outcomes in Healthcare workers 18 (UK-REACH): Protocol for a prospective longitudinal cohort study of healthcare and ancillary workers in UK healthcare settings Screening for Diabetes and Prediabetes. Endocrinology, 2018, 369-400 17 0.1 7 Promoting physical activity with self-management support for those with multimorbidity: a 16 1.6 randomised controlled trial. British Journal of General Practice, 2021, 71, e921-e930 Whole-genome analysis uncovers loss of blaZ associated with carriage isolates belonging to methicillin-resistant Staphylococcus aureus (MRSA) clone ST5-VI in Cape Verde. Journal of Global 15 1 3.4 Antimicrobial Resistance, 2021, 26, 77-83 The association of blood pressure variability with adverse outcomes in a primary care chronic 14 1.9 \circ kidney disease cohort. Journal of Hypertension, 2021, 39, 2067-2074 Developing a Conceptually Equivalent Type 2 Diabetes Risk Score for Indian Gujaratis in the UK. 13 3.9 O Journal of Diabetes Research, 2016, 2016, 8107108 Measuring quality of life in trials including patients on dialysis: how are transplants and mortality 0 incorporated into the analysis? A systematic review protocol. BMJ Open, 2021, 11, e048179 Evaluation of the design, conduct and reporting of randomised controlled trials in the 11 3 O haemodialysis population: a scoping review and interview study.. BMJ Open, 2022, 12, e058368 Behavioural interventions to promote physical activity in a multiethnic population at high risk of 10 diabetes: PROPELS three-arm RCT.. Health Technology Assessment, 2021, 25, 1-190 The effect of exercise training on adipose tissue insulin sensitivity: A systematic review and 10.6 0 9 meta-analysis.. Obesity Reviews, 2022, e13445 Understanding variations in reported epidemiology of major lower extremity amputation in the UK: a systematic review **2021**, 11, e053599

7	Effects of an Electronic Software "Prompt" With Health Care Professional Training on Cardiovascular and Renal Complications in a Multiethnic Population With Type 2 Diabetes and Microalbuminuria (the GP-Prompt Study): Results of a Pragmatic Cluster-Randomized Trial.	14.6
6	Diabetes Care, 2020, 43, 1893-1901 Screening for Diabetes and Prediabetes. Endocrinology, 2018, 1-33	0.1
5	Predictors of adverse outcome in the first and second waves of the COVID-19 pandemic: results from a UK centre <i>Therapeutic Advances in Infectious Disease</i> , 2022 , 9, 20499361221074569	2.8
4	Understanding variations in reported epidemiology of major lower extremity amputation in the UK: a systematic review. <i>BMJ Open</i> , 2021 , 11, e053599	3
3	Risk Assessment for Diabetes 2020 , 61-76	
2	Mind the gap: covariate constrained randomisation can protect against substantial power loss in parallel cluster randomised trials <i>BMC Medical Research Methodology</i> , 2022 , 22, 111	4.7
1	Fibro-inflammatory recovery and type 2 diabetes remission following a low calorie diet but not exercise training: a secondary analysis of the DIASTOLIC randomised controlled trial <i>Diabetic Medicine</i> 2022, e14884	3.5