

Yoshiyuki Tanaka

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6844817/publications.pdf>

Version: 2024-02-01

19
papers

461
citations

687363

13
h-index

794594

19
g-index

19
all docs

19
docs citations

19
times ranked

304
citing authors

#	ARTICLE	IF	CITATIONS
1	Fasciation in Strawberry Floral Organs and Possible Implications for Floral Transition. Horticulture Journal, 2022, 91, 58-67.	0.8	2
2	Morphological and gene expression characterization of maf-1, a floral chili pepper mutant caused by a nonsense mutation in CaLFY. Molecular Breeding, 2022, 42, .	2.1	1
3	Characterization and bulk segregant analysis of a novel seedless mutant tn-1 of chili pepper (<i>Capsicum</i>) Tj ETQq1 1,0,784314 rgBT /O	3.6	2
4	Capsaicinoid biosynthesis in the pericarp of chili pepper fruits is associated with a placental septum-like transcriptome profile and tissue structure. Plant Cell Reports, 2021, 40, 1859-1874.	5.6	7
5	Effects of Intermittent Low Temperature Storage Duration and Cycle on the Growth and Flowering of <i>Eustoma</i> (&Eustoma grandiflorum L.) Seedlings Raised in the Summer. Horticulture Journal, 2020, 89, 292-299.	0.8	1
6	Multiple Non-pungent <i>Capsicum chinense</i> Accessions with a Loss of Function <i>CaKR1</i> Allele Originating from South America. Horticulture Journal, 2020, 89, 460-465.	0.8	2
7	Positional differences of intronic transposons in <i>pAMT</i> affect the pungency level in chili pepper through altered splicing efficiency. Plant Journal, 2019, 100, 693-705.	5.7	23
8	Mutation in the putative ketoacyl-ACP reductase <i>CaKR1</i> induces loss of pungency in <i>Capsicum</i> . Theoretical and Applied Genetics, 2019, 132, 65-80.	3.6	43
9	Identification of a Novel Mutant <i>pAMT</i> Allele Responsible for Low-pungency and Capsinoid Production in Chili Pepper: Accession "No. A4034" (<i>Capsicum chinense</i>). Horticulture Journal, 2018, 87, 222-228.	0.8	14
10	Difference in capsaicinoid biosynthesis gene expression in the pericarp reveals elevation of capsaicinoid contents in chili peppers (<i>Capsicum chinense</i>). Plant Cell Reports, 2017, 36, 267-279.	5.6	39
11	Non-pungency in a Japanese Chili Pepper Landrace (<i>Capsicum annuum</i>) is Caused by a Novel Loss-of-function <i>Pun1</i> Allele. Horticulture Journal, 2017, 86, 61-69.	0.8	21
12	Multiple loss-of-function putative aminotransferase alleles contribute to low pungency and capsinoid biosynthesis in <i>Capsicum chinense</i> . Molecular Breeding, 2015, 35, 1.	2.1	27
13	Incidence of Blossom-end Rot in Relation to the Water-soluble Calcium Concentration in Tomato Fruits as Affected by Calcium Nutrition and Cropping Season. Japanese Society for Horticultural Science, 2014, 83, 282-289.	0.8	14
14	Analysis of Non-pungency, Aroma, and Origin of a <i>Capsicum chinense</i> Cultivar from a Caribbean Island. Japanese Society for Horticultural Science, 2014, 83, 244-251.	0.8	17
15	Application of marker-assisted selection in breeding of a new fresh pepper cultivar (<i>Capsicum annuum</i>) containing capsinoids, low-pungent capsaicinoid analogs. Scientia Horticulturae, 2014, 165, 242-245.	3.6	22
16	Newly Mutated putative-aminotransferase in Nonpungent Pepper (<i>Capsicum annuum</i>) Results in Biosynthesis of Capsinoids, Capsaicinoid Analogues. Journal of Agricultural and Food Chemistry, 2010, 58, 1761-1767.	5.2	53
17	Novel Loss-of-Function putative aminotransferase Alleles Cause Biosynthesis of Capsinoids, Nonpungent Capsaicinoid Analogues, in Mildly Pungent Chili Peppers (<i>Capsicum chinense</i>). Journal of Agricultural and Food Chemistry, 2010, 58, 11762-11767.	5.2	40
18	Functional loss of <i>pAMT</i> results in biosynthesis of capsinoids, capsaicinoid analogs, in <i>Capsicum annuum</i> cv. CH19 Sweet. Plant Journal, 2009, 59, 953-961.	5.7	98

#	ARTICLE	IF	CITATIONS
19	Assessment of Capsiconinoid Composition, Nonpungent Capsaicinoid Analogues, in <i>Capsicum</i> Cultivars. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 5407-5412.	5.2	35