

Anna-Karin Wikström

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/684754/publications.pdf>

Version: 2024-02-01

101
papers

3,299
citations

186265

28
h-index

168389

53
g-index

103
all docs

103
docs citations

103
times ranked

4528
citing authors

#	ARTICLE	IF	CITATIONS
1	Maternal Obesity and Risk of Preterm Delivery. JAMA - Journal of the American Medical Association, 2013, 309, 2362.	7.4	550
2	The risk of maternal ischaemic heart disease after gestational hypertensive disease. BJOG: an International Journal of Obstetrics and Gynaecology, 2005, 112, 1486-1491.	2.3	202
3	Placental Growth Factor and Soluble FMS-Like Tyrosine Kinase-1 in Early-Onset and Late-Onset Preeclampsia. Obstetrics and Gynecology, 2007, 109, 1368-1374.	2.4	172
4	Tobacco Use During Pregnancy and Preeclampsia Risk. Hypertension, 2010, 55, 1254-1259.	2.7	154
5	Maternal Body Mass Index, Height, and Risks of Preeclampsia. American Journal of Hypertension, 2012, 25, 120-125.	2.0	110
6	The Swedish Pregnancy Register – for quality of care improvement and research. Acta Obstetrica Et Gynecologica Scandinavica, 2018, 97, 466-476.	2.8	101
7	Association Between Pregnancy and Perinatal Outcomes Among Women With Epilepsy. JAMA Neurology, 2017, 74, 983.	9.0	90
8	Induction of labour at 41 weeks versus expectant management and induction of labour at 42 weeks (SWEdish Post-term Induction Study, SWEPIS): multicentre, open label, randomised, superiority trial. BMJ: British Medical Journal, 2019, 367, l6131.	2.3	87
9	Hyperemesis gravidarum and risks of placental dysfunction disorders: a population-based cohort study. BJOG: an International Journal of Obstetrics and Gynaecology, 2013, 120, 541-547.	2.3	86
10	Maternal Use of Swedish Snuff (Snus) and Risk of Stillbirth. Epidemiology, 2010, 21, 772-778.	2.7	66
11	Influence of smoking and snuff cessation on risk of preterm birth. European Journal of Epidemiology, 2012, 27, 297-304.	5.7	58
12	Delineating the association between mode of delivery and postpartum depression symptoms: a longitudinal study. Acta Obstetrica Et Gynecologica Scandinavica, 2018, 97, 301-311.	2.8	56
13	Pregnancy Weight Gain Before Diagnosis and Risk of Preeclampsia. Hypertension, 2018, 72, 433-441.	2.7	53
14	Influence of Snuff and Smoking Habits in Early Pregnancy on Risks for Stillbirth and Early Neonatal Mortality. Nicotine and Tobacco Research, 2014, 16, 78-83.	2.6	51
15	Angiotensin-1/Angiotensin-2 Ratio for Prediction of Preeclampsia. American Journal of Hypertension, 2009, 22, 891-895.	2.0	49
16	Prehypertension in Pregnancy and Risks of Small for Gestational Age Infant and Stillbirth. Hypertension, 2016, 67, 640-646.	2.7	49
17	Delineating the Association between Heavy Postpartum Haemorrhage and Postpartum Depression. PLoS ONE, 2016, 11, e0144274.	2.5	49
18	Maternal overweight and obesity and risk of pre-eclampsia in women with type 1 diabetes or type 2 diabetes. Diabetologia, 2016, 59, 2099-2105.	6.3	48

#	ARTICLE	IF	CITATIONS
19	Previous preeclampsia and risks of adverse outcomes in subsequent nonpreeclamptic pregnancies. <i>American Journal of Obstetrics and Gynecology</i> , 2011, 204, 148.e1-148.e6.	1.3	47
20	Placental abruption and long-term maternal cardiovascular disease mortality: a population-based registry study in Norway and Sweden. <i>European Journal of Epidemiology</i> , 2016, 31, 501-511.	5.7	45
21	Maternal Snuff Use and Smoking and the Risk of Oral Cleft Malformations - A Population-Based Cohort Study. <i>PLoS ONE</i> , 2014, 9, e84715.	2.5	44
22	Relationship of Maternal Snuff Use and Cigarette Smoking With Neonatal Apnea. <i>Pediatrics</i> , 2011, 128, 503-509.	2.1	40
23	Psychosocial Stress Related to the Loss of a Close Relative the Year Before or During Pregnancy and Risk of Preeclampsia. <i>Hypertension</i> , 2013, 62, 183-189.	2.7	37
24	Early-onset Preeclampsia in Lupus Pregnancy. <i>Paediatric and Perinatal Epidemiology</i> , 2017, 31, 29-36.	1.7	36
25	Thyroid Testing and Management of Hypothyroidism During Pregnancy: A Population-based Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 2687-2692.	3.6	34
26	Risk of placental dysfunction disorders after prior miscarriages: a population-based study. <i>American Journal of Obstetrics and Gynecology</i> , 2014, 211, 34.e1-34.e8.	1.3	34
27	Risk of retained placenta in women previously delivered by caesarean section: a population-based cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014, 121, 224-229.	2.3	33
28	Elective induction of labor and the risk of cesarean section in low-risk parous women: a cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2013, 92, 198-203.	2.8	32
29	Iodine deficiency in a study population of pregnant women in Sweden. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2015, 94, 1168-1174.	2.8	28
30	Born Small for Gestational Age and Poor School Performance – How Small Is Too Small?. <i>Hormone Research in Paediatrics</i> , 2017, 88, 215-223.	1.8	28
31	Association of Attention-Deficit/Hyperactivity Disorder With Teenage Birth Among Women and Girls in Sweden. <i>JAMA Network Open</i> , 2019, 2, e1912463.	5.9	28
32	Early postpartum changes in circulating pro- and anti-angiogenic factors in early-onset and late-onset pre-eclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2008, 87, 146-153.	2.8	27
33	Antepartum risk factors for moderate to severe neonatal hypoxic ischemic encephalopathy: a Swedish national cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 615-623.	2.8	27
34	Clinical risk assessment in early pregnancy for preeclampsia in nulliparous women: A population based cohort study. <i>PLoS ONE</i> , 2019, 14, e0225716.	2.5	26
35	High Maternal Body Mass Index in Early Pregnancy and Risks of Stillbirth and Infant Mortality – A Population-Based Sibling Study in Sweden. <i>American Journal of Epidemiology</i> , 2016, 184, 98-105.	3.4	25
36	Epidural Analgesia During Childbirth and Postpartum Depressive Symptoms: A Population-Based Longitudinal Cohort Study. <i>Anesthesia and Analgesia</i> , 2020, 130, 615-624.	2.2	25

#	ARTICLE	IF	CITATIONS
37	Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials. <i>PLoS Medicine</i> , 2020, 17, e1003436.	8.4	25
38	Fibrinogen and histidine-rich glycoprotein in early-onset preeclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2010, 89, 131-139.	2.8	24
39	Targeted Thyroid Testing During Pregnancy in Clinical Practice. <i>Obstetrics and Gynecology</i> , 2014, 124, 10-15.	2.4	23
40	Labour dystocia risk of recurrence and instrumental delivery in following labour a population-based cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012, 119, 1648-1656.	2.3	22
41	Consanguineous marriage, prepregnancy maternal characteristics and stillbirth risk: a population-based case-control study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2015, 94, 1095-1101.	2.8	22
42	Proton Pump Inhibitors and Preeclampsia Risk Among 157,720 Women. <i>Hypertension</i> , 2019, 73, 1097-1103.	2.7	22
43	Placental Expression of proBNP/NT-proBNP and Plasma Levels of NT-proBNP in Early- and Late-Onset Preeclampsia. <i>American Journal of Hypertension</i> , 2014, 27, 1225-1230.	2.0	21
44	Study protocol of SWEPISS a Swedish multicentre register based randomised controlled trial to compare induction of labour at 41 completed gestational weeks versus expectant management and induction at 42 completed gestational weeks. <i>BMC Pregnancy and Childbirth</i> , 2016, 16, 49.	2.4	20
45	Pregnancy outcome in women with polycystic ovary syndrome in relation to second-trimester testosterone levels. <i>Reproductive BioMedicine Online</i> , 2021, 42, 217-225.	2.4	20
46	Preeclampsia and Increased Permeability Over the Blood-Brain Barrier: A Role of Vascular Endothelial Growth Receptor 2. <i>American Journal of Hypertension</i> , 2021, 34, 73-81.	2.0	20
47	Increased Circulating Levels of the Antiangiogenic Factor Endostatin in Early-Onset But Not Late-Onset Preeclampsia. <i>Reproductive Sciences</i> , 2009, 16, 995-1000.	2.5	18
48	Placental location and pregnancy outcomes in nulliparous women: A population-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 988-996.	2.8	18
49	Plasma Levels of S100B in Preeclampsia and Association With Possible Central Nervous System Effects. <i>American Journal of Hypertension</i> , 2014, 27, 1105-1111.	2.0	17
50	From infancy to pregnancy: birth weight, body mass index, and the risk of gestational diabetes. <i>European Journal of Epidemiology</i> , 2012, 27, 799-805.	5.7	16
51	Interpregnancy Change in Smoking Habits and Risk of Preeclampsia: A Population-Based Study. <i>American Journal of Hypertension</i> , 2012, 25, 372-378.	2.0	16
52	Neighborhood deprivation and adverse perinatal outcomes in Sweden: A population-based register study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 1004-1013.	2.8	16
53	Cerebral Magnesium Levels in Preeclampsia; A Phosphorus Magnetic Resonance Spectroscopy Study. <i>American Journal of Hypertension</i> , 2017, 30, 667-672.	2.0	15
54	SSRI use during pregnancy and risk for postpartum haemorrhage: a national register-based cohort study in Sweden. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2020, 127, 1366-1373.	2.3	15

#	ARTICLE	IF	CITATIONS
55	Placental location and risk of retained placenta in women with a previous cesarean section: A population-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020, 99, 1666-1673.	2.8	15
56	Induction of labor versus expectant management of large-for-gestational-age infants in nulliparous women. <i>PLoS ONE</i> , 2017, 12, e0180748.	2.5	15
57	The Impact of Small for Gestational Age Births in Early and Late Preeclamptic Pregnancies for Preeclampsia Recurrence: a Cohort Study of Successive Pregnancies in Sweden. <i>Paediatric and Perinatal Epidemiology</i> , 2016, 30, 563-570.	1.7	14
58	Obstetric emergencies as antecedents to neonatal hypoxic ischemic encephalopathy, does parity matter?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 1396-1404.	2.8	14
59	Recurrence of placental dysfunction disorders across generations. <i>American Journal of Obstetrics and Gynecology</i> , 2011, 205, 454.e1-454.e8.	1.3	13
60	Serum Pentraxin 3 is associated with signs of arterial alteration in women with preeclampsia. <i>International Journal of Cardiology</i> , 2017, 241, 417-422.	1.7	13
61	Assessment of cerebral perfusion and edema in preeclampsia with intravoxel incoherent motion <scp>MRI</scp>. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 1212-1218.	2.8	13
62	Pregnancy and neonatal complications in women with polycystic ovary syndrome in relation to second-trimester anti-Müllerian hormone levels. <i>Reproductive BioMedicine Online</i> , 2019, 39, 141-148.	2.4	13
63	Prediction of Preeclampsia by Combining Serum Histidine-Rich Glycoprotein and Uterine Artery Doppler. <i>American Journal of Hypertension</i> , 2012, 25, 1305-10.	2.0	12
64	Early second-trimester plasma levels of NT-proBNP in women who subsequently develop early-onset preeclampsia. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2017, 30, 2163-2165.	1.5	12
65	Polycystic ovary syndrome and extremely preterm birth: A nationwide register-based study. <i>PLoS ONE</i> , 2021, 16, e0246743.	2.5	12
66	Opium use during pregnancy and risk of preterm delivery: A population-based cohort study. <i>PLoS ONE</i> , 2017, 12, e0176588.	2.5	11
67	Evaluation of the discrepancy between pH and lactate in combined fetal scalp blood sampling. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2011, 90, 1088-1093.	2.8	10
68	Cerebral Osmolytes and Plasma Osmolality in Pregnancy and Preeclampsia: A Proton Magnetic Resonance Spectroscopy Study. <i>American Journal of Hypertension</i> , 2018, 31, 847-853.	2.0	10
69	Women's childbirth experiences in the Swedish Post-term Induction Study (SWEPIS): a multicentre, randomised, controlled trial. <i>BMJ Open</i> , 2021, 11, e042340.	1.9	10
70	Chronic Hypertension in Women after Perinatal Exposure to Preeclampsia, Being Born Small for Gestational Age or Preterm. <i>Paediatric and Perinatal Epidemiology</i> , 2017, 31, 89-98.	1.7	9
71	Delivery outcome after trial of labor in nulliparous women 40 years or older: A nationwide population-based study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 1195-1203.	2.8	9
72	Preterm Delivery Phenotypes in Systemic Lupus Erythematosus Pregnancies. <i>American Journal of Perinatology</i> , 2019, 36, 964-968.	1.4	9

#	ARTICLE	IF	CITATIONS
73	Prenatal exposure to preeclampsia is associated with accelerated height gain in early childhood. PLoS ONE, 2018, 13, e0192514.	2.5	9
74	Gestational Diabetes Mellitus Risk in Pregnant Women With Systemic Lupus Erythematosus. Journal of Rheumatology, 2022, 49, 465-469.	2.0	9
75	Opium use during pregnancy and infant size at birth: a cohort study. BMC Pregnancy and Childbirth, 2018, 18, 358.	2.4	8
76	Slow fetal growth between first and early second trimester ultrasound scans and risk of small for gestational age (SGA) birth. PLoS ONE, 2017, 12, e0184853.	2.5	7
77	Pregnancy in metabolic healthy and unhealthy obese women. Acta Obstetrica Et Gynecologica Scandinavica, 2020, 99, 1640-1648.	2.8	7
78	Maternal Hypertensive Disorders in Pregnant Women With Systemic Lupus Erythematosus and Future Cardiovascular Outcomes. Arthritis Care and Research, 2021, 73, 574-579.	3.4	7
79	Hypoxic ischemic encephalopathy in offspring of immigrant women in Sweden: A population-based cohort study. Acta Obstetrica Et Gynecologica Scandinavica, 2021, 100, 2285-2293.	2.8	7
80	Factors associated with re-initiation of antidepressant treatment following discontinuation during pregnancy: a register-based cohort study. Archives of Women's Mental Health, 2020, 23, 709-717.	2.6	6
81	Efficacy and safety of oral misoprostol vs transvaginal balloon catheter for labor induction: An observational study within the SWEdish Postterm Induction Study (SWEPIIS). Acta Obstetrica Et Gynecologica Scandinavica, 2021, 100, 1463-1477.	2.8	5
82	Time matters—a Swedish cohort study of labor duration and risk of uterine rupture. Acta Obstetrica Et Gynecologica Scandinavica, 2021, 100, 1902-1909.	2.8	5
83	The association between postpartum hemorrhage and postpartum depression: A Swedish national register-based study. PLoS ONE, 2021, 16, e0255938.	2.5	5
84	Accelerated fetal growth in early pregnancy and risk of severe large-for-gestational-age and macrosomic infant: a cohort study in a low-risk population. Acta Obstetrica Et Gynecologica Scandinavica, 2017, 96, 1261-1268.	2.8	4
85	Evidence of under-reporting of early-onset preeclampsia using register data. Paediatric and Perinatal Epidemiology, 2021, 35, 596-600.	1.7	4
86	Experience of fetal scalp blood sampling during labor. Acta Obstetrica Et Gynecologica Scandinavica, 2014, 93, 113-117.	2.8	3
87	Accelerated fetal growth in early pregnancy and risk of preterm birth: a prospective cohort study. BMC Pregnancy and Childbirth, 2020, 20, 764.	2.4	3
88	Self-rated health before pregnancy and adverse birth outcomes in Sweden: A population-based register study. Birth, 2021, 48, 541-549.	2.2	3
89	Neuroticism is associated with higher antenatal care utilization in obstetric low-risk women. Acta Obstetrica Et Gynecologica Scandinavica, 2019, 98, 470-478.	2.8	2
90	Postnatal growth in children born small for gestational age with and without smoking mother. Pediatric Research, 2019, 85, 961-966.	2.3	2

#	ARTICLE	IF	CITATIONS
91	Study for Improving Maternal Pregnancy And Child ouTcomes (IMPACT): a study protocol for a Swedish prospective multicentre cohort study. <i>BMJ Open</i> , 2020, 10, e033851.	1.9	2
92	COVID-19 in Pregnancy and Early Childhood (COPE): study protocol for a prospective, multicentre biobank, survey and database cohort study. <i>BMJ Open</i> , 2021, 11, e049376.	1.9	2
93	Does Use of Low-Molecular-Weight Heparin during Pregnancy Influence the Risk of Prolonged Labor: A Population-Based Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0140422.	2.5	2
94	Study protocol: establishment of a multicentre pre-eclampsia database and biobank in Sweden: GO PROVE and UP MOST, a prospective cohort study. <i>BMJ Open</i> , 2021, 11, e049559.	1.9	2
95	Associations between capillary glucose during pregnancy and childhood growth to the age of five: a cohort study. <i>Scientific Reports</i> , 2022, 12, 1832.	3.3	2
96	Maternal Obesity and Risk of Preterm Delivery. <i>Survey of Anesthesiology</i> , 2014, 58, 70-71.	0.1	1
97	Plasma from women with preeclampsia impairs the blood brain barrier in an in vitro model. <i>Pregnancy Hypertension</i> , 2019, 17, S7.	1.4	1
98	Evaluation of the Discrepancy Between pH and Lactate in Combined Fetal Scalp Blood Sampling. <i>Obstetric Anesthesia Digest</i> , 2012, 32, 177-178.	0.1	0
99	Maternal Obesity and Risk of Preterm Delivery. <i>Obstetrical and Gynecological Survey</i> , 2013, 68, 731-732.	0.4	0
100	Induction of Labour at 41 Weeks Versus Expectant Management and Induction of Labour at 42 Weeks (SWEdish Post-Term Induction Study, SWEPIS): Multicentre, Open Label, Randomised, Superiority Trial. <i>Obstetrical and Gynecological Survey</i> , 2020, 75, 207-209.	0.4	0
101	Reply to: Re: Risk of preâ€eclampsia after gastric bypass: a matched cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2021, , .	2.3	0