

# Anna-Karin Wikström

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/684754/publications.pdf>

Version: 2024-02-01

101  
papers

3,299  
citations

186265

28  
h-index

168389

53  
g-index

103  
all docs

103  
docs citations

103  
times ranked

4528  
citing authors

#	ARTICLE	IF	CITATIONS
1	Gestational Diabetes Mellitus Risk in Pregnant Women With Systemic Lupus Erythematosus. <i>Journal of Rheumatology</i> , 2022, 49, 465-469.	2.0	9
2	Associations between capillary glucose during pregnancy and childhood growth to the age of five: a cohort study. <i>Scientific Reports</i> , 2022, 12, 1832.	3.3	2
3	Maternal Hypertensive Disorders in Pregnant Women With Systemic Lupus Erythematosus and Future Cardiovascular Outcomes. <i>Arthritis Care and Research</i> , 2021, 73, 574-579.	3.4	7
4	Pregnancy outcome in women with polycystic ovary syndrome in relation to second-trimester testosterone levels. <i>Reproductive BioMedicine Online</i> , 2021, 42, 217-225.	2.4	20
5	Preeclampsia and Increased Permeability Over the Blood-Brain Barrier: A Role of Vascular Endothelial Growth Receptor 2. <i>American Journal of Hypertension</i> , 2021, 34, 73-81.	2.0	20
6	Polycystic ovary syndrome and extremely preterm birth: A nationwide register-based study. <i>PLoS ONE</i> , 2021, 16, e0246743.	2.5	12
7	Women's childbirth experiences in the Swedish Post-term Induction Study (SWEPIIS): a multicentre, randomised, controlled trial. <i>BMJ Open</i> , 2021, 11, e042340.	1.9	10
8	Evidence of under-reporting of early-onset preeclampsia using register data. <i>Paediatric and Perinatal Epidemiology</i> , 2021, 35, 596-600.	1.7	4
9	Efficacy and safety of oral misoprostol vs transvaginal balloon catheter for labor induction: An observational study within the SWEdish Postterm Induction Study (SWEPIIS). <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 1463-1477.	2.8	5
10	Self-rated health before pregnancy and adverse birth outcomes in Sweden: A population-based register study. <i>Birth</i> , 2021, 48, 541-549.	2.2	3
11	Time matters—a Swedish cohort study of labor duration and risk of uterine rupture. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 1902-1909.	2.8	5
12	The association between postpartum hemorrhage and postpartum depression: A Swedish national register-based study. <i>PLoS ONE</i> , 2021, 16, e0255938.	2.5	5
13	Hypoxic ischemic encephalopathy in offspring of immigrant women in Sweden: A population-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 2285-2293.	2.8	7
14	COVID-19 in Pregnancy and Early Childhood (COPE): study protocol for a prospective, multicentre biobank, survey and database cohort study. <i>BMJ Open</i> , 2021, 11, e049376.	1.9	2
15	Reply to: Re: Risk of pre-eclampsia after gastric bypass: a matched cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2021, , .	2.3	0
16	Study protocol: establishment of a multicentre pre-eclampsia database and biobank in Sweden: GO PROVE and UP MOST, a prospective cohort study. <i>BMJ Open</i> , 2021, 11, e049559.	1.9	2
17	Study for Improving Maternal Pregnancy And Child ouTcomes (IMPACT): a study protocol for a Swedish prospective multicentre cohort study. <i>BMJ Open</i> , 2020, 10, e033851.	1.9	2
18	Induction of Labour at 41 Weeks Versus Expectant Management and Induction of Labour at 42 Weeks (SWEdish Post-Term Induction Study, SWEPIIS): Multicentre, Open Label, Randomised, Superiority Trial. <i>Obstetrical and Gynecological Survey</i> , 2020, 75, 207-209.	0.4	0

#	ARTICLE	IF	CITATIONS
19	Accelerated fetal growth in early pregnancy and risk of preterm birth: a prospective cohort study. <i>BMC Pregnancy and Childbirth</i> , 2020, 20, 764.	2.4	3
20	Pregnancy in metabolic healthy and unhealthy obese women. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020, 99, 1640-1648.	2.8	7
21	SSRI use during pregnancy and risk for postpartum haemorrhage: a national register-based cohort study in Sweden. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2020, 127, 1366-1373.	2.3	15
22	Factors associated with re-initiation of antidepressant treatment following discontinuation during pregnancy: a register-based cohort study. <i>Archives of Women's Mental Health</i> , 2020, 23, 709-717.	2.6	6
23	Placental location and risk of retained placenta in women with a previous cesarean section: A population-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020, 99, 1666-1673.	2.8	15
24	Epidural Analgesia During Childbirth and Postpartum Depressive Symptoms: A Population-Based Longitudinal Cohort Study. <i>Anesthesia and Analgesia</i> , 2020, 130, 615-624.	2.2	25
25	Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials. <i>PLoS Medicine</i> , 2020, 17, e1003436.	8.4	25
26	Neuroticism is associated with higher antenatal care utilization in obstetric low-risk women. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 470-478.	2.8	2
27	Plasma from women with preeclampsia impairs the blood brain barrier in an in vitro model. <i>Pregnancy Hypertension</i> , 2019, 17, S7.	1.4	1
28	Association of Attention-Deficit/Hyperactivity Disorder With Teenage Birth Among Women and Girls in Sweden. <i>JAMA Network Open</i> , 2019, 2, e1912463.	5.9	28
29	Delivery outcome after trial of labor in nulliparous women 40 years or older—A nationwide population-based study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 1195-1203.	2.8	9
30	Proton Pump Inhibitors and Preeclampsia Risk Among 157,720 Women. <i>Hypertension</i> , 2019, 73, 1097-1103.	2.7	22
31	Neighborhood deprivation and adverse perinatal outcomes in Sweden: A population-based register study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 1004-1013.	2.8	16
32	Placental location and pregnancy outcomes in nulliparous women: A population-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019, 98, 988-996.	2.8	18
33	Pregnancy and neonatal complications in women with polycystic ovary syndrome in relation to second-trimester anti-Müllerian hormone levels. <i>Reproductive BioMedicine Online</i> , 2019, 39, 141-148.	2.4	13
34	Postnatal growth in children born small for gestational age with and without smoking mother. <i>Pediatric Research</i> , 2019, 85, 961-966.	2.3	2
35	Induction of labour at 41 weeks versus expectant management and induction of labour at 42 weeks (SWEdish Post-term Induction Study, SWEPIS): multicentre, open label, randomised, superiority trial. <i>BMJ: British Medical Journal</i> , 2019, 367, l6131.	2.3	87
36	Clinical risk assessment in early pregnancy for preeclampsia in nulliparous women: A population based cohort study. <i>PLoS ONE</i> , 2019, 14, e0225716.	2.5	26

#	ARTICLE	IF	CITATIONS
37	Preterm Delivery Phenotypes in Systemic Lupus Erythematosus Pregnancies. <i>American Journal of Perinatology</i> , 2019, 36, 964-968.	1.4	9
38	Antepartum risk factors for moderate to severe neonatal hypoxic ischemic encephalopathy: a Swedish national cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 615-623.	2.8	27
39	Cerebral Osmolytes and Plasma Osmolality in Pregnancy and Preeclampsia: A Proton Magnetic Resonance Spectroscopy Study. <i>American Journal of Hypertension</i> , 2018, 31, 847-853.	2.0	10
40	The Swedish Pregnancy Register – for quality of care improvement and research. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 466-476.	2.8	101
41	Delineating the association between mode of delivery and postpartum depression symptoms: a longitudinal study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 301-311.	2.8	56
42	Opium use during pregnancy and infant size at birth: a cohort study. <i>BMC Pregnancy and Childbirth</i> , 2018, 18, 358.	2.4	8
43	Assessment of cerebral perfusion and edema in preeclampsia with intravoxel incoherent motion <scp>MRI</scp>. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 1212-1218.	2.8	13
44	Obstetric emergencies as antecedents to neonatal hypoxic ischemic encephalopathy, does parity matter?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2018, 97, 1396-1404.	2.8	14
45	Pregnancy Weight Gain Before Diagnosis and Risk of Preeclampsia. <i>Hypertension</i> , 2018, 72, 433-441.	2.7	53
46	Prenatal exposure to preeclampsia is associated with accelerated height gain in early childhood. <i>PLoS ONE</i> , 2018, 13, e0192514.	2.5	9
47	Chronic Hypertension in Women after Perinatal Exposure to Preeclampsia, Being Born Small for Gestational Age or Preterm. <i>Paediatric and Perinatal Epidemiology</i> , 2017, 31, 89-98.	1.7	9
48	Cerebral Magnesium Levels in Preeclampsia; A Phosphorus Magnetic Resonance Spectroscopy Study. <i>American Journal of Hypertension</i> , 2017, 30, 667-672.	2.0	15
49	Serum Pentraxin 3 is associated with signs of arterial alteration in women with preeclampsia. <i>International Journal of Cardiology</i> , 2017, 241, 417-422.	1.7	13
50	Early-onset Preeclampsia in Lupus Pregnancy. <i>Paediatric and Perinatal Epidemiology</i> , 2017, 31, 29-36.	1.7	36
51	Accelerated fetal growth in early pregnancy and risk of severe large-for-gestational-age and macrosomic infant: a cohort study in a low-risk population. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2017, 96, 1261-1268.	2.8	4
52	Born Small for Gestational Age and Poor School Performance – How Small Is Too Small?. <i>Hormone Research in Paediatrics</i> , 2017, 88, 215-223.	1.8	28
53	Association Between Pregnancy and Perinatal Outcomes Among Women With Epilepsy. <i>JAMA Neurology</i> , 2017, 74, 983.	9.0	90
54	Early second-trimester plasma levels of NT-proBNP in women who subsequently develop early-onset preeclampsia. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2017, 30, 2163-2165.	1.5	12

#	ARTICLE	IF	CITATIONS
55	Slow fetal growth between first and early second trimester ultrasound scans and risk of small for gestational age (SGA) birth. PLoS ONE, 2017, 12, e0184853.	2.5	7
56	Opium use during pregnancy and risk of preterm delivery: A population-based cohort study. PLoS ONE, 2017, 12, e0176588.	2.5	11
57	Induction of labor versus expectant management of large-for-gestational-age infants in nulliparous women. PLoS ONE, 2017, 12, e0180748.	2.5	15
58	Maternal overweight and obesity and risk of pre-eclampsia in women with type 1 diabetes or type 2 diabetes. Diabetologia, 2016, 59, 2099-2105.	6.3	48
59	High Maternal Body Mass Index in Early Pregnancy and Risks of Stillbirth and Infant Mortality—A Population-Based Sibling Study in Sweden. American Journal of Epidemiology, 2016, 184, 98-105.	3.4	25
60	The Impact of Small for Gestational Age Births in Early and Late Preeclamptic Pregnancies for Preeclampsia Recurrence: a Cohort Study of Successive Pregnancies in Sweden. Paediatric and Perinatal Epidemiology, 2016, 30, 563-570.	1.7	14
61	Study protocol of SWEPIs a Swedish multicentre register based randomised controlled trial to compare induction of labour at 41 completed gestational weeks versus expectant management and induction at 42 completed gestational weeks. BMC Pregnancy and Childbirth, 2016, 16, 49.	2.4	20
62	Prehypertension in Pregnancy and Risks of Small for Gestational Age Infant and Stillbirth. Hypertension, 2016, 67, 640-646.	2.7	49
63	Placental abruption and long-term maternal cardiovascular disease mortality: a population-based registry study in Norway and Sweden. European Journal of Epidemiology, 2016, 31, 501-511.	5.7	45
64	Delineating the Association between Heavy Postpartum Haemorrhage and Postpartum Depression. PLoS ONE, 2016, 11, e0144274.	2.5	49
65	Iodine deficiency in a study population of pregnant women in Sweden. Acta Obstetrica Et Gynecologica Scandinavica, 2015, 94, 1168-1174.	2.8	28
66	Consanguineous marriage, prepregnancy maternal characteristics and stillbirth risk: a population-based case-control study. Acta Obstetrica Et Gynecologica Scandinavica, 2015, 94, 1095-1101.	2.8	22
67	Does Use of Low-Molecular-Weight Heparin during Pregnancy Influence the Risk of Prolonged Labor: A Population-Based Cohort Study. PLoS ONE, 2015, 10, e0140422.	2.5	2
68	Maternal Snuff Use and Smoking and the Risk of Oral Cleft Malformations - A Population-Based Cohort Study. PLoS ONE, 2014, 9, e84715.	2.5	44
69	Plasma Levels of S100B in Preeclampsia and Association With Possible Central Nervous System Effects. American Journal of Hypertension, 2014, 27, 1105-1111.	2.0	17
70	Experience of fetal scalp blood sampling during labor. Acta Obstetrica Et Gynecologica Scandinavica, 2014, 93, 113-117.	2.8	3
71	Targeted Thyroid Testing During Pregnancy in Clinical Practice. Obstetrics and Gynecology, 2014, 124, 10-15.	2.4	23
72	Maternal Obesity and Risk of Preterm Delivery. Survey of Anesthesiology, 2014, 58, 70-71.	0.1	1

#	ARTICLE	IF	CITATIONS
73	Risk of placental dysfunction disorders after prior miscarriages: a population-based study. <i>American Journal of Obstetrics and Gynecology</i> , 2014, 211, 34.e1-34.e8.	1.3	34
74	Placental Expression of proBNP/NT-proBNP and Plasma Levels of NT-proBNP in Early- and Late-Onset Preeclampsia. <i>American Journal of Hypertension</i> , 2014, 27, 1225-1230.	2.0	21
75	Influence of Snuff and Smoking Habits in Early Pregnancy on Risks for Stillbirth and Early Neonatal Mortality. <i>Nicotine and Tobacco Research</i> , 2014, 16, 78-83.	2.6	51
76	Risk of retained placenta in women previously delivered by caesarean section: a population-based cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014, 121, 224-229.	2.3	33
77	Maternal Obesity and Risk of Preterm Delivery. <i>JAMA - Journal of the American Medical Association</i> , 2013, 309, 2362.	7.4	550
78	Elective induction of labor and the risk of cesarean section in low-risk parous women: a cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2013, 92, 198-203.	2.8	32
79	Thyroid Testing and Management of Hypothyroidism During Pregnancy: A Population-based Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 2687-2692.	3.6	34
80	Maternal Obesity and Risk of Preterm Delivery. <i>Obstetrical and Gynecological Survey</i> , 2013, 68, 731-732.	0.4	0
81	Psychosocial Stress Related to the Loss of a Close Relative the Year Before or During Pregnancy and Risk of Preeclampsia. <i>Hypertension</i> , 2013, 62, 183-189.	2.7	37
82	Hyperemesis gravidarum and risks of placental dysfunction disorders: a population-based cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2013, 120, 541-547.	2.3	86
83	Evaluation of the Discrepancy Between pH and Lactate in Combined Fetal Scalp Blood Sampling. <i>Obstetric Anesthesia Digest</i> , 2012, 32, 177-178.	0.1	0
84	From infancy to pregnancy: birth weight, body mass index, and the risk of gestational diabetes. <i>European Journal of Epidemiology</i> , 2012, 27, 799-805.	5.7	16
85	Interpregnancy Change in Smoking Habits and Risk of Preeclampsia: A Population-Based Study. <i>American Journal of Hypertension</i> , 2012, 25, 372-378.	2.0	16
86	Labour dystocia—risk of recurrence and instrumental delivery in following labour—a population-based cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012, 119, 1648-1656.	2.3	22
87	Prediction of Preeclampsia by Combining Serum Histidine-Rich Glycoprotein and Uterine Artery Doppler. <i>American Journal of Hypertension</i> , 2012, 25, 1305-10.	2.0	12
88	Maternal Body Mass Index, Height, and Risks of Preeclampsia. <i>American Journal of Hypertension</i> , 2012, 25, 120-125.	2.0	110
89	Influence of smoking and snuff cessation on risk of preterm birth. <i>European Journal of Epidemiology</i> , 2012, 27, 297-304.	5.7	58
90	Evaluation of the discrepancy between pH and lactate in combined fetal scalp blood sampling. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2011, 90, 1088-1093.	2.8	10

#	ARTICLE	IF	CITATIONS
91	Previous preeclampsia and risks of adverse outcomes in subsequent nonpreeclamptic pregnancies. American Journal of Obstetrics and Gynecology, 2011, 204, 148.e1-148.e6.	1.3	47
92	Recurrence of placental dysfunction disorders across generations. American Journal of Obstetrics and Gynecology, 2011, 205, 454.e1-454.e8.	1.3	13
93	Relationship of Maternal Snuff Use and Cigarette Smoking With Neonatal Apnea. Pediatrics, 2011, 128, 503-509.	2.1	40
94	Tobacco Use During Pregnancy and Preeclampsia Risk. Hypertension, 2010, 55, 1254-1259.	2.7	154
95	Maternal Use of Swedish Snuff (Snus) and Risk of Stillbirth. Epidemiology, 2010, 21, 772-778.	2.7	66
96	Fibrinogen and histidine-rich glycoprotein in early-onset preeclampsia. Acta Obstetrica Et Gynecologica Scandinavica, 2010, 89, 131-139.	2.8	24
97	Increased Circulating Levels of the Antiangiogenic Factor Endostatin in Early-Onset But Not Late-Onset Preeclampsia. Reproductive Sciences, 2009, 16, 995-1000.	2.5	18
98	Angiopoietin-1/Angiopoietin-2 Ratio for Prediction of Preeclampsia. American Journal of Hypertension, 2009, 22, 891-895.	2.0	49
99	Early postpartum changes in circulating pro- and anti-angiogenic factors in early-onset and late-onset preeclampsia. Acta Obstetrica Et Gynecologica Scandinavica, 2008, 87, 146-153.	2.8	27
100	Placental Growth Factor and Soluble FMS-Like Tyrosine Kinase-1 in Early-Onset and Late-Onset Preeclampsia. Obstetrics and Gynecology, 2007, 109, 1368-1374.	2.4	172
101	The risk of maternal ischaemic heart disease after gestational hypertensive disease. BJOG: an International Journal of Obstetrics and Gynaecology, 2005, 112, 1486-1491.	2.3	202