

Florian Herse

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6848175/florian-herse-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

79
papers

2,737
citations

31
h-index

50
g-index

81
ext. papers

3,220
ext. citations

6.4
avg, IF

4.46
L-index

#	Paper	IF	Citations
79	Kidney Injury Caused by Preeclamptic Pregnancy Recovers Postpartum in a Transgenic Rat Model. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
78	Maternal Angiotensin Increases Placental Leptin in Early Gestation via an Alternative Renin-Angiotensin System Pathway: Suggesting a Link to Preeclampsia. <i>Hypertension</i> , 2021 , 77, 1723-1736	8.5	5
77	Preexisting hypertension and pregnancy-induced hypertension reveal molecular differences in placental proteome in rodents. <i>Physiological Genomics</i> , 2021 , 53, 259-268	3.6	1
76	Intrauterine Exposure to Diabetic Milieu Does Not Induce Diabetes and Obesity in Male Adulthood in a Novel Rat Model. <i>Hypertension</i> , 2021 , 77, 202-215	8.5	3
75	Diabetic pregnancy as a novel risk factor for cardiac dysfunction in the offspring-the heart as a target for fetal programming in rats. <i>Diabetologia</i> , 2021 , 64, 2829-2842	10.3	0
74	Circulating Maternal sFLT1 (Soluble fms-Like Tyrosine Kinase-1) Is Sufficient to Impair Spiral Arterial Remodeling in a Preeclampsia Mouse Model. <i>Hypertension</i> , 2021 , 78, 1067-1079	8.5	4
73	RNA interference therapeutics targeting angiotensinogen ameliorate preeclamptic phenotype in rodent models. <i>Journal of Clinical Investigation</i> , 2020 , 130, 2928-2942	15.9	12
72	Statins Reverse Postpartum Cardiovascular Dysfunction in a Rat Model of Preeclampsia. <i>Hypertension</i> , 2020 , 75, 202-210	8.5	16
71	High-sensitivity cardiac troponin I in women with a history of early-onset preeclampsia. <i>Journal of Hypertension</i> , 2020 , 38, 1948-1954	1.9	2
70	Effects of empagliflozin and target-organ damage in a novel rodent model of heart failure induced by combined hypertension and diabetes. <i>Scientific Reports</i> , 2020 , 10, 14061	4.9	3
69	Aldosterone, Salt, and Potassium Intakes as Predictors of Pregnancy Outcome, Including Preeclampsia. <i>Hypertension</i> , 2019 , 74, 391-398	8.5	12
68	Cardiovascular Programming During and After Diabetic Pregnancy: Role of Placental Dysfunction and IUGR. <i>Frontiers in Endocrinology</i> , 2019 , 10, 215	5.7	5
67	The Pediatric Cell Atlas: Defining the Growth Phase of Human Development at Single-Cell Resolution. <i>Developmental Cell</i> , 2019 , 49, 10-29	10.2	39
66	Human sFLT1 Leads to Severe Changes in Placental Differentiation and Vascularization in a Transgenic hsFLT1/rtTA FGR Mouse Model. <i>Frontiers in Endocrinology</i> , 2019 , 10, 165	5.7	12
65	Placental CX3CL1 is Deregulated by Angiotensin II and Contributes to a Pro-Inflammatory Trophoblast-Monocyte Interaction. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	8
64	Regulatory antibodies against GPCR in women ten years after early-onset preeclampsia. <i>Frontiers in Bioscience - Landmark</i> , 2019 , 24, 1462-1476	2.8	1
63	Interleukin-4 supplementation improves the pathophysiology of hypertension in response to placental ischemia in RUPP rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2019 , 316, R165-R171	3.2	13

62	Diabetes Mellitus in Pregnancy Leads to Growth Restriction and Epigenetic Modification of the Gene in Rat Fetuses. <i>Hypertension</i> , 2018 , 71, 911-920	8.5	17
61	Continuous Blood Glucose Monitoring Reveals Enormous Circadian Variations in Pregnant Diabetic Rats. <i>Frontiers in Endocrinology</i> , 2018 , 9, 271	5.7	3
60	Nitric oxide-sensitive guanylyl cyclase stimulation improves experimental heart failure with preserved ejection fraction. <i>JCI Insight</i> , 2018 , 3,	9.9	17
59	Downregulation of p53 drives autophagy during human trophoblast differentiation. <i>Cellular and Molecular Life Sciences</i> , 2018 , 75, 1839-1855	10.3	21
58	Modeling Superimposed Preeclampsia Using Ang II (Angiotensin II) Infusion in Pregnant Stroke-Prone Spontaneously Hypertensive Rats. <i>Hypertension</i> , 2018 , 72, 208-218	8.5	8
57	Soluble B7-H4 blood serum levels are elevated in women at high risk for preeclampsia in the first trimester, as well as in patients with confirmed preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2018 , 80, e12988	3.8	7
56	Placental DAPK1 and autophagy marker LC3B-II are dysregulated by TNF- α in a gestational age-dependent manner. <i>Histochemistry and Cell Biology</i> , 2017 , 147, 695-705	2.4	14
55	Placental expression of sFlt-1 and PlGF in early preeclampsia vs. early IUGR vs. age-matched healthy pregnancies. <i>Hypertension in Pregnancy</i> , 2017 , 36, 151-160	2	27
54	Vitamin D supplementation reduces some AT1-AA-induced downstream targets implicated in preeclampsia including hypertension. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2017 , 312, R125-R131	3.2	13
53	Increased placental sFlt-1 but unchanged PlGF expression in late-onset preeclampsia. <i>Hypertension in Pregnancy</i> , 2017 , 36, 175-185	2	12
52	Placental-Specific Overexpression of sFlt-1 Alters Trophoblast Differentiation and Nutrient Transporter Expression in an IUGR Mouse Model. <i>Journal of Cellular Biochemistry</i> , 2017 , 118, 1316-1329	4.7	23
51	Soluble (pro)renin receptor in preeclampsia and diabetic pregnancies. <i>Journal of the American Society of Hypertension</i> , 2017 , 11, 644-652		9
50	Disturbed Placental Imprinting in Preeclampsia Leads to Altered Expression of DLX5, a Human-Specific Early Trophoblast Marker. <i>Circulation</i> , 2017 , 136, 1824-1839	16.7	31
49	Proliferation of endogenous regulatory T cells improve the pathophysiology associated with placental ischaemia of pregnancy. <i>American Journal of Reproductive Immunology</i> , 2017 , 78, e12724	3.8	18
48	Natural Killer Cell Reduction and Uteroplacental Vasculopathy. <i>Hypertension</i> , 2016 , 68, 964-73	8.5	12
47	Relaxin Treatment in an Ang-II-Based Transgenic Preeclamptic-Rat Model. <i>PLoS ONE</i> , 2016 , 11, e0150743	3.7	7
46	Convergent Evolution Within CEA Gene Families in Mammals: Hints for Species-Specific Selection Pressures 2016 , 37-53		0
45	Reduced uterine perfusion pressure T-helper 17 cells cause pathophysiology associated with preeclampsia during pregnancy. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2016 , 311, R1192-R1199	3.2	36

44	Vitamin D supplementation improves pathophysiology in a rat model of preeclampsia. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2016 , 310, R346-54	3.2	24
43	CD74-Downregulation of Placental Macrophage-Trophoblastic Interactions in Preeclampsia. <i>Circulation Research</i> , 2016 , 119, 55-68	15.7	34
42	Vitamin D depletion does not affect key aspects of the preeclamptic phenotype in a transgenic rodent model for preeclampsia. <i>Journal of the American Society of Hypertension</i> , 2016 , 10, 597-607.e1		3
41	Placental fractalkine mediates adhesion of THP-1 monocytes to villous trophoblast. <i>Histochemistry and Cell Biology</i> , 2015 , 143, 565-74	2.4	14
40	Placental fractalkine is up-regulated in severe early-onset preeclampsia. <i>American Journal of Pathology</i> , 2015 , 185, 1334-43	5.8	29
39	Regulatory T cells ameliorate intrauterine growth retardation in a transgenic rat model for preeclampsia. <i>Hypertension</i> , 2015 , 65, 1298-306	8.5	25
38	Increased apoptosis, altered oxygen signaling, and antioxidant defenses in first-trimester pregnancies with high-resistance uterine artery blood flow. <i>American Journal of Pathology</i> , 2015 , 185, 2731-41	5.8	36
37	Blockade of CD40 ligand for intercellular communication reduces hypertension, placental oxidative stress, and AT1-AA in response to adoptive transfer of CD4+ T lymphocytes from RUPP rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015 , 309, R1243-50	3.2	11
36	IL-10 supplementation increases Tregs and decreases hypertension in the RUPP rat model of preeclampsia. <i>Hypertension in Pregnancy</i> , 2015 , 34, 291-306	2	55
35	An increased population of regulatory T cells improves the pathophysiology of placental ischemia in a rat model of preeclampsia. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015 , 309, R884-91	3.2	52
34	Administration of interleukin-17 soluble receptor C suppresses TH17 cells, oxidative stress, and hypertension in response to placental ischemia during pregnancy. <i>Hypertension</i> , 2013 , 62, 1068-73	8.5	78
33	Amyloid- β peptides activate β -adrenergic cardiovascular receptors. <i>Hypertension</i> , 2013 , 62, 966-72	8.5	17
32	Endothelin-1, oxidative stress, and endogenous angiotensin II: mechanisms of angiotensin II type I receptor autoantibody-enhanced renal and blood pressure response during pregnancy. <i>Hypertension</i> , 2013 , 62, 886-92	8.5	65
31	Interfering with Gal-1-mediated angiogenesis contributes to the pathogenesis of preeclampsia. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 11451-6	11.5	71
30	Angiotensin II type 1 receptor autoantibody (AT1-AA)-mediated pregnancy hypertension. <i>American Journal of Reproductive Immunology</i> , 2013 , 69, 413-8	3.8	69
29	CD4 T Cells Play a Critical Role in Mediating Hypertension in Response to Placental Ischemia. <i>Journal of Hypertension: Open Access</i> , 2013 , 2,		20
28	CD19+CD5+ cells as indicators of preeclampsia. <i>Hypertension</i> , 2012 , 59, 861-8	8.5	87
27	IL-17-mediated oxidative stress is an important stimulator of AT1-AA and hypertension during pregnancy. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2012 , 303, R353-8	3.2	92

26	Cardiovascular risk markers in pregnancies complicated by diabetes mellitus or preeclampsia. <i>Pregnancy Hypertension</i> , 2012 , 2, 403-10	2.6	8
25	Activating autoantibodies to the angiotensin II type I receptor play an important role in mediating hypertension in response to adoptive transfer of CD4+ T lymphocytes from placental ischemic rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2012 , 302, R1197-201	3.2	58
24	Cytochrome P450 subfamily 2J polypeptide 2 expression and circulating epoxyeicosatrienoic metabolites in preeclampsia. <i>Circulation</i> , 2012 , 126, 2990-9	16.7	48
23	Cardiovascular biomarker midregional proatrial natriuretic peptide during and after preeclamptic pregnancies. <i>Hypertension</i> , 2012 , 59, 395-401	8.5	18
22	Hypertension in response to AT1-AA: role of reactive oxygen species in pregnancy-induced hypertension. <i>American Journal of Hypertension</i> , 2011 , 24, 835-40	2.3	60
21	Hypertension in response to placental ischemia during pregnancy: role of B lymphocytes. <i>Hypertension</i> , 2011 , 57, 865-71	8.5	92
20	Angiotensin II type 1 receptor antibodies and increased angiotensin II sensitivity in pregnant rats. <i>Hypertension</i> , 2011 , 58, 77-84	8.5	107
19	Heparin strongly induces soluble fms-like tyrosine kinase 1 release in vivo and in vitro--brief report. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 2972-4	9.4	45
18	Adipose tissue-derived soluble fms-like tyrosine kinase 1 is an obesity-relevant endogenous paracrine adipokine. <i>Hypertension</i> , 2011 , 58, 37-42	8.5	20
17	Alpha1A-adrenergic receptor-directed autoimmunity induces left ventricular damage and diastolic dysfunction in rats. <i>PLoS ONE</i> , 2010 , 5, e9409	3.7	11
16	Inhibition of trophoblast-induced spiral artery remodeling reduces placental perfusion in rat pregnancy. <i>Hypertension</i> , 2010 , 56, 304-10	8.5	60
15	Increased angiotensin II in the mesometrial triangle of a transgenic rat model of preeclampsia. <i>Hypertension</i> , 2010 , 55, 562-6	8.5	21
14	Effects of circulating and local uteroplacental angiotensin II in rat pregnancy. <i>Hypertension</i> , 2010 , 56, 311-8	8.5	54
13	Prevalence of agonistic autoantibodies against the angiotensin II type 1 receptor and soluble fms-like tyrosine kinase 1 in a gestational age-matched case study. <i>Hypertension</i> , 2009 , 53, 393-8	8.5	78
12	A recently evolved novel trophoblast-enriched secreted form of fms-like tyrosine kinase-1 variant is up-regulated in hypoxia and preeclampsia. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 2524-30	5.6	64
11	Circulating and placental growth-differentiation factor 15 in preeclampsia and in pregnancy complicated by diabetes mellitus. <i>Hypertension</i> , 2009 , 54, 106-12	8.5	37
10	Circulating and uteroplacental adipocytokine concentrations in preeclampsia. <i>Reproductive Sciences</i> , 2009 , 16, 584-90	3	27
9	Dietary n-3 polyunsaturated fatty acids and direct renin inhibition improve electrical remodeling in a model of high human renin hypertension. <i>Hypertension</i> , 2008 , 51, 540-6	8.5	79

8	Trophoblasts reduce the vascular smooth muscle cell proatherogenic response. <i>Hypertension</i> , 2008 , 51, 554-9	8.5	26
7	Uterine vascular function in a transgenic preeclampsia rat model. <i>Hypertension</i> , 2008 , 51, 547-53	8.5	69
6	Autoantibodies to the angiotensin type I receptor in response to placental ischemia and tumor necrosis factor alpha in pregnant rats. <i>Hypertension</i> , 2008 , 52, 1168-72	8.5	133
5	Potential relevance of alpha(1)-adrenergic receptor autoantibodies in refractory hypertension. <i>PLoS ONE</i> , 2008 , 3, e3742	3.7	74
4	AT1-receptor autoantibodies and uteroplacental RAS in pregnancy and pre-eclampsia. <i>Journal of Molecular Medicine</i> , 2008 , 86, 697-703	5.5	57
3	Dysregulation of the circulating and tissue-based renin-angiotensin system in preeclampsia. <i>Hypertension</i> , 2007 , 49, 604-11	8.5	187
2	Agonistic angiotensin II type 1 receptor autoantibodies in postpartum women with a history of preeclampsia. <i>Hypertension</i> , 2007 , 49, 612-7	8.5	98
1	Expression of the protein phosphatase 1 inhibitor KEPI is downregulated in breast cancer cell lines and tissues and involved in the regulation of the tumor suppressor EGR1 via the MEK-ERK pathway. <i>Biological Chemistry</i> , 2007 , 388, 489-95	4.5	12