

# Kenneth F Raffa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/684878/publications.pdf>

Version: 2024-02-01

214  
papers

15,021  
citations

17405

63  
h-index

22102

113  
g-index

217  
all docs

217  
docs citations

217  
times ranked

10826  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Numbers matter: how irruptive bark beetles initiate transition to self-sustaining behavior during landscape-altering outbreaks. <i>Oecologia</i> , 2022, 198, 681-698.  | 0.9 | 9         |
| 2  | Spread rates do not necessarily predict outbreak dynamics in a broadly distributed invasive insect. <i>Forest Ecology and Management</i> , 2022, 520, 120357.   | 1.4 | 3         |
| 3  | Root Secondary Metabolites in <i>Populus tremuloides</i> : Effects of Simulated Climate Warming, Defoliation, and Genotype. <i>Journal of Chemical Ecology</i> , 2021, 47, 313-321.   | 0.9 | 9         |
| 4  | Bark Beetle Outbreaks in Europe: State of Knowledge and Ways Forward for Management. <i>Current Forestry Reports</i> , 2021, 7, 138-165.  | 3.4 | 133       |
| 5  | Predicting non-native insect impact: focusing on the trees to see the forest. <i>Biological Invasions</i> , 2021, 23, 3921-3936.  | 1.2 | 5         |
| 6  | Growth and defense characteristics of whitebark pine ( <i>Pinus albicaulis</i> ) and lodgepole pine ( <i>Pinus</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Montana, USA. <i>Forest Ecology and Management</i> , 2021, 493, 119286.     | 1.4 | 5         |
| 7  | Climate-induced outbreaks in high-elevation pines are driven primarily by immigration of bark beetles from historical hosts. <i>Global Change Biology</i> , 2021, 27, 5786-5805.  | 4.2 | 5         |
| 8  | Combined drought and bark beetle attacks deplete non-structural carbohydrates and promote death of mature pine trees. <i>Plant, Cell and Environment</i> , 2021, 44, 3866-3881.   | 2.8 | 16        |
| 9  | Physical contact, volatiles, and acoustic signals contribute to monogamy in an invasive aggregating bark beetle. <i>Insect Science</i> , 2020, 27, 1285-1297.   | 1.5 | 4         |
| 10 | Why do entomologists and plant pathologists approach trophic relationships so differently? Identifying biological distinctions to foster synthesis. <i>New Phytologist</i> , 2020, 225, 609-620.                                      | 3.5 | 14        |
| 11 | Tree defence and bark beetles in a drying world: carbon partitioning, functioning and modelling. <i>New Phytologist</i> , 2020, 225, 26-36.   | 3.5 | 144       |
| 12 | Phenological responses to prior-season defoliation and soil-nutrient availability vary among early- and late-flushing aspen ( <i>Populus tremuloides</i> Michx.) genotypes. <i>Forest Ecology and Management</i> , 2020, 458, 117771. | 1.4 | 5         |
| 13 | Relationships between conifer constitutive and inducible defenses against bark beetles change across levels of biological and ecological scale. <i>Oikos</i> , 2020, 129, 1093-1107.  | 1.2 | 12        |
| 14 | Evolutionary history predicts high-impact invasions by herbivorous insects. <i>Ecology and Evolution</i> , 2019, 9, 12216-12230.  | 0.8 | 28        |
| 15 | Drought-Mediated Changes in Tree Physiological Processes Weaken Tree Defenses to Bark Beetle Attack. <i>Journal of Chemical Ecology</i> , 2019, 45, 888-900.  | 0.9 | 67        |
| 16 | Anatomical defences against bark beetles relate to degree of historical exposure between species and are allocated independently of chemical defences within trees. <i>Plant, Cell and Environment</i> , 2019, 42, 633-646.           | 2.8 | 27        |
| 17 | Pine Engravers Carry Bacterial Communities Whose Members Reduce Concentrations of Host Monoterpenes With Variable Degrees of Redundancy, Specificity, and Capability. <i>Environmental Entomology</i> , 2018, 47, 638-645.            | 0.7 | 28        |
| 18 | Predators and competitors of the mountain pine beetle <i>Dendroctonus ponderosae</i> (Coleoptera:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 <i>Forest Entomology</i> , 2018, 20, 402-413.   | 0.7 | 2         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Genetic variation in aspen phytochemical patterns structures windows of opportunity for gypsy moth larvae. <i>Oecologia</i> , 2018, 187, 471-482.   | 0.9 | 18        |
| 20 | Strategic Development of Tree Resistance Against Forest Pathogen and Insect Invasions in Defense-Free Space. <i>Frontiers in Ecology and Evolution</i> , 2018, 6, .   | 1.1 | 31        |
| 21 | Seasonal and Regional Distributions, Degree-Day Models, and Phoresy Rates of the Major Sap Beetle (Coleoptera: Nitidulidae) Vectors of the Oak Wilt Fungus, <i>Bretziella fagacearum</i> , in Wisconsin. <i>Environmental Entomology</i> , 2018, 47, 1152-1164.             | 0.7 | 13        |
| 22 | Sound-Triggered Production of Antiaggregation Pheromone Limits Overcrowding of <i>Dendroctonus valens</i> Attacking Pine Trees. <i>Chemical Senses</i> , 2017, 42, bjw102.  | 1.1 | 9         |
| 23 | Gallery and acoustic traits related to female body size mediate male mate choice in a bark beetle. <i>Animal Behaviour</i> , 2017, 125, 41-50.  | 0.8 | 11        |
| 24 | Recent and future climate suitability for whitebark pine mortality from mountain pine beetles varies across the western US. <i>Forest Ecology and Management</i> , 2017, 399, 132-142.  | 1.4 | 24        |
| 25 | Defence syndromes in lodgepole "whitebark pine ecosystems relate to degree of historical exposure to mountain pine beetles. <i>Plant, Cell and Environment</i> , 2017, 40, 1791-1806.   | 2.8 | 61        |
| 26 | Spatial and temporal components of induced plant responses in the context of herbivore life history and impact on host. <i>Functional Ecology</i> , 2017, 31, 2034-2050.  | 1.7 | 23        |
| 27 | Bacterial Communities Associated With the Pine Wilt Disease Vector <i>Monochamus alternatus</i> (Coleoptera: Cerambycidae) During Different Larval Instars. <i>Journal of Insect Science</i> , 2017, 17, .  | 0.6 | 7         |
| 28 | Supercooling points of diapausing forest tent caterpillar (Lepidoptera: Lasiocampidae) eggs. <i>Canadian Entomologist</i> , 2016, 148, 512-519.   | 0.4 | 8         |
| 29 | Behaviours of phoretic mites (Acari) associated with <i>Ips pini</i> and <i>Ips grandicollis</i> (Coleoptera: Curculionidae) during host-tree colonization. <i>Agricultural and Forest Entomology</i> , 2016, 18, 108-118.  | 0.7 | 9         |
| 30 | Climate influences on whitebark pine mortality from mountain pine beetle in the Greater Yellowstone Ecosystem. <i>Ecological Applications</i> , 2016, 26, 2507-2524.  | 1.8 | 66        |
| 31 | Spatial variability in tree regeneration after wildfire delays and dampens future bark beetle outbreaks. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 13075-13080.   | 3.3 | 65        |
| 32 | Oviposition and feeding on red pine by jack pine budworm at a previously unrecorded scale. <i>Agricultural and Forest Entomology</i> , 2016, 18, 214-222.   | 0.7 | 1         |
| 33 | Evaluation of tree mortality and parasitoid recoveries on the contiguous western invasion front of emerald ash borer. <i>Agricultural and Forest Entomology</i> , 2016, 18, 327-339.  | 0.7 | 6         |
| 34 | Interactions between Bacteria And Aspen Defense Chemicals at the Phyllosphere "Herbivore Interface. <i>Journal of Chemical Ecology</i> , 2016, 42, 193-201.   | 0.9 | 39        |
| 35 | Contributions by Host Trees and Insect Activity to Bacterial Communities in <i>Dendroctonus valens</i> (Coleoptera: Curculionidae) Galleries, and Their High Overlap With Other Microbial Assemblages of Bark Beetles. <i>Environmental Entomology</i> , 2016, 45, 348-356. | 0.7 | 23        |
| 36 | Rapid Induction of Multiple Terpenoid Groups by Ponderosa Pine in Response to Bark Beetle-Associated Fungi. <i>Journal of Chemical Ecology</i> , 2016, 42, 1-12.  | 0.9 | 76        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Effects of winter temperatures, spring degree-day accumulation, and insect population source on phenological synchrony between forest tent caterpillar and host trees. <i>Forest Ecology and Management</i> , 2016, 362, 241-250.            | 1.4 | 50        |
| 38 | Structure of Phoretic Mite Assemblages Across Subcortical Beetle Species at a Regional Scale. <i>Environmental Entomology</i> , 2016, 45, 53-65.   | 0.7 | 13        |
| 39 | Evolution of High Cellulolytic Activity in Symbiotic <i>Streptomyces</i> through Selection of Expanded Gene Content and Coordinated Gene Expression. <i>PLoS Biology</i> , 2016, 14, e1002475.   | 2.6 | 68        |
| 40 | Mountain Pine Beetle Dynamics and Reproductive Success in Post-Fire Lodgepole and Ponderosa Pine Forests in Northeastern Utah. <i>PLoS ONE</i> , 2016, 11, e0164738.   | 1.1 | 11        |
| 41 | Tree response and mountain pine beetle attack preference, reproduction and emergence timing in mixed whitebark and lodgepole pine stands. <i>Agricultural and Forest Entomology</i> , 2015, 17, 421-432.                                     | 0.7 | 59        |
| 42 | Tree mortality from drought, insects, and their interactions in a changing climate. <i>New Phytologist</i> , 2015, 208, 674-683.   | 3.5 | 641       |
| 43 | Do Phoretic Mites Influence the Reproductive Success of <i>Ips grandicollis</i> (Coleoptera: Curculionidae)? <i>Environmental Entomology</i> , 2015, 44, 1498-1511.  | 0.7 | 7         |
| 44 | Experimental climate warming alters aspen and birch phytochemistry and performance traits for an outbreak insect herbivore. <i>Global Change Biology</i> , 2015, 21, 2698-2710.  | 4.2 | 69        |
| 45 | Economics and Politics of Bark Beetles. , 2015, , 585-613.   |     | 43        |
| 46 | Natural History and Ecology of Bark Beetles. , 2015, , 1-40.   |     | 105       |
| 47 | Foliar bacterial communities of trembling aspen in a common garden. <i>Canadian Journal of Microbiology</i> , 2015, 61, 143-149.   | 0.8 | 10        |
| 48 | Evaluating Predators and Competitors in Wisconsin Red Pine Forests for Attraction to Mountain Pine Beetle Pheromones for Anticipatory Biological Control. <i>Environmental Entomology</i> , 2015, 44, 1161-1171.                             | 0.7 | 9         |
| 49 | Contrasting Patterns of Diterpene Acid Induction by Red Pine and White Spruce to Simulated Bark Beetle Attack, and Interspecific Differences in Sensitivity Among Fungal Associates. <i>Journal of Chemical Ecology</i> , 2015, 41, 524-532. | 0.9 | 15        |
| 50 | Bacteria influence mountain pine beetle brood development through interactions with symbiotic and antagonistic fungi: implications for climate-driven host range expansion. <i>Oecologia</i> , 2015, 179, 467-485.                           | 0.9 | 39        |
| 51 | Aspen Defense Chemicals Influence Midgut Bacterial Community Composition of Gypsy Moth. <i>Journal of Chemical Ecology</i> , 2015, 41, 75-84.  | 0.9 | 50        |
| 52 | New Insights into the Consequences of Post-Windthrow Salvage Logging Revealed by Functional Structure of Saproxylous Beetles Assemblages. <i>PLoS ONE</i> , 2014, 9, e101757.  | 1.1 | 62        |
| 53 | Effects of an invasive herbivore at the single plant scale do not extend to population-scale seedling dynamics. <i>Canadian Journal of Forest Research</i> , 2014, 44, 8-16.   | 0.8 | 6         |
| 54 | Convergent Bacterial Microbiotas in the Fungal Agricultural Systems of Insects. <i>MBio</i> , 2014, 5, e02077.   | 1.8 | 96        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Prevalence of <i>Borrelia burgdorferi</i> and <i>Anaplasma phagocytophilum</i> in <i>Ixodes scapularis</i> (Acari: Ixodidae) Nymphs Collected in Managed Red Pine Forests in Wisconsin. <i>Journal of Medical Entomology</i> , 2014, 51, 694-701.                    | 0.9 | 35        |
| 56 | Populations of uncultivated American cranberry in sphagnum bog communities harbor novel assemblages of Actinobacteria with antifungal properties. <i>Botany</i> , 2014, 92, 589-595.   | 0.5 | 8         |
| 57 | Acquisition and Structuring of Midgut Bacterial Communities in Gypsy Moth (Lepidoptera: Erebidae) Larvae. <i>Environmental Entomology</i> , 2014, 43, 595-604.   | 0.7 | 106       |
| 58 | Terpenes Tell Different Tales at Different Scales: Glimpses into the Chemical Ecology of Conifer - Bark Beetle - Microbial Interactions. <i>Journal of Chemical Ecology</i> , 2014, 40, 1-20.  | 0.9 | 94        |
| 59 | Plant-associated bacteria degrade defense chemicals and reduce their adverse effects on an insect defoliator. <i>Oecologia</i> , 2014, 175, 901-910.   | 0.9 | 106       |
| 60 | Cellulolytic <i>Streptomyces</i> Strains Associated with Herbivorous Insects Share a Phylogenetically Linked Capacity To Degrade Lignocellulose. <i>Applied and Environmental Microbiology</i> , 2014, 80, 4692-4701.  | 1.4 | 70        |
| 61 | Simulated climate warming alters phenological synchrony between an outbreak insect herbivore and host trees. <i>Oecologia</i> , 2014, 175, 1041-1049.  | 0.9 | 92        |
| 62 | A Tale of Convergence. <i>Journal of Chemical Ecology</i> , 2014, 40, 415-416.   | 0.9 | 0         |
| 63 | Influence of Diet and Density on Laboratory Cannibalism Behaviors in Gypsy Moth Larvae ( <i>Lymantria</i> ) Tj ETQq1 1 0.784314 rgBT /Ove<br>0.4 6   | 0.7 | 181       |
| 64 | Responses of two parasitoids, the exotic <i>Spathius agrili</i> Yang and the native <i>Spathius floridanus</i> Ashmead, to volatile cues associated with the emerald ash borer, <i>Agrilus planipennis</i> Fairmaire. <i>Biological Control</i> , 2014, 79, 110-117. | 1.4 | 15        |
| 65 | Bacteria Associated with a Tree-Killing Insect Reduce Concentrations of Plant Defense Compounds. <i>Journal of Chemical Ecology</i> , 2013, 39, 1003-1006.   | 0.9 | 227       |
| 66 | Minimization of chloroplast contamination in 16S rRNA gene pyrosequencing of insect herbivore bacterial communities. <i>Journal of Microbiological Methods</i> , 2013, 95, 149-155.  | 0.7 | 181       |
| 67 | Using delimiting surveys to characterize the spatiotemporal dynamics facilitates the management of an invasive non-native insect. <i>Population Ecology</i> , 2013, 55, 545-555.   | 0.7 | 14        |
| 68 | Dispersal and edge behaviour of bark beetles and predators inhabiting red pine plantations. <i>Agricultural and Forest Entomology</i> , 2013, 15, 1-11.  | 0.7 | 30        |
| 69 | Belowground herbivory in red pine stands initiates a cascade that increases abundance of Lyme disease vectors. <i>Forest Ecology and Management</i> , 2013, 302, 354-362.  | 1.4 | 9         |
| 70 | Mites Phoretic on <i>Ips pini</i> (Coleoptera: Curculionidae: Scolytinae) in Wisconsin Red Pine Stands. <i>Annals of the Entomological Society of America</i> , 2013, 106, 204-213.  | 1.3 | 15        |
| 71 | Mountain Pine Beetles Colonizing Historical and Native Host Trees Are Associated with a Bacterial Community Highly Enriched in Genes Contributing to Terpene Metabolism. <i>Applied and Environmental Microbiology</i> , 2013, 79, 3468-3475.                        | 1.4 | 236       |
| 72 | Temperature-driven range expansion of an irruptive insect heightened by weakly coevolved plant defenses. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 2193-2198.  | 3.3 | 169       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | Trap Lure Blend of Pine Volatiles and Bark Beetle Pheromones for <i>Monochamus</i> spp. (Coleoptera: Cerambycidae) in Pine Forests of Canada and the United States. <i>Journal of Economic Entomology</i> , 2013, 106, 1684-1692. | 0.8 | 30        |
| 74 | Wildfire provides refuge from local extinction but is an unlikely driver of outbreaks by mountain pine beetle. <i>Ecological Monographs</i> , 2012, 82, 69-84.  | 2.4 | 47        |
| 75 | Consequences of Climate Warming and Altered Precipitation Patterns for Plant-Insect and Multitrophic Interactions. <i>Plant Physiology</i> , 2012, 160, 1719-1727.  | 2.3 | 279       |
| 76 | Effects of biotic disturbances on forest carbon cycling in the United States and Canada. <i>Global Change Biology</i> , 2012, 18, 7-34.   | 4.2 | 418       |
| 77 | Variable host phenology does not pose a barrier to invasive weevils in a northern hardwood forest. <i>Agricultural and Forest Entomology</i> , 2012, 14, 276-285.   | 0.7 | 8         |
| 78 | What explains landscape patterns of tree mortality caused by bark beetle outbreaks in Greater Yellowstone?. <i>Global Ecology and Biogeography</i> , 2012, 21, 556-567.   | 2.7 | 69        |
| 79 | Efficacy of tree defense physiology varies with bark beetle population density: a basis for positive feedback in eruptive species. <i>Canadian Journal of Forest Research</i> , 2011, 41, 1174-1188.                              | 0.8 | 250       |
| 80 | The interdependence of mechanisms underlying climate-driven vegetation mortality. <i>Trends in Ecology and Evolution</i> , 2011, 26, 523-532.   | 4.2 | 839       |
| 81 | Altered GAI activity of hybrid aspen has minimal effects on the performance of a polyphagous weevil, <i>Polydrusus sericeus</i> . <i>Entomologia Experimentalis Et Applicata</i> , 2011, 138, 104-109.                            | 0.7 | 1         |
| 82 | Prior host feeding experience influences ovipositional but not feeding preference in a polyphagous insect herbivore. <i>Entomologia Experimentalis Et Applicata</i> , 2011, 138, 137-145.   | 0.7 | 33        |
| 83 | Cellulose-degrading bacteria associated with the invasive woodwasp <i>Sirex noctilio</i> . <i>ISME Journal</i> , 2011, 5, 1323-1331.  | 4.4 | 154       |
| 84 | Potential insight for drug discovery from high-fidelity receptor-mediated transduction mechanisms in insects. <i>Expert Opinion on Drug Discovery</i> , 2011, 6, 1091-1101.   | 2.5 | 0         |
| 85 | Fire Injury Reduces Inducible Defenses of Lodgepole Pine against Mountain Pine Beetle. <i>Journal of Chemical Ecology</i> , 2011, 37, 1184-1192.  | 0.9 | 33        |
| 86 | Responses of Bark Beetle-Associated Bacteria to Host Monoterpenes and Their Relationship to Insect Life Histories. <i>Journal of Chemical Ecology</i> , 2011, 37, 808-817.  | 0.9 | 73        |
| 87 | Presence and Diversity of <i>Streptomyces</i> in <i>Dendroctonus</i> and Sympatric Bark Beetle Galleries Across North America. <i>Microbial Ecology</i> , 2011, 61, 759-768.  | 1.4 | 63        |
| 88 | From Commensal to Pathogen: Translocation of <i>Enterococcus faecalis</i> from the Midgut to the Hemocoel of <i>Manduca sexta</i> . <i>MBio</i> , 2011, 2, e00065-11.   | 1.8 | 133       |
| 89 | Temporal and Species Variation in Cold Hardiness Among Invasive Rhizophagous Weevils (Coleoptera: Tj ETQq1 1 0.784314 rgBT /Over<br>104, 59-67.   | 1.3 | 7         |
| 90 | Robustness of the Bacterial Community in the Cabbage White Butterfly Larval Midgut. <i>Microbial Ecology</i> , 2010, 59, 199-211.   | 1.4 | 142       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | Symbiosis research, technology, and education: Proceedings of the 6th International Symbiosis Society Congress held in Madison Wisconsin, USA, August 2009. <i>Symbiosis</i> , 2010, 51, 1-12.   | 1.2 | 1         |
| 92  | Chemical modulators of the innate immune response alter gypsy moth larval susceptibility to <i>Bacillus thuringiensis</i> . <i>BMC Microbiology</i> , 2010, 10, 129.   | 1.3 | 48        |
| 93  | Too close for comfort: effect of trap spacing distance and pattern on statistical inference of behavioral choice tests in the field. <i>Entomologia Experimentalis Et Applicata</i> , 2010, 136, 66-71.                                  | 0.7 | 13        |
| 94  | Performance of the invasive weevil <i>Polydrusus sericeus</i> is influenced by atmospheric CO <sub>2</sub> and host species. <i>Agricultural and Forest Entomology</i> , 2010, 12, 285-292.  | 0.7 | 11        |
| 95  | Variation in Complex Semiochemical Signals Arising From Insects and Host Plants. <i>Environmental Entomology</i> , 2010, 39, 874-882.  | 0.7 | 12        |
| 96  | Geographic Variation in Bacterial Communities Associated With the Red Turpentine Beetle (Coleoptera: Curculionidae). <i>Environmental Entomology</i> , 2010, 39, 406-414.  | 0.7 | 64        |
| 97  | Laboratory Performance of Two Polyphagous Invasive Weevils on the Predominant Woody Plant Species of a Northern Hardwood Community. <i>Environmental Entomology</i> , 2010, 39, 1242-1248.   | 0.7 | 8         |
| 98  | Host Plant Phenology Affects Performance of an Invasive Weevil, <i>Phyllobius oblongus</i> (Coleoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50   | 0.7 | 18        |
| 99  | Effect of Clonal Variation Among Hybrid Poplars on Susceptibility of Gypsy Moth (Lepidoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 1 2010, 103, 718-725.   | 0.8 | 4         |
| 100 | Predisposition to bark beetle attack by root herbivores and associated pathogens: Roles in forest decline, gap formation, and persistence of endemic bark beetle populations. <i>Forest Ecology and Management</i> , 2010, 259, 374-382. | 1.4 | 43        |
| 101 | Assemblage of Hymenoptera arriving at logs colonized by <i>Ips pini</i> (Coleoptera: Curculionidae: Tj ETQq1 1 0.784314 rgBT /Overlock 0.4 6   | 0.4 | 6         |
| 102 | Contributions of gut bacteria to <i>Bacillus thuringiensis</i> -induced mortality vary across a range of Lepidoptera. <i>BMC Biology</i> , 2009, 7, 11.  | 1.7 | 156       |
| 103 | Survey and phylogenetic analysis of culturable microbes in the oral secretions of three bark beetle species. <i>Entomologia Experimentalis Et Applicata</i> , 2009, 131, 138-147.  | 0.7 | 36        |
| 104 | Mate-finding failure as an important cause of Allee effects along the leading edge of an invading insect population. <i>Entomologia Experimentalis Et Applicata</i> , 2009, 133, 307-314.  | 0.7 | 69        |
| 105 | Resident Microbiota of the Gypsy Moth Midgut Harbors Antibiotic Resistance Determinants. <i>DNA and Cell Biology</i> , 2009, 28, 109-117.  | 0.9 | 79        |
| 106 | Movement of outbreak populations of mountain pine beetle: influences of spatiotemporal patterns and climate. <i>Ecography</i> , 2008, 31, 348-358.   | 2.1 | 166       |
| 107 | The enemy of my enemy is still my enemy: competitors add to predator load of a tree-killing bark beetle. <i>Agricultural and Forest Entomology</i> , 2008, 10, 411-421.  | 0.7 | 27        |
| 108 | Cross-scale Drivers of Natural Disturbances Prone to Anthropogenic Amplification: The Dynamics of Bark Beetle Eruptions. <i>BioScience</i> , 2008, 58, 501-517.  | 2.2 | 1,410     |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 109 | Spatial-Temporal Modeling of Forest Gaps Generated by Colonization From Below- and Above-Ground Bark Beetle Species. <i>Journal of the American Statistical Association</i> , 2008, 103, 162-177.  | 1.8 | 23        |
| 110 | <i>Bursaphelenchus rufipennis</i> n. sp. (Nematoda: Parasitaphelenchinae) and redescription of <i>Ektaphelenchus obtusus</i> (Nematoda: Ektaphelenchinae), associates from nematangia on the hind wings of <i>Dendroctonus rufipennis</i> (Coleoptera: Scolytidae). <i>Nematology</i> , 2008, 10, 925-955. | 0.2 | 30        |
| 111 | Gut Microbiota of an Invasive Subcortical Beetle, <i>Agrilus planipennis</i> Fairmaire, Across Various Life Stages. <i>Environmental Entomology</i> , 2008, 37, 1344-1353.   | 0.7 | 71        |
| 112 | Gut Microbiota of an Invasive Subcortical Beetle, <i>Agrilus planipennis</i> Fairmaire, Across Various Life Stages. <i>Environmental Entomology</i> , 2008, 37, 1344-1353.   | 0.7 | 64        |
| 113 | Multipartite Symbioses Among Fungi, Mites, Nematodes, and the Spruce Beetle, <i>Dendroctonus rufipennis</i> . <i>Environmental Entomology</i> , 2008, 37, 956-963.   | 0.7 | 39        |
| 114 | Preoutbreak Dynamics of a Recently Established Invasive Herbivore: Roles of Natural Enemies and Habitat Structure in Stage-Specific Performance of Gypsy Moth (Lepidoptera: Lymantriidae) Populations in Northeastern Wisconsin. <i>Environmental Entomology</i> , 2008, 37, 1174-1184.                    | 0.7 | 7         |
| 115 | Parasitoids and Dipteran Predators Exploit Volatiles from Microbial Symbionts to Locate Bark Beetles. <i>Environmental Entomology</i> , 2008, 37, 150-161.   | 0.7 | 34        |
| 116 | Signal Mimics Derived from a Metagenomic Analysis of the Gypsy Moth Gut Microbiota. <i>Applied and Environmental Microbiology</i> , 2007, 73, 3669-3676.   | 1.4 | 66        |
| 117 | Interactions among intraspecific competition, emergence patterns, and host selection behaviour in <i>Ips pini</i> (Coleoptera: Scolytinae). <i>Ecological Entomology</i> , 2007, 32, 162-171.  | 1.1 | 28        |
| 118 | Continuous Time Modelling of Dynamical Spatial Lattice Data Observed at Sparsely Distributed Times. <i>Journal of the Royal Statistical Society Series B: Statistical Methodology</i> , 2007, 69, 701-713.   | 1.1 | 5         |
| 119 | Phylogeography of spruce beetles ( <i>Dendroctonus rufipennis</i> Kirby) (Curculionidae: Scolytinae) in North America. <i>Molecular Ecology</i> , 2007, 16, 2560-2573.   | 2.0 | 56        |
| 120 | Can chemical communication be cryptic? Adaptations by herbivores to natural enemies exploiting prey semiochemistry. <i>Oecologia</i> , 2007, 153, 1009-1019.   | 0.9 | 35        |
| 121 | Landscape level analysis of mountain pine beetle in British Columbia, Canada: spatiotemporal development and spatial synchrony within the present outbreak. <i>Ecography</i> , 2006, 29, 427-441.  | 2.1 | 197       |
| 122 | Response of ground beetle (Carabidae) assemblages to logging history in northern hardwood-hemlock forests. <i>Forest Ecology and Management</i> , 2006, 222, 335-347.  | 1.4 | 61        |
| 123 | Is the outbreak status of <i>Thrips calcaratus</i> Uzel in North America due to altered host relationships?. <i>Forest Ecology and Management</i> , 2006, 225, 200-206.  | 1.4 | 2         |
| 124 | Sources of Insect and Plant Volatiles Attractive to Cottonwood Leaf Beetles Feeding on Hybrid Poplar. <i>Journal of Chemical Ecology</i> , 2006, 32, 2585-2594.  | 0.9 | 17        |
| 125 | Bacteria in oral secretions of an endophytic insect inhibit antagonistic fungi. <i>Ecological Entomology</i> , 2006, 31, 636-645.  | 1.1 | 184       |
| 126 | Characterization of Gut-Associated Bacteria in Larvae and Adults of the Southern Pine Beetle, <i>Dendroctonus frontalis</i> Zimmermann. <i>Environmental Entomology</i> , 2006, 35, 1710-1717.   | 0.7 | 74        |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 127 | Characterization of Gut-Associated Bacteria in Larvae and Adults of the Southern Pine Beetle, <i>Dendroctonus frontalis</i> Zimmermann. <i>Environmental Entomology</i> , 2006, 35, 1710-1717.  | 0.7 | 41        |
| 128 | Bacteria Associated with the Guts of Two Wood-Boring Beetles: <i>Anoplophora glabripennis</i> and <i>Saperda vestita</i> (Cerambycidae). <i>Environmental Entomology</i> , 2006, 35, 625-629.   | 0.7 | 121       |
| 129 | Midgut bacteria required for <i>Bacillus thuringiensis</i> insecticidal activity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 15196-15199.  | 3.3 | 379       |
| 130 | Selective manipulation of predators using pheromones: responses to frontalin and ipsdienol pheromone components of bark beetles in the Great Lakes region. <i>Agricultural and Forest Entomology</i> , 2005, 7, 193-200.  | 0.7 | 26        |
| 131 | Modeling flight activity and population dynamics of the pine engraver, <i>Ips pini</i> , in the Great Lakes region: effects of weather and predators over short time scales. <i>Population Ecology</i> , 2005, 47, 61-69.   | 0.7 | 30        |
| 132 | Interactions Among Conifer Terpenoids and Bark Beetles Across Multiple Levels of Scale: An Attempt to Understand Links Between Population Patterns and Physiological Processes. <i>Recent Advances in Phytochemistry</i> , 2005, 39, 79-118.  | 0.5 | 118       |
| 133 | Contrasts in Cellulolytic Activities of Gut Microorganisms Between the Wood Borer, <i>Saperda vestita</i> (Coleoptera: Cerambycidae), and the Bark Beetles, <i>Ips pini</i> and <i>Dendroctonus frontalis</i> (Coleoptera: Curculionidae). <i>Environmental Entomology</i> , 2005, 34, 541-547. | 0.7 | 111       |
| 134 | Quantifying sources of variation in the frequency of fungi associated with spruce beetles: Implications for hypothesis testing and sampling methodology in bark beetle-symbiont relationships. <i>Forest Ecology and Management</i> , 2005, 217, 187-202.                                       | 1.4 | 38        |
| 135 | Effects of Diterpene Acids on Components of a Conifer Bark Beetle-Fungal Interaction: Tolerance by <i>Ips pini</i> and Sensitivity by Its Associate <i>Ophiostoma ips</i> . <i>Environmental Entomology</i> , 2005, 34, 486-493.  | 0.7 | 71        |
| 136 | Components of Antagonism and Mutualism in <i>Ips pini</i> -Fungal Interactions: Relationship to a Life History of Colonizing Highly Stressed and Dead Trees. <i>Environmental Entomology</i> , 2004, 33, 28-34.   | 0.7 | 58        |
| 137 | Species Assemblage Arriving at and Emerging from Trees Colonized by <i>Ips pini</i> in the Great Lakes Region: Partitioning by Time Since Colonization, Season, and Host Species. <i>Annals of the Entomological Society of America</i> , 2004, 97, 117-129.                                    | 1.3 | 21        |
| 138 | Does aggregation benefit bark beetles by diluting predation? Links between a group-colonisation strategy and the absence of emergent multiple predator effects. <i>Ecological Entomology</i> , 2004, 29, 129-138.   | 1.1 | 72        |
| 139 | Census of the Bacterial Community of the Gypsy Moth Larval Midgut by Using Culturing and Culture-Independent Methods. <i>Applied and Environmental Microbiology</i> , 2004, 70, 293-300.  | 1.4 | 472       |
| 140 | Behavior of Adult and Larval <i>Platysoma cylindrica</i> (Coleoptera: Histeridae) and Larval <i>Medetera bistriata</i> (Diptera: Dolichopodidae) During Subcortical Predation of <i>Ips pini</i> (Coleoptera: Scolytidae). <i>Journal of Insect Behavior</i> , 2004, 17, 115-128.               | 0.4 | 17        |
| 141 | Gender- and sequence-dependent predation within group colonizers of defended plants: a constraint on cheating among bark beetles?. <i>Oecologia</i> , 2004, 138, 253-258.   | 0.9 | 20        |
| 142 | Density-dependent effects of multiple predators sharing a common prey in an endophytic habitat. <i>Oecologia</i> , 2004, 139, 418-426.  | 0.9 | 12        |
| 143 | FEEDBACK BETWEEN INDIVIDUAL HOST SELECTION BEHAVIOR AND POPULATION DYNAMICS IN AN ERUPTIVE HERBIVORE. <i>Ecological Monographs</i> , 2004, 74, 101-116.   | 2.4 | 125       |
| 144 | Phloeophagous and predaceous insects responding to synthetic pheromones of bark beetles inhabiting white spruce stands in the Great Lakes region. <i>Journal of Chemical Ecology</i> , 2003, 29, 1651-1663.   | 0.9 | 20        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 145 | Leaf ontogeny influences leaf phenolics and the efficacy of genetically expressed <i>Bacillus thuringiensis cry1A(a)</i> d-endotoxin in hybrid poplar against gypsy moth. <i>Journal of Chemical Ecology</i> , 2003, 29, 2585-2602.                                | 0.9 | 24        |
| 146 | Effect of varying monoterpene concentrations on the response of <i>Ips pini</i> (Coleoptera: Scolytidae) to its aggregation pheromone: implications for pest management and ecology of bark beetles. <i>Agricultural and Forest Entomology</i> , 2003, 5, 269-274. | 0.7 | 95        |
| 147 | Seasonal Activity of Adult, Ground-occurring Beetles (Coleoptera) in Forests of Northeastern Wisconsin and the Upper Peninsula of Michigan. <i>American Midland Naturalist</i> , 2003, 149, 121-133.   | 0.2 | 21        |
| 148 | Spatial analysis of forest gaps resulting from bark beetle colonization of red pines experiencing belowground herbivory and infection. <i>Forest Ecology and Management</i> , 2003, 177, 145-153.  | 1.4 | 12        |
| 149 | Fate of Conifer Terpenes in a Polyphagous Folivore: Evidence for Metabolism by Gypsy Moth (Lepidoptera: Lymantriidae). <i>Journal of Entomological Science</i> , 2003, 38, 583-601.  | 0.2 | 7         |
| 150 | Density-mediated responses of bark beetles to host allelochemicals: a link between individual behaviour and population dynamics. <i>Ecological Entomology</i> , 2002, 27, 484-492.   | 1.1 | 36        |
| 151 | Desiccation of <i>Pinus</i> foliage induced by conifer sawfly oviposition: effect on egg viability. <i>Ecological Entomology</i> , 2002, 27, 618-621.  | 1.1 | 11        |
| 152 | Population Dynamics of <i>Ips pini</i> and <i>Ips grandicollis</i> in Red Pine Plantations in Wisconsin: Within- and Between-Year Associations with Predators, Competitors, and Habitat Quality. <i>Environmental Entomology</i> , 2002, 31, 1043-1051.            | 0.7 | 46        |
| 153 | Heritability of Host Acceptance and Gallery Construction Behaviors of the Bark Beetle <i>Ips pini</i> (Coleoptera: Scolytidae). <i>Environmental Entomology</i> , 2002, 31, 1276-1281.   | 0.7 | 22        |
| 154 | Bark beetles and fungal associates colonizing white spruce in the Great Lakes region. <i>Canadian Journal of Forest Research</i> , 2002, 32, 1137-1150.  | 0.8 | 28        |
| 155 | Association of declining red pine stands with reduced populations of bark beetle predators, seasonal increases in root colonizing insects, and incidence of root pathogens. <i>Forest Ecology and Management</i> , 2002, 164, 221-236.                             | 1.4 | 60        |
| 156 | How many choices can your test animal compare effectively? Evaluating a critical assumption of behavioral preference tests. <i>Oecologia</i> , 2002, 133, 422-429.   | 0.9 | 62        |
| 157 | Relative effects of exophytic predation, endophytic predation, and intraspecific competition on a subcortical herbivore: consequences to the reproduction of <i>Ips pini</i> and <i>Thanasimus dubius</i> . <i>Oecologia</i> , 2002, 133, 483-491.                 | 0.9 | 36        |
| 158 | Mixed messages across multiple trophic levels: the ecology of bark beetle chemical communication systems. <i>Chemoecology</i> , 2001, 11, 49-65.   | 0.6 | 171       |
| 159 | Modulation of predator attraction to pheromones of two prey species by stereochemistry of plant volatiles. <i>Oecologia</i> , 2001, 127, 444-453.  | 0.9 | 75        |
| 160 | Kairomonal range of generalist predators in specialized habitats: responses to multiple phloeophagous species emitting pheromones vs. host odors. <i>Entomologia Experimentalis Et Applicata</i> , 2001, 99, 205-210.  | 0.7 | 37        |
| 161 | EFFECTS OF FOLIVORY ON SUBCORTICAL PLANT DEFENSES: CAN DEFENSE THEORIES PREDICT INTERGUILD PROCESSES?. <i>Ecology</i> , 2001, 82, 1387-1400.   | 1.5 | 82        |
| 162 | Effect of Host Tree Seasonal Phenology on Substrate Suitability for the Pine Engraver (Coleoptera: <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i> ) <i>Entomology</i> , 2001, 94, 844-849.  | 0.8 | 20        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 163 | Title is missing!. Journal of Chemical Ecology, 2000, 26, 823-840.  | 0.9 | 39        |
| 164 | Title is missing!. Journal of Chemical Ecology, 2000, 26, 1923-1939.  | 0.9 | 24        |
| 165 | Title is missing!. Journal of Chemical Ecology, 2000, 26, 2527-2548.  | 0.9 | 75        |
| 166 | BIOSYNTHESIS OF CONIFEROPHAGOUS BARK BEETLE PHEROMONES AND CONIFER ISOPRENOIDS: EVOLUTIONARY PERSPECTIVE AND SYNTHESIS. Canadian Entomologist, 2000, 132, 697-753.  | 0.4 | 120       |
| 167 | Improved Population Monitoring of Bark Beetles and Predators by Incorporating Disparate Behavioral Responses to Semiochemicals. Environmental Entomology, 2000, 29, 618-629.  | 0.7 | 64        |
| 168 | Synergy Between Zwittermicin A and <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> Against Gypsy Moth (Lepidoptera: Lymantriidae). Environmental Entomology, 2000, 29, 101-107.  | 0.7 | 80        |
| 169 | Influences of Host Chemicals and Internal Physiology on the Multiple Steps of Postlanding Host Acceptance Behavior of <i>Ips pini</i> (Coleoptera: Scolytidae). Environmental Entomology, 2000, 29, 442-453.                          | 0.7 | 86        |
| 170 | Exploiting Behavioral Disparities Among Predators and Prey to Selectively Remove Pests: Maximizing the Ratio of Bark Beetles to Predators Removed During Semiochemically Based Trap-Out. Environmental Entomology, 2000, 29, 651-660. | 0.7 | 59        |
| 171 | Compound effects of induced plant responses on insect herbivores and parasitoids: implications for tritrophic interactions. Ecological Entomology, 2000, 25, 171-179.   | 1.1 | 102       |
| 172 | Effects of forest management practices on the diversity of ground-occurring beetles in mixed northern hardwood forests of the Great Lakes Region. Forest Ecology and Management, 2000, 139, 135-155.                                  | 1.4 | 98        |
| 173 | Title is missing!. Journal of Chemical Ecology, 1999, 25, 861-880.  | 0.9 | 64        |
| 174 | Title is missing!. Journal of Chemical Ecology, 1999, 25, 1771-1797.  | 0.9 | 28        |
| 175 | Partitioning of <sup>14</sup> C-labeled photosynthate to allelochemicals and primary metabolites in source and sink leaves of aspen: evidence for secondary metabolite turnover. Oecologia, 1999, 119, 408-418.                       | 0.9 | 63        |
| 176 | Effects of elicitation treatment and genotypic variation on induced resistance in <i>Populus</i> : impacts on gypsy moth (Lepidoptera: Lymantriidae) development and feeding behavior. Oecologia, 1999, 120, 295-303.                 | 0.9 | 79        |
| 177 | Effects of Selected <i>Larix laricina</i> Terpenoids on <i>Lymantria dispar</i> (Lepidoptera: Lymantriidae) Development and Behavior. Environmental Entomology, 1999, 28, 148-154.  | 0.7 | 24        |
| 178 | Title is missing!. Journal of Chemical Ecology, 1998, 24, 501-523.  | 0.9 | 30        |
| 179 | Productivity, drought tolerance and pest status of hybrid <i>Populus</i> : tree improvement and silvicultural implications. Biomass and Bioenergy, 1998, 14, 1-20.  | 2.9 | 41        |
| 180 | Endogenous and exogenous factors affecting parasitism of gypsy moth egg masses by <i>Ooencyrtus kuvanae</i> . Entomologia Experimentalis Et Applicata, 1998, 88, 123-135.   | 0.7 | 22        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 181 | Association of within-tree jack pine budworm feeding patterns with canopy level and within-needle variation of water, nutrient, and monoterpene concentrations. Canadian Journal of Forest Research, 1998, 28, 228-233.   | 0.8 | 14        |
| 182 | Heritability Estimates of Development Time and Size Characters in the Gypsy Moth (Lepidoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 1998, 27, 415-418.  | 0.7 | 5         |
| 183 | Effect of nitrogen availability on the growth and phytochemistry of hybrid poplar and the efficacy of the <math>Bacillus thuringiensis</math> <math>cry1A</math> (<math>a</math> <math>d</math>)-endotoxin on gypsy moth. Canadian Journal of Forest Research, 1998, 28, 1055-1067. | 0.8 | 9         |
| 184 | Effects of Host Diet on the Orientation, Development, and Subsequent Generations of the Gypsy Moth (Lepidoptera: Lymantriidae) Egg Parasitoid Ooencyrtus kuvanae (Hymenoptera: Encyrtidae). Environmental Entomology, 1997, 26, 1276-1282.  | 0.7 | 10        |
| 185 | Effects of Selected Midwestern Larval Host Plants on Performance by Two Strains of the Gypsy Moth (Lepidoptera: Lymantriidae) Parasitoid Cotesia melanoscela (Hymenoptera: Braconidae). Environmental Entomology, 1997, 26, 1155-1166.  | 0.7 | 19        |
| 186 | Individual and social components of wood ant response to conifer sawfly defence (Hymenoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 1998, 27, 415-418.   | 0.8 | 18        |
| 187 | Combined chemical defenses against an insect-fungal complex. Journal of Chemical Ecology, 1996, 22, 1367-1388.  | 0.9 | 126       |
| 188 | Defoliation tolerance affects the spatial and temporal distributions of larch sawfly and natural enemy populations. Ecological Entomology, 1996, 21, 259-269.   | 1.1 | 13        |
| 189 | Effects of biotic and abiotic stress on induced accumulation of terpenes and phenolics in red pines inoculated with bark beetle-vectored fungus. Journal of Chemical Ecology, 1995, 21, 601-626.  | 0.9 | 122       |
| 190 | Contributions of female oviposition patterns and larval behavior to group defense in conifer sawflies (hymenoptera: diprionidae). Oecologia, 1995, 103, 24-33.  | 0.9 | 53        |
| 191 | Interaction of pre-attack and induced monoterpene concentrations in host conifer defense against bark beetle-fungal complexes. Oecologia, 1995, 102, 285-295.   | 0.9 | 243       |
| 192 | Defoliation intensity and larval age interact to affect sawfly performance on previously injured Pinus resinosa. Oecologia, 1995, 102, 24-30.   | 0.9 | 15        |
| 193 | Field Evaluation of Transgenic Poplar Expressing a Bacillus thuringiensis cry1A (&sup2;) d-Endotoxin Gene Against Forest Tent Caterpillar (Lepidoptera: Lasiocampidae) and Gypsy Moth (Lepidoptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 1998, 28, 228-233.                  | 0.7 | 52        |
| 194 | Responses of Gypsy Moth (Lepidoptera: Lymantriidae) and Forest Tent Caterpillar (Lepidoptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2 1994, 23, 1030-1041.   | 0.7 | 52        |
| 195 | Maturation of the Male Pales Weevil (Coleoptera: Curculionidae) Reproductive System and its Effect on Male Response to Females. Annals of the Entomological Society of America, 1992, 85, 571-577.  | 1.3 | 16        |
| 196 | Comparison of insect, fungal, and mechanically induced defoliation of larch: effects on plant productivity and subsequent host susceptibility. Oecologia, 1992, 90, 411-416.  | 0.9 | 39        |
| 197 | The effect of host variability on growth and performance of the introduced pine sawfly, Diprionsimilis. Canadian Journal of Forest Research, 1991, 21, 1668-1674.   | 0.8 | 8         |
| 198 | Temporal and Spatial Disparities Among Bark Beetles, Predators, and Associates Responding to Synthetic Bark Beetle Pheromones: Ips pini (Coleoptera: Scolytidae) in Wisconsin. Environmental Entomology, 1991, 20, 1665-1679.   | 0.7 | 59        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 199 | Dispersal Patterns and Mark-and-Recapture Estimates of Two Pine Root Weevil Species, <i>Hylobius pales</i> and <i>Pachylobius picivorus</i> (Coleoptera: Curculionidae), in Christmas Tree Plantations. <i>Environmental Entomology</i> , 1990, 19, 1829-1836. | 0.7 | 22        |
| 200 | Chiral escape of bark beetles from predators responding to a bark beetle pheromone. <i>Oecologia</i> , 1989, 80, 566-569.  | 0.9 | 76        |
| 201 | Genetic Engineering of Trees to Enhance Resistance to Insects. <i>BioScience</i> , 1989, 39, 524-534.  | 2.2 | 85        |
| 202 | Computation of response factors for quantitative analysis of monoterpenes by gas-liquid chromatography. <i>Journal of Chemical Ecology</i> , 1988, 14, 1385-1390.  | 0.9 | 35        |
| 203 | Response of red and jack pines to inoculation with microbial associates of the pine engraver, <i>Ips pini</i> (Coleoptera: Scolytidae). <i>Canadian Journal of Forest Research</i> , 1988, 18, 581-586.  | 0.8 | 19        |
| 204 | Host resistance to invasion by lower stem and root infesting insects of pine: response to controlled inoculations with the fungal associate <i>Leptographium terebrantis</i> . <i>Canadian Journal of Forest Research</i> , 1988, 18, 675-681.                 | 0.8 | 21        |
| 205 | Seasonal and long-term responses of host trees to microbial associates of the pine engraver, <i>Ips pini</i> . <i>Canadian Journal of Forest Research</i> , 1988, 18, 1624-1634.   | 0.8 | 15        |
| 206 | Effect of Host Plant on Cannibalism Rates by Fall Armyworm (Lepidoptera: Noctuidae) Larvae. <i>Environmental Entomology</i> , 1987, 16, 672-675.   | 0.7 | 31        |
| 207 | Influence of Host Plant on Deterrence by Azadirachtin of Feeding by Fall Armyworm Larvae (Lepidoptera: Noctuidae). <i>Journal of Economic Entomology</i> , 1987, 80, 384-387.  | 0.8 | 35        |
| 208 | Maintenance of innate feeding preferences by a polyphagous insect despite ingestion of applied deleterious chemicals. <i>Entomologia Experimentalis Et Applicata</i> , 1987, 44, 221-227.  | 0.7 | 13        |
| 209 | Interacting Selective Pressures in Conifer-Bark Beetle Systems: A Basis for Reciprocal Adaptations?. <i>American Naturalist</i> , 1987, 129, 234-262.  | 1.0 | 182       |
| 210 | Evolution of Optimal Group Attack, with Particular Reference to Bark Beetles (Coleoptera: Scolytidae). <i>Evolution</i> , 1987, 41, 50-62.   | 1.5 | 91        |
| 211 | ACCUMULATION OF MONOTERPENES AND ASSOCIATED VOLATILES FOLLOWING INOCULATION OF GRAND FIR WITH A FUNGUS TRANSMITTED BY THE FIR ENGRAVER, <i>Scolytus ventralis</i> (Coleoptera: Scolytidae). <i>Environmental Entomology</i> , 1987, 16, 784-791.               | 1.0 | 119       |
| 212 | Physiological Differences Between Lodgepole Pines Resistant and Susceptible to the Mountain Pine Beetle 1 and Associated Microorganisms 2. <i>Environmental Entomology</i> , 1982, 11, 486-492.  | 0.7 | 183       |
| 213 | Potential Alternate Hosts of the Gypsy Moth 1 Parasite <i>Apanteles porthetriae</i> 234. <i>Environmental Entomology</i> , 1977, 6, 57-59.   | 0.7 | 8         |
| 214 | The impact is in the details: evaluating a standardized protocol and scale for determining non-native insect impact. <i>NeoBiota</i> , 0, 55, 61-83.   | 1.0 | 7         |