
Helga K Ising

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6877997/publications.pdf Version: 2024-02-01



HELCA K ISINC

#	Article	IF	CITATIONS
1	The association between migrant status and transition in an ultra-high risk for psychosis population. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 943-952.	1.6	5
2	Designing and Testing of a Health-Economic Markov Model for Prevention and Treatment of Early Psychosis. Expert Review of Pharmacoeconomics and Outcomes Research, 2020, 20, 269-279.	0.7	11
3	Cost-Effectiveness of Virtual Reality Cognitive Behavioral Therapy for Psychosis: Health-Economic Evaluation Within a Randomized Controlled Trial. Journal of Medical Internet Research, 2020, 22, e17098.	2.1	35
4	Ultrahigh risk for developing psychosis and psychotic personality organization. Microbial Biotechnology, 2019, 13, 673-676.	0.9	4
5	Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk: Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data Meta-Analysis. Frontiers in Psychiatry, 2019, 10, 345.	1.3	29
6	CBT in the prevention of psychosis and other severe mental disorders in patients with an at risk mental state: A review and proposed next steps. Schizophrenia Research, 2019, 203, 88-93.	1.1	31
7	Four-Year Cost-effectiveness of Cognitive Behavior Therapy for Preventing First-episode Psychosis: The Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. Schizophrenia Bulletin, 2017, 43, sbw084.	2.3	29
8	The Prodromal Questionnaire: a case for IRT-based adaptive testing of psychotic experiences?. International Journal of Methods in Psychiatric Research, 2017, 26, e1518.	1.1	10
9	The effect of childhood adversity on 4-year outcome in individuals at ultra high risk for psychosis in the Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. Psychiatry Research, 2017, 247, 55-62.	1.7	23
10	Exposing therapists to trauma-focused treatment in psychosis: effects on credibility, expected burden, and harm expectancies. Högre Utbildning, 2016, 7, 31712.	1.4	10
11	COMT Val 158 Met genotype and cannabis use in people with an At Risk Mental State for psychosis: Exploring Gene x Environment interactions. Schizophrenia Research, 2016, 174, 24-28.	1.1	22
12	Predicting trauma-focused treatment outcome in psychosis. Schizophrenia Research, 2016, 176, 239-244.	1.1	11
13	Four-Year Follow-up of Cognitive Behavioral Therapy in Persons at Ultra-High Risk for Developing Psychosis: The Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. Schizophrenia Bulletin, 2016, 42, 1243-1252.	2.3	57
14	The distribution of self-reported psychotic-like experiences in non-psychotic help-seeking mental health patients in the general population; a factor mixture analysis. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 349-358.	1.6	16
15	PREVENTIVE PSYCHOTHERAPY. Schizophrenia Research, 2014, 153, S42-S43.	1.1	1
16	Depression and social anxiety in help-seeking patients with an ultra-high risk for developing psychosis. Psychiatry Research, 2013, 209, 309-313.	1.7	55
17	Cognitive Behavioral Therapy for Subjects at Ultrahigh Risk for Developing Psychosis: A Randomized Controlled Clinical Trial. Schizophrenia Bulletin, 2012, 38, 1180-1188.	2.3	168
18	The Validity of the 16-Item Version of the Prodromal Questionnaire (PQ-16) to Screen for Ultra High Risk of Developing Psychosis in the General Help-Seeking Population. Schizophrenia Bulletin, 2012, 38, 1288-1296.	2.3	304

#	Article	IF	CITATIONS
19	A single blind randomized controlled trial of cognitive behavioural therapy in a help-seeking population with an At Risk Mental State for psychosis: the Dutch Early Detection and Intervention Evaluation (EDIE-NL) trial. Trials, 2010, 11, 30.	0.7	46