Clodagh M Toomey

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6880270/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Establishing outcome measures in early knee osteoarthritis. Nature Reviews Rheumatology, 2019, 15, 438-448.	3.5	88
2	Protein Supplementation at Breakfast and Lunch for 24 Weeks beyond Habitual Intakes Increases Whole-Body Lean Tissue Mass in Healthy Older Adults. Journal of Nutrition, 2016, 146, 65-69.	1.3	74
3	Exercise Therapy in Juvenile Idiopathic Arthritis: A Systematic Review and Meta-Analysis. Archives of Physical Medicine and Rehabilitation, 2018, 99, 178-193.e1.	0.5	71
4	Ultrasound Measurement of Subcutaneous Adipose Tissue Thickness Accurately Predicts Total and Segmental Body Fat of Young Adults. Ultrasound in Medicine and Biology, 2012, 38, 28-34.	0.7	70
5	A Review of Body Composition Measurement in the Assessment of Health. Topics in Clinical Nutrition, 2015, 30, 16-32.	0.2	52
6	Higher Fat Mass Is Associated With a History of Knee Injury in Youth Sport. Journal of Orthopaedic and Sports Physical Therapy, 2017, 47, 80-87.	1.7	49
7	Association between MRI-defined osteoarthritis, pain, function and strength 3–10 years following knee joint injury in youth sport. British Journal of Sports Medicine, 2018, 52, 934-939.	3.1	48
8	Measurement of maximal isometric torque and muscle quality of the knee extensors and flexors in healthy 50―to 70â€yearâ€old women. Clinical Physiology and Functional Imaging, 2017, 37, 448-455.	0.5	44
9	The effect of hydration status on the measurement of lean tissue mass by dual-energy X-ray absorptiometry. European Journal of Applied Physiology, 2017, 117, 567-574.	1.2	40
10	Technical considerations for accurate measurement of subcutaneous adipose tissue thickness using B-mode ultrasound. Ultrasound, 2011, 19, 91-96.	0.3	38
11	Health-related Outcomes after a Youth Sport–related Knee Injury. Medicine and Science in Sports and Exercise, 2019, 51, 255-263.	0.2	38
12	Generalised equations for the prediction of percentage body fat by anthropometry in adult men and women aged 18–81 years. British Journal of Nutrition, 2013, 109, 678-685.	1.2	28
13	Implementing a junior high school-based programme to reduce sports injuries through neuromuscular training (iSPRINT): a cluster randomised controlled trial (RCT). British Journal of Sports Medicine, 2020, 54, 913-919.	3.1	27
14	Twelve weeks' progressive resistance training combined with protein supplementation beyond habitual intakes increases upper leg lean tissue mass, muscle strength and extended gait speed in healthy older women. Biogerontology, 2017, 18, 881-891.	2.0	26
15	The Association Between Moderate and Vigorous Physical Activity and Time to Medical Clearance to Return to Play Following Sport-Related Concussion in Youth Ice Hockey Players. Frontiers in Neurology, 2019, 10, 588.	1.1	20
16	Muscle strength can better differentiate between gradations of functional performance than muscle quality in healthy 50–70 y women. Brazilian Journal of Physical Therapy, 2017, 21, 457-464.	1.1	17
17	Gait Adaptations in Youth With Juvenile Idiopathic Arthritis. Arthritis Care and Research, 2020, 72, 917-924.	1.5	14
18	Body composition analysis of inter-county Gaelic athletic association players measured by dual energy X-ray absorptiometry. Journal of Sports Sciences, 2016, 34, 1015-1020.	1.0	8

Clodagh M Toomey

#	Article	IF	CITATIONS
19	Secondary consequences of juvenile idiopathic arthritis in children and adolescents with knee involvement: physical activity, adiposity, fitness, and functional performance. Rheumatology International, 2022, 42, 319-327.	1.5	8
20	Health-Related Outcomes 3-15 Years Following Ankle Sprain Injury in Youth Sport: What Does the Future Hold?. Foot and Ankle International, 2022, 43, 21-31.	1.1	7
21	Changes in exertion-related symptoms in adults and youth who have sustained a sport-related concussion. Journal of Science and Medicine in Sport, 2021, 24, 2-6.	0.6	6
22	Seasonal changes in body composition of inter-county Gaelic Athletic Association hurlers. Journal of Sports Sciences, 2017, 35, 2427-2432.	1.0	5
23	Visual rating of movement quality in individuals with and without a history of intra-articular knee injury. Physiotherapy Theory and Practice, 2021, 37, 1474-1480.	0.6	5
24	Adiposity as a Risk Factor for Sport Injury in Youth: A Systematic Review. Clinical Journal of Sport Medicine, 2022, 32, 418-426.	0.9	5
25	Does a history of youth sport-related knee injury still impact accelerometer-measured levels of physical activity after 3–12 years?. Physical Therapy in Sport, 2022, 55, 90-97.	0.8	5
26	More Than Just Adolescence: Differences in Fatigue Between Youth With Cerebral Palsy and Typically Developing Peers. Annals of Rehabilitation Medicine, 2021, 45, 197-203.	0.6	4
27	What Does the Future Hold? Health-Related Quality of Life 3–12 Years Following a Youth Sport-Related Knee Injury. International Journal of Environmental Research and Public Health, 2021, 18, 6877.	1.2	3
28	Knee Injury and Osteoarthritis Outcome Score (KOOS) Responder Criteria and Minimal Detectable Change 3–12 Years Following a Youth Sport-Related Knee Injury. Journal of Clinical Medicine, 2021, 10, 522.	1.0	2
29	15â€The association between physical activity and 3–15 year history of sport-related intra-articular knee injury: a matched cohort design. , 2018, , .		1
30	A SYSTEMATIC REVIEW OF THE ASSOCIATION BETWEEN ADIPOSITY AND SPORT INJURY RISK IN YOUTH. British Journal of Sports Medicine, 2017, 51, 396.2-397.	3.1	0
31	The association between moderate and vigorous physical activity and time to medical clearance to return to play following sport-related concussion in youth ice-hockey players. British Journal of Sports Medicine, 2017, 51, A44.1-A44.	3.1	0
32	6â€The consequences of knee joint injury in youth sport. , 2018, , .		0
33	051â€Implementing a school prevention program to reduce injuries through neuromuscular training (iSPRINT): a cluster-randomized controlled trial 2021		0