

Levente Laczkó³

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6883415/publications.pdf>

Version: 2024-02-01

18

papers

134

citations

1307594

7

h-index

1281871

11

g-index

18

all docs

18

docs citations

18

times ranked

221

citing authors

#	ARTICLE	IF	CITATIONS
1	The effect of gut passage by waterbirds on the seed coat and pericarp of diaspores lacking “external flesh”. Evidence for widespread adaptation to endozoochory in angiosperms. <i>PLoS ONE</i> , 2019, 14, e0226551.	2.5	26
2	Iterative allogamy–autogamy transitions drive actual and incipient speciation during the ongoing evolutionary radiation within the orchid genus <i>Epipactis</i> (Orchidaceae). <i>Annals of Botany</i> , 2019, 124, 481-497.	2.9	24
3	Evolutionary history of the Pasque-flowers (<i>Pulsatilla</i> , Ranunculaceae): Molecular phylogenetics, systematics and rDNA evolution. <i>Molecular Phylogenetics and Evolution</i> , 2019, 135, 45-61.	2.7	18
4	Phylogenetic and Morphological Analysis of Birch Mice (Genus <i>Sicista</i> , Family Sminthidae, Rodentia) in the Kazak Cradle with Description of a New Species. <i>Journal of Mammalian Evolution</i> , 2019, 26, 147-163.	1.8	12
5	Out of Colchis: The Colonization of Europe by <i>Primula vulgaris</i> Huds. (Primulaceae). <i>Acta Societatis Botanicorum Poloniae</i> , 2020, 89, .	0.8	10
6	Molecular phylogenetics, seed morphometrics, chromosome number evolution and systematics of European <i>Elatine</i> L. (Elatinaceae) species. <i>PeerJ</i> , 2016, 4, e2800.	2.0	10
7	Multi-Locus Genetic Identification of a Newly Discovered Population Reveals a Deep Genetic Divergence in European Blind Mole Rats (Rodentia: Spalacidae: <i>Nannospalax</i>). <i>Annales Zoologici Fennici</i> , 2020, 57, 89.	0.6	8
8	The first archaeobotanical evidence of <i>Lagenaria siceraria</i> from the territory of Hungary: histology, phytoliths and (a)DNA. <i>Vegetation History and Archaeobotany</i> , 2017, 26, 125-142.	2.1	7
9	Comparative analysis of human papillomavirus type 6 complete genomes originated from head and neck and anogenital disorders. <i>Infection, Genetics and Evolution</i> , 2019, 71, 140-150.	2.3	5
10	Is <i>Nymphaea lotus</i> var. <i>thermalis</i> a Tertiary relict in Europe?. <i>Aquatic Botany</i> , 2019, 155, 1-4.	1.6	4
11	<i>Hepatica transsilvanica</i> Fuss (Ranunculaceae) is an Allotetraploid Relict of the Tertiary Flora in Europe – Molecular Phylogenetic Evidence. <i>Acta Societatis Botanicorum Poloniae</i> , 2020, 89, .	0.8	4
12	The Phylogenetic Position of <i>Vincetoxicum pannonicum</i> (Borhidi) Holub Supports the Species' Allopolyploid Hybrid Origin. <i>Acta Societatis Botanicorum Poloniae</i> , 2020, 89, .	0.8	2
13	The <code><scp>RadOrgMiner</scp></code> pipeline: Automated genotyping of organellar loci from <code><scp>RADseq</scp></code> data. <i>Methods in Ecology and Evolution</i> , 2022, 13, 1962-1975.	5.2	2
14	Seasonal variation of genotypes and reproductive plasticity in a facultative clonal freshwater invertebrate animal (<i>Hydra oligactis</i>) living in a temperate lake. <i>Ecology and Evolution</i> , 2022, 12, .	1.9	2
15	Title is missing!. , 2019, 14, e0226551.	0	0
16	Title is missing!. , 2019, 14, e0226551.	0	0
17	Title is missing!. , 2019, 14, e0226551.	0	0
18	Title is missing!. , 2019, 14, e0226551.	0	0