Birgitt Wiese

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6892951/publications.pdf

Version: 2024-02-01

		159358	189595
136	3,391	30	50
papers	citations	h-index	g-index
139	139	139	5048
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	AD dementia risk in late MCI, in early MCI, and in subjective memory impairment. Alzheimer's and Dementia, 2014, 10, 76-83.	0.4	414
2	Prevalence and risk factors for depression in non-demented primary care attenders aged 75Âyears and older. Journal of Affective Disorders, 2008, 111, 153-163.	2.0	156
3	The essence of frailty: A systematic review and qualitative synthesis on frailty concepts and definitions. European Journal of Internal Medicine, 2018, 56, 3-10.	1.0	133
4	Longitudinal Predictors of Institutionalization in Old Age. PLoS ONE, 2015, 10, e0144203.	1.1	128
5	Cognitive performance before and after the onset of subjective cognitiveÂdecline in old age. Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring, 2015, 1, 194-205.	1.2	110
6	Impact of Depression on Health Care Utilization and Costs among Multimorbid Patients – Results from the MultiCare Cohort Study. PLoS ONE, 2014, 9, e91973.	1.1	88
7	A modified Delphi study to determine the level of consensus across the European Union on the structures, processes and desired outcomes of the management of polypharmacy in older people. PLoS ONE, 2017, 12, e0188348.	1.1	71
8	Differential Risk of Incident Alzheimer's Disease Dementia in Stable Versus Unstable Patterns of Subjective Cognitive Decline. Journal of Alzheimer's Disease, 2016, 54, 1135-1146.	1.2	70
9	Association between multimorbidity patterns and chronic pain in elderly primary care patients: a cross-sectional observational study. BMC Family Practice, 2016, 17, 68.	2.9	65
10	Gender differences in the effect of social support on health-related quality of life: results of a population-based prospective cohort study in old age in Germany. Quality of Life Research, 2016, 25, 1159-1168.	1.5	61
11	Determinants of health-related quality of life in older primary care patients: results of the longitudinal observational AgeCoDe Study. British Journal of General Practice, 2015, 65, e716-e723.	0.7	59
12	Prospective Associations between Single Foods, Alzheimer's Dementia and Memory Decline in the Elderly. Nutrients, 2018, 10, 852.	1.7	57
13	Comparison of different quantitative coronary analysis systems: ARTREK, CAAS, and CMS., 1996, 37, 14-22.		56
14	The CERAD Neuropsychological Assessment Battery Total Score Detects and Predicts Alzheimer Disease Dementia with High Diagnostic Accuracy. American Journal of Geriatric Psychiatry, 2014, 22, 1017-1028.	0.6	56
15	Mild cognitive impairment and quality of life in the oldest old: a closer look. Quality of Life Research, 2020, 29, 1675-1683.	1.5	53
16	Elevated HbA1c is Associated with Increased Risk of Incident Dementia in Primary Care Patients. Journal of Alzheimer's Disease, 2015, 44, 1203-1212.	1.2	52
17	The assessment of met and unmet care needs in the oldest old with and without depression using the Camberwell Assessment of Need for the Elderly (CANE): Results of the AgeMooDe study. Journal of Affective Disorders, 2016, 193, 309-317.	2.0	49
18	The impact of social engagement on health-related quality of life and depressive symptoms in old age - evidence from a multicenter prospective cohort study in Germany. Health and Quality of Life Outcomes, 2017, 15, 140.	1.0	48

#	Article	IF	Citations
19	Longitudinal predictors of informal and formal caregiving time in community-dwelling dementia patients. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 607-616.	1.6	46
20	Collaborative nurse-led self-management support for primary care patients with anxiety, depressive or somatic symptoms: Cluster-randomised controlled trial (findings of the SMADS study). International Journal of Nursing Studies, 2016, 63, 101-111.	2.5	42
21	A case study of polypharmacy management in nine European countries: Implications for change management and implementation. PLoS ONE, 2018, 13, e0195232.	1.1	42
22	The role of spousal loss in the development of depressive symptoms in the elderly â€" Implications for diagnostic systems. Journal of Affective Disorders, 2014, 161, 97-103.	2.0	41
23	Prevalence and determinants of overweight and obesity in old age in Germany. BMC Geriatrics, 2015, 15, 83.	1.1	41
24	Is function in instrumental activities of daily living a useful feature in predicting <scp>A</scp> Izheimer's disease dementia in subjective cognitive decline?. International Journal of Geriatric Psychiatry, 2019, 34, 193-203.	1.3	41
25	Excess health care costs of late-life depression – Results of the AgeMooDe study. Journal of Affective Disorders, 2016, 199, 139-147.	2.0	40
26	Systematic development and adjustment of the German version of the Supportive and Palliative Care Indicators Tool (SPICT-DE). BMC Palliative Care, 2018, 17, 27.	0.8	37
27	The Costs of Dementia From the Societal Perspective: Is Care Provided in the Community Really Cheaper than Nursing Home Care?. Journal of the American Medical Directors Association, 2014, 15, 117-126.	1.2	35
28	Subjective cognitive decline is longitudinally associated with lower health-related quality of life. International Psychogeriatrics, 2017, 29, 1939-1950.	0.6	35
29	Treatment preferences for depression in the elderly. International Psychogeriatrics, 2017, 29, 389-398.	0.6	34
30	AgeWell.de – study protocol of a pragmatic multi-center cluster-randomized controlled prevention trial against cognitive decline in older primary care patients. BMC Geriatrics, 2019, 19, 203.	1.1	34
31	Effect of Visual Impairment on Physical and Cognitive Function in Old Age: Findings of a Populationâ€Based Prospective Cohort Study in Germany. Journal of the American Geriatrics Society, 2016, 64, 2311-2316.	1.3	33
32	Impact of coronary heart disease on cognitive decline in Alzheimer's disease: a prospective longitudinal cohort study in primary care. British Journal of General Practice, 2017, 67, e111-e117.	0.7	33
33	Which types of mental work demands may be associated with reduced risk of dementia?. Alzheimer's and Dementia, 2017, 13, 431-440.	0.4	33
34	Disentangling the complex relation of disability and depressive symptoms in old age – findings of a multicenter prospective cohort study in Germany. International Psychogeriatrics, 2017, 29, 885-895.	0.6	30
35	What Is Successful Aging? A Psychometric Validation Study of Different Construct Definitions. Gerontologist, The, 2019, 59, 738-748.	2.3	30
36	Subjective Cognitive Decline May Be a Stronger Predictor of Incident Dementia in Women than in Men. Journal of Alzheimer's Disease, 2019, 68, 1469-1478.	1.2	28

#	Article	IF	Citations
37	A qualitative study on older primary care patients' perspectives on depression and its treatments - potential barriers to and opportunities for managing depression. BMC Family Practice, 2018, 19, 2.	2.9	26
38	Combined treatment with memantine and galantamineâ€CR comparedÂwith galantamineâ€CR only in antidementia drug naà ve patients with mildâ€toâ€moderate Alzheimer's disease. Alzheimer's and Dementia: Translational Research and Clinical Interventions, 2015, 1, 198-204.	1.8	25
39	Social inequalities in patient-reported outcomes among older multimorbid patients – results of the MultiCare cohort study. International Journal for Equity in Health, 2015, 14, 17.	1.5	25
40	Changes in Social Network Size Are Associated With Cognitive Changes in the Oldest-Old. Frontiers in Psychiatry, 2020, 11, 330.	1.3	25
41	Late-Life Depressive Symptoms andÂLifetime History of Major Depression: Cognitive Deficits are Largely Due toÂlncipient Dementia rather than Depression. Journal of Alzheimer's Disease, 2016, 54, 185-199.	1.2	23
42	How does social support affect functional impairment in late life? Findings of a multicenter prospective cohort study in Germany. Age and Ageing, 2017, 46, 813-820.	0.7	23
43	Are older men more vulnerable to depression than women after losing their spouse? Evidence from three German old-age cohorts (AgeDifferent.de platform). Journal of Affective Disorders, 2019, 256, 650-657.	2.0	23
44	The Impact of Depressive Symptoms on Healthcare Costs in Late Life: Longitudinal Findings From the AgeMooDe Study. American Journal of Geriatric Psychiatry, 2017, 25, 131-141.	0.6	22
45	Potentially inappropriate medication: Association between the use of antidepressant drugs and the subsequent risk for dementia. Journal of Affective Disorders, 2018, 226, 28-35.	2.0	22
46	Prevalence of Anxiety Symptoms and Their Association With Loss Experience in a Large Cohort Sample of the Oldest-Old. Results of the AgeCoDe/AgeQualiDe Study. Frontiers in Psychiatry, 2019, 10, 285.	1.3	22
47	Complex coevolution of depression and health-related quality of life in old age. Quality of Life Research, 2015, 24, 2713-2722.	1.5	20
48	Incident Subjective Cognitive Decline Does Not Predict Mortality in the Elderly – Results from the Longitudinal German Study on Ageing, Cognition, and Dementia (AgeCoDe). PLoS ONE, 2016, 11, e0147050.	1.1	20
49	Obesity and Associated Lifestyle in a Large Sample of Multi-Morbid German Primary Care Attendees. PLoS ONE, 2014, 9, e102587.	1.1	19
50	Associations between Dementia Outcomes and Depressive Symptoms, Leisure Activities, and Social Support. Dementia and Geriatric Cognitive Disorders Extra, 2014, 4, 481-493.	0.6	19
51	Advance directives and power of attorney for health care in the oldest-old – results of the AgeQualiDe study. BMC Geriatrics, 2017, 17, 85.	1.1	19
52	Optimal care at the end of life (OPAL): study protocol of a prospective interventional mixed-methods study with pretest-posttest-design in a primary health care setting considering the view of general practitioners, relatives of deceased patients and health care stakeholders. BMC Health Services Research, 2019, 19, 486.	0.9	18
53	Multidisciplinary intervention to improve medication safety in nursing home residents: protocol of a cluster randomised controlled trial (HIOPP-3-iTBX study). BMC Geriatrics, 2019, 19, 24.	1.1	18
54	Quality indicators for the evaluation of end-of-life care in Germany – a retrospective cross-sectional analysis of statutory health insurance data. BMC Palliative Care, 2020, 19, 187.	0.8	18

#	Article	IF	Citations
55	Comparison of FORTA, PRISCUS and EU(7)-PIM lists on identifying potentially inappropriate medication and its impact on cognitive function in multimorbid elderly German people in primary care: a multicentre observational study. BMJ Open, 2021, 11, e050344.	0.8	18
56	Women Outperform Men in Verbal Episodic Memory Even in Oldest-Old Age: 13-Year Longitudinal Results of the AgeCoDe/AgeQualiDe Study. Journal of Alzheimer's Disease, 2019, 69, 857-869.	1.2	17
57	Recurrent urinary tract infections and complications after symptomatic versus antibiotic treatment: follow-up of a randomised controlled trial. GMS German Medical Science, 2016, 14, Doc01.	2.7	17
58	Correlation between Oxford Elbow Score and Single Assessment Numeric Evaluation: Is one simple question enough?. Journal of Shoulder and Elbow Surgery, 2020, 29, 1223-1229.	1.2	16
59	Comparative effects of (S)-ketamine and racemic (R/S)-ketamine on psychopathology, state of consciousness and neurocognitive performance in healthy volunteers. European Neuropsychopharmacology, 2021, 44, 92-104.	0.3	16
60	The Role of Social Isolation and the Development of Depression. A Comparison of the Widowed and Married Oldest Old in Germany. International Journal of Environmental Research and Public Health, 2021, 18, 6986.	1.2	16
61	Recruitment and Baseline Characteristics of Participants in the AgeWell.de Study—A Pragmatic Cluster-Randomized Controlled Lifestyle Trial against Cognitive Decline. International Journal of Environmental Research and Public Health, 2021, 18, 408.	1.2	15
62	Social Support and Functional Decline in the Oldest Old. Gerontology, 2022, 68, 200-208.	1.4	15
63	Prevalence of pain and its associated factors among the oldest-olds in different care settings – results of the AgeQualiDe study. BMC Family Practice, 2018, 19, 85.	2.9	14
64	A novel superior medication-based chronic disease score predicted all-cause mortality in independent geriatric cohorts. Journal of Clinical Epidemiology, 2019, 105, 112-124.	2.4	14
65	Correlates of hospitalization among the oldest old: results of the AgeCoDe–AgeQualiDe prospective cohort study. Aging Clinical and Experimental Research, 2020, 32, 1295-1301.	1.4	14
66	Prediction of dementia of Alzheimer type by different types of subjective cognitive decline. Alzheimer's and Dementia, 2020, 16, 1745-1749.	0.4	14
67	Impact of falls on depressive symptoms among the oldest old: Results from the AgeQualiDe study. International Journal of Geriatric Psychiatry, 2018, 33, 1383-1388.	1.3	13
68	Correlates of institutionalization among the oldest oldâ€"Evidence from the multicenter AgeCoDeâ€AgeQualiDe study. International Journal of Geriatric Psychiatry, 2021, 36, 1095-1102.	1.3	13
69	A comparison of GP and GDS diagnosis of depression in late life among multimorbid patients – Results of the MultiCare study. Journal of Affective Disorders, 2014, 168, 276-283.	2.0	12
70	Prevalence and factors associated with obesity among the oldest old. Archives of Gerontology and Geriatrics, 2020, 89, 104069.	1.4	12
71	Perceived stress and study-related behavior and experience patterns of medical students: a cross-sectional study. BMC Medical Education, 2022, 22, 122.	1.0	12
72	Optimization of Low-Cost Culture Media for the Production of Biomass and Bacteriocin by a Urogenital Lactobacillus salivarius Strain. Probiotics and Antimicrobial Proteins, 2010, 2, 2-11.	1.9	11

#	Article	IF	Citations
73	Self-efficacy in multimorbid elderly patients with osteoarthritis in primary care—influence on pain-related disability. Clinical Rheumatology, 2015, 34, 1761-1767.	1.0	11
74	Mortality in Incident Cognitive Impairment: Results of the Prospective AgeCoDe Study. Journal of the American Geriatrics Society, 2017, 65, 738-746.	1.3	11
75	Prevalence and determinants of driving habits in the oldest old: Results of the multicenter prospective AgeCoDe-AgeQualiDe study. Archives of Gerontology and Geriatrics, 2019, 82, 245-250.	1.4	11
76	Study on advance care planning in care dependent community-dwelling older persons in Germany (STADPLAN): protocol of a cluster-randomised controlled trial. BMC Geriatrics, 2020, 20, 142.	1.1	11
77	Unmet care needs of the oldest old with late-life depression: A comparison of patient, caring relative and general practitioner perceptions $\hat{a} \in \mathbb{C}$ Results of the AgeMooDe study. Journal of Affective Disorders, 2016, 205, 182-189.	2.0	10
78	Frequent attendance in primary care in the oldest old: evidence from the AgeCoDe-AgeQualiDe study. Aging Clinical and Experimental Research, 2020, 32, 2629-2638.	1.4	10
79	Is the Short Form of the Mini-Mental State Examination (MMSE) a better screening instrument for dementia in older primary care patients than the original MMSE? Results of the German study on ageing, cognition, and dementia in primary care patients (AgeCoDe) Psychological Assessment, 2015, 27, 895-904.	1.2	9
80	Health behaviour, social support, socio-economic status and the 5-year progression of multimorbidity: Results from the MultiCare Cohort Study. Journal of Comorbidity, 2019, 9, 2235042X1988356.	3.9	9
81	Determinants of incident dementia in different old age groups: results of the prospective AgeCoDe/AgeQualiDe study. International Psychogeriatrics, 2020, 32, 645-659.	0.6	9
82	EQ-5D-3L health status and health state utilities of the oldest-old (85 +) in Germany: results from the AgeCoDe-AgeQualiDe study. Quality of Life Research, 2020, 29, 3223-3232.	1.5	9
83	Associations and correlates of general versus specific successful ageing components. European Journal of Ageing, 2021, 18, 549-563.	1.2	9
84	Which Factors Contribute to Frailty among the Oldest Old? Results of the Multicentre Prospective AgeCoDe and AgeQualiDe Study. Gerontology, 2020, 66, 460-466.	1.4	9
85	Systematic identification of critically ill and dying patients in primary care using the German version of the Supportive and Palliative Care Indicators Tool (SPICT-DE). GMS German Medical Science, 2020, 18, Doc02.	2.7	9
86	Narrative medicine-based intervention in primary care to reduce polypharmacy: results from the cluster-randomised controlled trial MultiCare AGENDA. BMJ Open, 2018, 8, e017653.	0.8	9
87	Satisfaction with general practice care in German patients with multimorbidity: a cross-sectional study. Family Practice, 2017, 34, 452-458.	0.8	8
88	Needs and health care costs in old age: a longitudinal perspective: results from the AgeMooDe study. Aging and Mental Health, 2020, 24, 1763-1768.	1.5	8
89	Family conferences and shared prioritisation to improve patient safety in the frail elderly (COFRAIL): study protocol of a cluster randomised intervention trial in primary care. Trials, 2020, 21, 285.	0.7	8
90	Do selfâ€reported hearing and visual impairments predict longitudinal dementia in older adults?. Journal of the American Geriatrics Society, 2021, 69, 1519-1528.	1.3	8

#	Article	IF	CITATIONS
91	Driving status and health-related quality of life among the oldest old: a population-based examination using data from the AgeCoDe–AgeQualiDe prospective cohort study. Aging Clinical and Experimental Research, 2021, 33, 3109-3115.	1.4	8
92	A geriatric assessment in general practice: prevalence, location, impact and doctor-patient perceptions of pain. BMC Family Practice, 2016, 17, 8.	2.9	7
93	Cross-cultural validation of the German version of the Four-Dimensional Symptom Questionnaire (4DSQ) in multimorbid elderly people. Quality of Life Research, 2018, 27, 2691-2697.	1.5	7
94	Determinants of health-care costs in the oldest-old in Germany. Journal of the Economics of Ageing, 2019, 14, 100200.	0.6	7
95	Sex-Specific Associations Between Depressive Symptoms and Risk for Subsequent Dementia. Journal of Alzheimer's Disease, 2020, 74, 151-161.	1.2	7
96	Help-seeking for psychological distress and its association with anxiety in the oldest old – results from the AgeQualiDe cohort study. Aging and Mental Health, 2021, 25, 923-929.	1.5	7
97	Anticholinergic drug burden according to the anticholinergic drug scale and the German anticholinergic burden and their impact on cognitive function in multimorbid elderly German people: a multicentre observational study. BMJ Open, 2021, 11, e044230.	0.8	7
98	Incidence of Anxiety in Latest Life and Risk Factors. Results of the AgeCoDe/AgeQualiDe Study. International Journal of Environmental Research and Public Health, 2021, 18, 12786.	1.2	7
99	International nifedipine trial on anti-atherosclerotic therapy (INTACT) — methodologic implications and results of a coronary angiographic follow-up study using computer-assisted film analysis. International Journal of Cardiovascular Imaging, 1990, 6, 117-133.	0.2	6
100	Older patients' perceived burdens of their health problems: a cross-sectional analysis in 74 German general practices. Patient Preference and Adherence, 2015, 9, 811.	0.8	6
101	The impact of depressive symptoms on utilization of home care by the elderly: Longitudinal results from the AgeMooDe study. Journal of Affective Disorders, 2016, 204, 247-254.	2.0	6
102	Unmet needs in the depressed primary care elderly and their relation to severity of depression: results from the AgeMooDe study. Aging and Mental Health, 2018, 22, 1038-1045.	1.5	6
103	Late-Life Depressive Symptoms Are Associated With Functional Impairment Cross-sectionally and Over Time: Results of the AgeMooDe Study. Journals of Gerontology - Series B Psychological Sciences and Social Sciences, 2020, 75, 811-820.	2.4	6
104	Post-discharge adjustment of medication in geriatric patients. Zeitschrift Fur Gerontologie Und Geriatrie, 2020, 53, 663-670.	0.8	6
105	Comparison of different quantitative coronary analysis systems: ARTREK, CAAS, and CMS., 1996, 37, 14.		6
106	Survival of Beneficial Vaginal Lactobacilli (BVL) to Different Gastrointestinal Tract Conditions. Current Pharmaceutical Design, 2020, 26, 3608-3618.	0.9	6
107	Factors Leading to Institutionalization among the Oldest Old: Longitudinal Findings from the AgeCoDe-AgeQualiDe Study. Gerontology, 2022, 68, 894-902.	1.4	6
108	Physical Activity Determinants in Older German Adults at Increased Dementia Risk with Multimorbidity: Baseline Results of the AgeWell.de Study. International Journal of Environmental Research and Public Health, 2022, 19, 3164.	1.2	6

#	Article	IF	CITATIONS
109	Social support and health-related quality of life among the oldest old — longitudinal evidence from the multicenter prospective AgeCoDe-AgeQualiDe study. Quality of Life Research, 2021, , 1.	1.5	6
110	Categorical and dimensional perspectives on depression in elderly primary care patients – Results of the AgeMooDe study. International Journal of Methods in Psychiatric Research, 2018, 27, .	1.1	5
111	Does transpersonal trust moderate the association between chronic conditions and general practitioner visits in the oldest old? Results of the AgeCoDe and AgeQualiDe study. Geriatrics and Gerontology International, 2019, 19, 705-710.	0.7	5
112	The value of the GP's clinical judgement in predicting dementia: a multicentre prospective cohort study among patients in general practice. British Journal of General Practice, 2019, 69, e786-e793.	0.7	5
113	Screening of autochthonous vaginal beneficial lactobacilli strains by their growth at high temperatures for technological applications. Antonie Van Leeuwenhoek, 2020, 113, 1393-1409.	0.7	5
114	Improving the continuity and coordination of ambulatory care through feedback and facilitated dialogue—a study protocol for a cluster-randomised trial to evaluate the ACD study (Accountable) Tj ETQq0 0 0) r gB7 /Ov	erl o ck 10 Tf 5
115	Gender Specifics of Healthy Ageing in Older Age as Seen by Women and Men (70+): A Focus Group Study. International Journal of Environmental Research and Public Health, 2022, 19, 3137.	1.2	5
116	Quality of end-of-life care in general practice – a pre–post comparison of a two-tiered intervention. , 2022, 23, 90.		5
117	Association of anxiety symptoms with health care use and costs in people aged 85 and over. International Journal of Geriatric Psychiatry, 2019, 34, 765-776.	1.3	4
118	Depressive Symptoms and Frailty Among the Oldest Old: Evidence From a Multicenter Prospective Study. Journal of the American Medical Directors Association, 2021, 22, 577-582.e2.	1.2	4
119	Effects of <scp>APOE</scp> e4â€ellele and mental work demands on cognitive decline in old age: Results from the German Study on Ageing, Cognition, and Dementia in Primary Care Patients (<scp>AgeCoDe</scp>). International Journal of Geriatric Psychiatry, 2021, 36, 152-162.	1.3	4
120	Association of Alcohol and Tobacco Consumption with Depression Severity in the Oldest Old. Results from the Age Different Old Age Cohort Platform. International Journal of Environmental Research and Public Health, 2021, 18, 7959.	1.2	4
121	Frailty and Autonomy among the Oldest Old: Evidence from the Multicenter Prospective AgeCoDe-AgeQualiDe Study. Gerontology, 2021, 67, 591-598.	1.4	3
122	Factors Contributing to Persistent Frequent Attendance in Primary Care Among the Oldest Old: Longitudinal Evidence From the AgeCoDe-AgeQualiDe Study. Frontiers in Medicine, 2022, 9, 815419.	1.2	3
123	Mild Depressive Symptoms Predict Mortality from Heart Failure in Elderly Men but Not Women. Journal of the American Geriatrics Society, 2015, 63, 2408-2410.	1.3	2
124	Lifestyle Aspects As A Predictor Of Pain Among Oldest-Old Primary Care Patients – A Longitudinal Cohort Study. Clinical Interventions in Aging, 2019, Volume 14, 1881-1888.	1.3	2
125	Nonsensus in the treatment of proximal humerus fractures: uncontrolled, blinded, comparative behavioural analysis between Homo chirurgicus accidentus and Macaca sylvanus. BMJ, The, 2020, 371, m4429.	3.0	2
126	Health-related quality of life and ego integrity among the oldest old – Evidence from the multicenter AgeCoDe-AgeQualiDe study. Archives of Gerontology and Geriatrics, 2021, 95, 104408.	1.4	2

#	Article	IF	CITATIONS
127	Incidence and risk factors of depressive symptoms in the highest age groups and competing mortality risk. Evidence from the AgeCoDe-AqeQualiDe prospective cohort study. Journal of Affective Disorders, 2022, 308, 494-501.	2.0	2
128	Non-random relations in drug use expressed as patterns comprising prescription and over-the-counter drugs in multimorbid elderly patients in primary care: Data of the exploratory analysis of the multicentre, observational cohort study MultiCare. European Journal of General Practice, 2021, 27, 119-129.	0.9	1
129	Disorientation in Time and Place in Old Age: Longitudinal Evidence from Three Old Age Cohorts in Germany (AgeDifferent.de Platform). Journal of Alzheimer's Disease, 2021, 79, 1589-1599.	1.2	1
130	Do oldest old individuals perceive receipt of informal care as a restriction or support of their autonomy?. Aging and Mental Health, 2022, 26, 1862-1873.	1.5	1
131	GPs' perspective on End-of-Life Care - an evaluation based on the German version of the General Practice End of Life Care Index. GMS German Medical Science, 2020, 18, Doc10.	2.7	1
132	Inter- and intraobserver reliability of morphological Mutch classification for greater tuberosity fractures of the proximal humerus: A comparison of x-ray, two-, and three-dimensional CT imaging. PLoS ONE, 2021, 16, e0259646.	1.1	1
133	Association of mental demands in the workplace with cognitive function in older adults at increased risk for dementia. BMC Geriatrics, 2021, 21, 688.	1.1	1
134	Risk for disorientation: Longitudinal evidence from three oldâ€age cohorts in Germany (AgeDifferent.de). Alzheimer's and Dementia, 2020, 16, e039676.	0.4	0
135	AgeWell.de (Germany). Alzheimer's and Dementia, 2020, 16, e046950.	0.4	0
136	The moderating effects of social support and depressive symptoms on pain among elderly multimorbid patientsâ€"data from the multicentre, prospective, observational cohort study MultiCare. Aging and Mental Health, 2021, , 1-7.	1.5	0