

Mieko Nakamura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6898803/publications.pdf>

Version: 2024-02-01

56
papers

1,274
citations

331670

21
h-index

377865

34
g-index

60
all docs

60
docs citations

60
times ranked

1650
citing authors

#	ARTICLE	IF	CITATIONS
1	Serum Lutein and Zeaxanthin Are Inversely Associated with High-Sensitivity C-Reactive Protein in Non-Smokers: The Mikkabi Study. <i>Antioxidants</i> , 2022, 11, 259.	5.1	4
2	Association between functional foods and cardiometabolic health in a real-life setting: a longitudinal observational study using objective diet records from an electronic purchase system. <i>Food and Function</i> , 2022, , .	4.6	1
3	Trends in Prevalence, Treatment, and Control of Hypertension According to 40-Year-Old Life Expectancy at Prefectures in Japan from the National Health and Nutrition Surveys. <i>Nutrients</i> , 2022, 14, 1219.	4.1	3
4	Dietary Changes during the COVID-19 Pandemic: A Longitudinal Study Using Objective Sequential Diet Records from an Electronic Purchase System in a Workplace Cafeteria in Japan. <i>Nutrients</i> , 2021, 13, 1606.	4.1	8
5	149Interaction of socioeconomic status on the association between guardiansâ€™ sense of coherence and childrenâ€™s caries. <i>International Journal of Epidemiology</i> , 2021, 50, .	1.9	0
6	Does the Association between Guardiansâ€™ Sense of Coherence and their Childrenâ€™s Untreated Caries Differ According to Socioeconomic Status?. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 1619.	2.6	1
7	Association between Food Spending and Dietary Intake in Workers. <i>The Japanese Journal of Nutrition and Dietetics</i> , 2020, 78, 188-197.	0.1	0
8	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , 2019, 29, 133-138.	2.4	9
9	Health Effects of Î²-Cryptoxanthin and Î²-Cryptoxanthin-Enriched Satsuma Mandarin Juice. , 2019, , 393-417.		1
10	The development and validation of the Ethical Sensitivity Questionnaire for Nursing Students. <i>BMC Medical Education</i> , 2019, 19, 215.	2.4	20
11	Low Zinc, Copper, and Manganese Intake is Associated with Depression and Anxiety Symptoms in the Japanese Working Population: Findings from the Eating Habit and Well-Being Study. <i>Nutrients</i> , 2019, 11, 847.	4.1	64
12	Dietary patterns and abnormal glucose tolerance among Japanese: findings from the National Health and Nutrition Survey, 2012. <i>Public Health Nutrition</i> , 2019, 22, 2460-2468.	2.2	13
13	Decisional Conflict in Home Medical Care in a Family-Oriented Society: Family Members' Perspectives on Surrogate Decision Making from a Multicenter Cohort Study. <i>Journal of Palliative Medicine</i> , 2019, 22, 814-822.	1.1	7
14	Association between Dietary Intake and Autistic Traits in Japanese Working Adults: Findings from the Eating Habit and Well-Being Study. <i>Nutrients</i> , 2019, 11, 3010.	4.1	5
15	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , 2019, 24, 1.	3.4	84
16	Impact of Patients' Expressed Wishes on Their Surrogate Decision Makers' Preferred Decision-Making Roles in Japan. <i>Journal of Palliative Medicine</i> , 2018, 21, 354-360.	1.1	8
17	Association between the time perspective and type of involvement in bullying among adolescents: A cross-sectional study in Japan. <i>Japan Journal of Nursing Science</i> , 2018, 15, 156-166.	1.3	5
18	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S59-S65.	2.4	11

#	ARTICLE	IF	CITATIONS
19	Dietary intake and dinner timing among shift workers in Japan. <i>Journal of Occupational Health</i> , 2018, 60, 467-474.	2.1	11
20	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , 2018, 28, S23-S28.	2.4	10
21	Dietary patterns among Japanese adults: findings from the National Health and Nutrition Survey, 2012. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2018, 27, 1120-1130.	0.4	16
22	Promotion of Salt Reduction in Shizuoka Prefecture. <i>The Japanese Journal of Nutrition and Dietetics</i> , 2018, 76, 34-43.	0.1	0
23	Orthopedic, ophthalmic, and psychiatric diseases primarily affect activity limitation for Japanese males and females: Based on the Comprehensive Survey of Living Conditions. <i>Journal of Epidemiology</i> , 2017, 27, 75-79.	2.4	11
24	Association of food access and neighbor relationships with diet and underweight among community-dwelling older Japanese. <i>Journal of Epidemiology</i> , 2017, 27, 546-551.	2.4	34
25	The Influence of Speech-Language-Hearing Therapy Duration on the Degree of Improvement in Poststroke Language Impairment. <i>Rehabilitation Research and Practice</i> , 2017, 2017, 1-8.	0.6	0
26	Class Average Score for Teacher Support and Relief of Depression in Adolescents: A Population Study in Japan. <i>Journal of School Health</i> , 2016, 86, 173-180.	1.6	6
27	High serum carotenoids are associated with lower risk for developing elevated serum alanine aminotransferase among Japanese subjects: the Mikkabi cohort study. <i>British Journal of Nutrition</i> , 2016, 115, 1462-1469.	2.3	30
28	Serum β -cryptoxanthin and β -carotene derived from Satsuma mandarin and brachial artery ankle pulse wave velocity: The Mikkabi cohort study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016, 26, 808-814.	2.6	23
29	A modified DASH diet is one possible solution for overcoming the unfavorable link between vegetable and salt intake in the Japanese diet. <i>Hypertension Research</i> , 2016, 39, 756-757.	2.7	2
30	Poor Oral Health and Diet in Relation to Weight Loss, Stable Underweight, and Obesity in Community-Dwelling Older Adults: A Cross-Sectional Study From the JAGES 2010 Project. <i>Journal of Epidemiology</i> , 2016, 26, 322-329.	2.4	32
31	High Vitamin C Intake with High Serum β -Cryptoxanthin Associated with Lower Risk for Osteoporosis in Post-Menopausal Japanese Female Subjects: Mikkabi Cohort Study. <i>Journal of Nutritional Science and Vitaminology</i> , 2016, 62, 185-191.	0.6	26
32	Association of Hay Fever with the Failure of <i>Helicobacter pylori</i> Primary Eradication. <i>Internal Medicine</i> , 2016, 55, 1729-1734.	0.7	3
33	Characteristics of pneumonia deaths after an earthquake and tsunami: an ecological study of 5.7 million participants in 131 municipalities, Japan. <i>BMJ Open</i> , 2016, 6, e009190.	1.9	17
34	High serum carotenoids associated with lower risk for the metabolic syndrome and its components among Japanese subjects: Mikkabi cohort study. <i>British Journal of Nutrition</i> , 2015, 114, 1674-1682.	2.3	40
35	High-serum carotenoids associated with lower risk for developing type 2 diabetes among Japanese subjects: Mikkabi cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2015, 3, e000147.	2.8	65
36	Is high job control a risk factor for poor quality of life in workers with high autism spectrum tendencies? A cross-sectional survey at a factory in Japan. <i>Journal of Occupational Health</i> , 2015, 57, 419-426.	2.1	5

#	ARTICLE	IF	CITATIONS
37	Development of a Short Food Frequency Questionnaire for Nutritional Assessment of People Living in a Regional Community and Determination of Its Validity and Reproducibility. The Japanese Journal of Nutrition and Dietetics, 2015, 73, 182-194.	0.1	7
38	Weather and headache onset: a large-scale study of headache medicine purchases. International Journal of Biometeorology, 2015, 59, 447-451.	3.0	20
39	Postpartum smoking relapse among women who quit during pregnancy: Cross-sectional study in Japan. Journal of Obstetrics and Gynaecology Research, 2013, 39, 1505-1512.	1.3	21
40	Consumption of Dairy Products and Death From Cardiovascular Disease in the Japanese General Population: The NIPPON DATA80. Journal of Epidemiology, 2013, 23, 47-54.	2.4	51
41	Occupational Factors and Problem Drinking among a Japanese Working Population. Industrial Health, 2013, 51, 490-500.	1.0	12
42	Dietary Reference Intakes for Japanese 2010: Lifestage. Journal of Nutritional Science and Vitaminology, 2012, 59, S103-S109.	0.6	7
43	High Serum Carotenoids Associated with Lower Risk for Bone Loss and Osteoporosis in Post-Menopausal Japanese Female Subjects: Prospective Cohort Study. PLoS ONE, 2012, 7, e52643.	2.5	91
44	Association between Food Group Intake and Serum Total Cholesterol in the Japanese Population: NIPPON DATA 80/90. Journal of Epidemiology, 2010, 20, S576-S581.	2.4	14
45	Synergistic interaction of cigarette smoking and alcohol drinking with serum carotenoid concentrations: findings from a middle-aged Japanese population. British Journal of Nutrition, 2009, 102, 1211-1219.	2.3	41
46	Associations of serum carotenoid concentrations with the metabolic syndrome: interaction with smoking. British Journal of Nutrition, 2008, 100, 1297-1306.	2.3	62
47	High β -carotene and β -cryptoxanthin are associated with low pulse wave velocity. Atherosclerosis, 2006, 184, 363-369.	0.8	46
48	Serum carotenoid concentrations are inversely associated with serum aminotransferases in hyperglycemic subjects. Diabetes Research and Clinical Practice, 2006, 71, 82-91.	2.8	34
49	Dietary Supplement Use by Community-living Population in Japan: Data from the National Institute for Longevity Sciences Longitudinal Study of Aging (NILS-LSA). Journal of Epidemiology, 2006, 16, 249-260.	2.4	52
50	The Homeostasis Model Assessment-insulin Resistance Index Is Inversely Associated with Serum Carotenoids in Non-diabetic Subjects. Journal of Epidemiology, 2006, 16, 71-78.	2.4	80
51	High Serum Carotenoids Are Inversely Associated with Serum Gamma-glutamyltransferase in Alcohol Drinkers within Normal Liver Function. Journal of Epidemiology, 2005, 15, 180-186.	2.4	68
52	A nationwide epidemiological survey regarding sudden deafness in 2001-factors influencing its prognosis-. Audiology Japan, 2004, 47, 109-118.	0.1	8
53	Feasibility and Effect on Blood Pressure of 6-Week Trial of Low Sodium Soy Sauce and Miso (Fermented) Tj ETQq1 1.0.784314 rgBT /Cv	1.6	32
54	Nutrient Intake in Japanese Adults. From the National Nutrition Survey, 1995-99.. Journal of Nutritional Science and Vitaminology, 2002, 48, 433-441.	0.6	22

#	ARTICLE	IF	CITATIONS
55	Japanese and Western diet and risk of idiopathic sudden deafness: a case-control study using pooled controls. <i>International Journal of Epidemiology</i> , 2001, 30, 608-615.	1.9	30
56	Cross-sectional Study on Serum Thiobarbituric Acid Reactive Substances(TBARS) in Akabane, Japan, with Special Reference to Dietary Habits, Smoking and Obesity.. <i>The Japanese Journal of Nutrition and Dietetics</i> , 1993, 51, 221-233.	0.1	0