

# Ali Sarafraz-Yazdi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6901962/publications.pdf>

Version: 2024-02-01

68  
papers

3,175  
citations

172457

29  
h-index

155660

55  
g-index

70  
all docs

70  
docs citations

70  
times ranked

2730  
citing authors

#	ARTICLE	IF	CITATIONS
1	Liquid-phase microextraction. <i>TrAC - Trends in Analytical Chemistry</i> , 2010, 29, 1-14.	11.4	654
2	Application of molecularly-imprinted polymers in solid-phase microextraction techniques. <i>TrAC - Trends in Analytical Chemistry</i> , 2015, 73, 81-90.	11.4	153
3	Separation and determination of amitriptyline and nortriptyline by dispersive liquid-liquid microextraction combined with gas chromatography flame ionization detection. <i>Talanta</i> , 2008, 75, 1293-1299.	5.5	138
4	Surfactant-based extraction methods. <i>TrAC - Trends in Analytical Chemistry</i> , 2011, 30, 918-929.	11.4	138
5	Determination of non-steroidal anti-inflammatory drugs in water samples by solid-phase microextraction based sol-gel technique using poly(ethylene glycol) grafted multi-walled carbon nanotubes coated fiber. <i>Analytica Chimica Acta</i> , 2012, 720, 134-141.	5.4	105
6	Surfactant enhanced liquid-phase microextraction of basic drugs of abuse in hair combined with high performance liquid chromatography. <i>Journal of Chromatography A</i> , 2005, 1094, 1-8.	3.7	98
7	Two-step hollow fiber-based, liquid-phase microextraction combined with high-performance liquid chromatography: A new approach to determination of aromatic amines in water. <i>Journal of Chromatography A</i> , 2005, 1082, 136-142.	3.7	95
8	A novel solid-phase microextraction using coated fiber based sol-gel technique using poly(ethylene glycol) and o-xylene in water samples with gas chromatography-flame ionization detector. <i>Journal of Chromatography A</i> , 2011, 1218, 5757-5764.	3.7	86
9	A combination of ultrasound and inorganic catalyst: removal of 2-chlorophenol from aqueous solution. <i>Ultrasonics Sonochemistry</i> , 2005, 12, 137-141.	8.2	85
10	Separation and determination of benzene, toluene, ethylbenzene and o-xylene compounds in water using directly suspended droplet microextraction coupled with gas chromatography-flame ionization detector. <i>Talanta</i> , 2009, 78, 936-941.	5.5	84
11	BTEX determination in water matrices using HF-LPME with gas chromatography-flame ionization detector. <i>Chemosphere</i> , 2008, 71, 671-676.	8.2	80
12	Design, synthesis and evaluation of a molecularly imprinted polymer for hollow fiber-based solid phase microextraction of chlorogenic acid in medicinal plants. <i>Journal of Chromatography A</i> , 2012, 1229, 24-29.	3.7	69
13	Liquid-liquid-liquid phase microextraction of aromatic amines in water using crown ethers by high-performance liquid chromatography with monolithic column. <i>Talanta</i> , 2005, 66, 664-669.	5.5	67
14	A solid phase microextraction coating based on ionic liquid sol-gel technique for determination of benzene, toluene, ethylbenzene and o-xylene in water samples using gas chromatography flame ionization detector. <i>Journal of Chromatography A</i> , 2013, 1300, 104-111.	3.7	59
15	Determination of non-steroidal anti-inflammatory drugs in urine by hollow-fiber liquid membrane-protected solid-phase microextraction based on sol-gel fiber coating. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 908, 67-75.	2.3	55
16	Determination of 3-nitroaniline in water samples by directly suspended droplet three-phase liquid-phase microextraction using 18-crown-6 ether and high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 5086-5091.	3.7	53
17	Carbon nanofibers decorated with magnetic nanoparticles as a new sorbent for the magnetic solid phase extraction of selected polycyclic aromatic hydrocarbons from water samples. <i>New Journal of Chemistry</i> , 2015, 39, 5621-5627.	2.8	47
18	Comparison of Hollow Fiber and Single-Drop Liquid-Phase Microextraction Techniques for HPLC Determination of Aniline Derivatives in Water. <i>Chromatographia</i> , 2006, 63, 563-569.	1.3	45

#	ARTICLE	IF	CITATIONS
19	A combination of ultrasound and a bio-catalyst: removal of 2-chlorophenol from aqueous solution. <i>Ultrasonics Sonochemistry</i> , 2006, 13, 37-41.	8.2	43
20	Determination of furan in food samples using two solid phase microextraction fibers based on sol-gel technique with gas chromatography-flame ionisation detector. <i>Food Chemistry</i> , 2012, 131, 698-704.	8.2	43
21	Modeling of solid-phase tea waste extraction for the removal of manganese and cobalt from water samples by using PSO-artificial neural network and response surface methodology. <i>Arabian Journal of Chemistry</i> , 2017, 10, S1663-S1673.	4.9	43
22	Comparative study of the three sol-gel based solid phase microextraction fibers in extraction of BTEX from water samples using gas chromatography-flame ionization detection. <i>Analytical Methods</i> , 2010, 2, 746.	2.7	40
23	A new high-speed hollow fiber based liquid phase microextraction method using volatile organic solvent for determination of aromatic amines in environmental water samples prior to high-performance liquid chromatography. <i>Talanta</i> , 2009, 79, 472-478.	5.5	38
24	Directly Suspended Droplet Microextraction and Analysis of Amitriptyline and Nortriptyline by GC. <i>Chromatographia</i> , 2007, 66, 613-617.	1.3	37
25	Comparative study of the sol-gel based solid phase microextraction fibers in extraction of naphthalene, fluorene, anthracene and phenanthrene from saffron samples extractants. <i>Mikrochimica Acta</i> , 2012, 176, 317-325.	5.0	37
26	Determination of monocyclic aromatic amines using headspace solid-phase microextraction based on sol-gel technique prior to GC. <i>Journal of Separation Science</i> , 2013, 36, 1629-1635.	2.5	36
27	Pre-concentration of non-steroidal anti-inflammatory drugs in water using dispersive liquid-liquid and single-drop microextraction with high-performance liquid chromatography. <i>Journal of Separation Science</i> , 2012, 35, 2476-2483.	2.5	35
28	Determination of Triazole Fungicides Using Hollow Fiber Liquid Phase Microextraction Prior to Gas Chromatography-Mass Spectrometry Analysis. <i>Industrial &amp; Engineering Chemistry Research</i> , 2012, 51, 3101-3107.	3.7	34
29	Application of Sol-Gel Based Poly(ethylene glycol)/Multiwalled Carbon Nanotubes Coated Fiber for SPME of Methyl tert-Butyl Ether in Environmental Water Samples. <i>Chromatographia</i> , 2010, 72, 923-931.	1.3	33
30	Headspace solid phase microextraction of volatile aromatic hydrocarbons using a steel wire coated with an electrochemically prepared nanocomposite consisting of polypyrrole, carbon nanotubes, and titanium oxide. <i>Mikrochimica Acta</i> , 2015, 182, 217-225.	5.0	31
31	Determination of volatile organic compounds in environmental water samples using three solid-phase microextraction fibers based on sol-gel technique with gas chromatography-flame ionization detector. <i>Analytical Methods</i> , 2011, 3, 1877.	2.7	30
32	Determination of melamine in soil samples using surfactant-enhanced hollow fiber liquid phase microextraction followed by HPLC-UV using experimental design. <i>Journal of Advanced Research</i> , 2015, 6, 957-966.	9.5	29
33	Magnetic solid-phase extraction of polycyclic aromatic hydrocarbons in water samples by Fe <sub>3</sub> O <sub>4</sub> @polypyrrole/carbon nanotubes. <i>Journal of Separation Science</i> , 2016, 39, 2746-2753.	2.5	29
34	New application of chitosan-grafted polyaniline in dispersive solid-phase extraction for the separation and determination of phthalate esters in milk using high-performance liquid chromatography. <i>Journal of Separation Science</i> , 2017, 40, 1739-1746.	2.5	29
35	Directly Suspended Droplet Three Liquid Phase Microextraction of Diclofenac Prior to LC. <i>Chromatographia</i> , 2008, 67, 49-53.	1.3	25
36	Determination of phenolic compounds in water and urine samples using solid-phase microextraction based on sol-gel technique prior to GC-FID. <i>Analytical Methods</i> , 2012, 4, 4316.	2.7	25

#	ARTICLE	IF	CITATIONS
37	Ionic Liquid-Based Submerged Single Drop Microextraction: a New Method for the Determination of Aromatic Amines in Environmental Water Samples. <i>Chromatographia</i> , 2010, 72, 867-873.	1.3	24
38	Determination of Trace of Methyl tert-Butyl Ether in Water Using Liquid Drop Headspace Sampling and GC. <i>Chromatographia</i> , 2004, 60, 699-702.	1.3	23
39	Determination of Hg(II) in Natural Waters by Diphenylation by Single-Drop Microextraction: GC. <i>Chromatographia</i> , 2010, 71, 1049-1054.	1.3	21
40	LC Determination of Mono-Substituted Phenols in Water Using Liquid-Liquid Phase Microextraction. <i>Chromatographia</i> , 2005, 62, 49-54.	1.3	20
41	Capillary gas chromatographic determination of copper and nickel using microwave-induced plasma atomic emission detection. <i>Journal of Chromatography A</i> , 1993, 636, 271-276.	3.7	19
42	Title is missing!. <i>Journal of Inclusion Phenomena and Macrocyclic Chemistry</i> , 2002, 43, 231-237.	1.6	18
43	Comparative study of direct immersion and headspace single drop microextraction techniques for BTEX determination in water samples using GC-FID. <i>International Journal of Environmental Analytical Chemistry</i> , 2010, 90, 1036-1047.	3.3	18
44	Headspace solid-phase microextraction using poly(ethylene glycol) grafted multi-walled carbon nanotube fibers for the determination of methyl tert-butyl ether in water samples. <i>Analytical Methods</i> , 2012, 4, 3701.	2.7	18
45	Determination of Aromatic Amines Using Solid-Phase Microextraction Based on an Ionic Liquid-Mediated Sol-Gel Technique. <i>Journal of Chromatographic Science</i> , 2016, 54, 677-681.	1.4	18
46	Development of novel magnetic solid-phase extraction sorbent based on Fe <sub>3</sub> O <sub>4</sub> /carbon nanosphere/polypyrrole composite and their application to the enrichment of polycyclic aromatic hydrocarbons from water samples prior to GC-FID analysis. <i>Journal of the Iranian Chemical Society</i> , 2018, 15, 153-161.	2.2	18
47	Preparation of an ionic liquid-mediated carbon nanotube-poly(dimethylsiloxane) fiber by sol-gel technique for determination of polycyclic aromatic hydrocarbons in urine samples using head-space solid-phase microextraction coupled with gas chromatography. <i>Journal of the Iranian Chemical Society</i> , 2014, 11, 969-977.	2.2	17
48	New polypyrrole-carbon nanotubes-silicon dioxide solid-phase microextraction fiber for the preconcentration and determination of benzene, toluene, ethylbenzene, and <i>o</i> -xylene using gas liquid chromatography. <i>Journal of Separation Science</i> , 2014, 37, 2605-2612.	2.5	17
49	Ionic-liquid-mediated poly(dimethylsiloxane)- grafted carbon nanotube fiber prepared by the sol-gel technique for the head space solid-phase microextraction of methyl <i>tert</i> -butyl ether using GC. <i>Journal of Separation Science</i> , 2014, 37, 127-134.	2.5	16
50	Development of a poly(ethylene glycol)-graphene oxide sol-gel coating for solid-phase microextraction of aromatic amines in water samples with a gas chromatography-flame ionization detector method. <i>New Journal of Chemistry</i> , 2015, 39, 1287-1294.	2.8	15
51	Effect of cathodic electrolyte on the performance of electrochemical hydride generation from graphite cathode. <i>Talanta</i> , 2004, 64, 644-649.	5.5	14
52	Separation and Determination of Amitriptyline and Nortriptyline in Biological Samples Using Single-Drop Microextraction with GC. <i>Chromatographia</i> , 2011, 73, 549-557.	1.3	14
53	Gas chromatography of copper (II), nickel (II), palladium (II) and vanadium (IV) using atomic emission detector and $\beta$ -ketoamine Schiff bases derived from stilbenediamines. <i>Chromatographia</i> , 2002, 56, 729-732.	1.3	13
54	PRE-CONCENTRATION AND DETERMINATION OF $\beta$ -BLOCKERS USING CARBON NANOTUBE-ASSISTED PSEUDO-STIRBAR HOLLOW FIBER SOLID-LIQUID-PHASE MICROEXTRACTION AND HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY WITH FLUORESCENCE DETECTION. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2013, 36, 750-769.	1.0	13

#	ARTICLE	IF	CITATIONS
55	Determination of Mercury in Real Water Samples Using in situ Derivatization Followed by Sol-Gel-Solid-Phase Microextraction with Gas Chromatography-Flame Ionization Detection. <i>Journal of Chromatographic Science</i> , 2014, 52, 81-87.	1.4	13
56	Polyethylene glycol grafted flower-like cupric nano oxide for the hollow fiber solid phase microextraction of hexaconazole, penconazole, and diniconazole in vegetable samples. <i>Journal of Separation Science</i> , 2016, 39, 3137-3144.	2.5	12
57	Gas chromatography of chloride and bromide as phenylboronic acid/mercuric nitrate derivatives with microwave induced plasma atomic emission detection. <i>Journal of Chromatography A</i> , 1992, 594, 395-399.	3.7	11
58	Platinum-selective capillary gas chromatographic determination with microwave-induced plasma atomic emission detection. <i>Journal of Chromatography A</i> , 1998, 824, 223-229.	3.7	11
59	A non-ionic surfactant-mediated sol-gel coating for solid-phase microextraction of benzene, toluene, ethylbenzene and o-xylene in water samples using a gas chromatography-flame ionization detector. <i>New Journal of Chemistry</i> , 2014, 38, 4486.	2.8	11
60	Membrane extraction with sorbent interface-gas chromatography as an effective and fast means for continuous monitoring of thermal degradation products of polyacrylonitrile. <i>Analyst, The</i> , 2002, 127, 912-916.	3.5	9
61	An insight into the determination of trace levels of benzodiazepines in biometric systems: Use of crab shell powder as an environmentally friendly biosorbent. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2018, 1092, 58-64.	2.3	9
62	Simultaneous derivatization and extraction of iodine from milk samples by hollow fiber liquid-phase microextraction followed by gas chromatography-electron capture detection. <i>Journal of the Iranian Chemical Society</i> , 2013, 10, 643-651.	2.2	8
63	SIMULTANEOUS DETERMINATION OF VITAMIN A AND E IN INFANT MILK FORMULAS USING SEMI-MICRO LIQUID-LIQUID EXTRACTION FOLLOWED BY HPLC-UV. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2014, 37, 391-403.	1.0	8
64	Determination of chlorophenols in environmental water samples using directly suspended droplet liquid-liquid-liquid phase microextraction prior to high-performance liquid chromatography. <i>International Journal of Environmental Analytical Chemistry</i> , 2010, 90, 1108-1118.	3.3	6
65	Determination of Hg(II) in Environmental Water Samples Using DLLME Method Prior to GC-FID. <i>Chromatographia</i> , 2013, 76, 861-865.	1.3	5
66	Carbon nanotube assisted sol-gel based hollow fiber solidphase microextraction combined with pre-heating injectionhigh performance liquid chromatography as a novel sample preparation method for determination of nitroaromatics. <i>Sample Preparation</i> , 2013, 1, 1-9.	0.4	5
67	Combination of artificial neural network and genetic algorithm method for modeling of methylene blue adsorption onto wood sawdust from water samples. <i>Toxicology and Industrial Health</i> , 2016, 32, 437-446.	1.4	5
68	Fluorine-tin oxide (FTO) electrode modified with platinum nanoparticles dispersed into montmorillonite clay as an effective and low cost catalyst for ethanol electrooxidation. <i>RSC Advances</i> , 2016, 6, 113240-113248.	3.6	4