

Roco Gallego-Durn

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6918189/rocio-gallego-duran-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

34
papers

618
citations

16
h-index

24
g-index

39
ext. papers

962
ext. citations

4.7
avg, IF

3.8
L-index

#	Paper	IF	Citations
34	Long non-coding RNA H19 as biomarker for hepatocellular carcinoma.. <i>Liver International</i> , 2022 ,	7.9	2
33	Liver injury in non-alcoholic fatty liver disease is associated with urea cycle enzyme dysregulation.. <i>Scientific Reports</i> , 2022 , 12, 3418	4.9	1
32	Systematic review and meta-analysis: analysis of variables influencing the interpretation of clinical trial results in NAFLD.. <i>Journal of Gastroenterology</i> , 2022 , 57, 357	6.9	1
31	Development and Validation of a Clinical-Genetic Risk Score to Predict Hepatic Encephalopathy in Patients With Liver Cirrhosis. <i>American Journal of Gastroenterology</i> , 2021 , 116, 1238-1247	0.7	2
30	Extra virgin olive oil improved body weight and insulin sensitivity in high fat diet-induced obese LDLr-/-Leiden mice without attenuation of steatohepatitis. <i>Scientific Reports</i> , 2021 , 11, 8250	4.9	3
29	Definite and indeterminate nonalcoholic steatohepatitis share similar clinical features and prognosis: A longitudinal study of 1893 biopsy-proven nonalcoholic fatty liver disease subjects. <i>Liver International</i> , 2021 , 41, 2076-2086	7.9	5
28	Management of NAFLD patients with advanced fibrosis. <i>Liver International</i> , 2021 , 41 Suppl 1, 95-104	7.9	3
27	Caucasian lean subjects with non-alcoholic fatty liver disease share long-term prognosis of non-lean: time for reappraisal of BMI-driven approach?. <i>Gut</i> , 2021 ,	19.2	27
26	Copy number variation and expression of exportin-4 associates with severity of fibrosis in metabolic associated fatty liver disease. <i>EBioMedicine</i> , 2021 , 70, 103521	8.8	5
25	Co-expression gene network analysis reveals novel regulatory pathways involved in porto-sinusoidal vascular disease. <i>Journal of Hepatology</i> , 2021 , 75, 924-934	13.4	2
24	Long-term outcomes and predictive ability of non-invasive scoring systems in patients with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2021 , 75, 786-794	13.4	21
23	Bacterial antigen translocation and age as BMI-independent contributing factors on systemic inflammation in NAFLD patients. <i>Liver International</i> , 2020 , 40, 2182-2193	7.9	6
22	Significant fibrosis predicts new-onset diabetes mellitus and arterial hypertension in patients with NASH. <i>Journal of Hepatology</i> , 2020 , 73, 17-25	13.4	26
21	Genetic variation in the TLL1 gene is not associated with fibrosis in patients with metabolic associated fatty liver disease. <i>PLoS ONE</i> , 2020 , 15, e0243590	3.7	0
20	Hepatitis C Virus Clearance by Direct-Acting Antivirals Agents Improves Endothelial Dysfunction and Subclinical Atherosclerosis: HEP-CAR Study. <i>Clinical and Translational Gastroenterology</i> , 2020 , 11, e00203	4.2	9
19	Development and Validation of Hepamet Fibrosis Scoring System-A Simple, Noninvasive Test to Identify Patients With Nonalcoholic Fatty Liver Disease With Advanced Fibrosis. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 216-225.e5	6.9	46
18	A variant in the MICA gene is associated with liver fibrosis progression in chronic hepatitis C through TGF- β dependent mechanisms. <i>Scientific Reports</i> , 2019 , 9, 1439	4.9	6

17	Metabolic characterization of two different non-alcoholic fatty liver disease pre-clinical mouse models. <i>Revista Espanola De Enfermedades Digestivas</i> , 2019 , 111, 301-307	0.9	2
16	Analysis of the burden and variability in the management of NAFLD patients in the clinical practice: unifying the required criteria. <i>Revista Espanola De Enfermedades Digestivas</i> , 2019 , 111, 270-274	0.9	2
15	Impaired brain glymphatic flow in experimental hepatic encephalopathy. <i>Journal of Hepatology</i> , 2019 , 70, 40-49	13.4	33
14	A polymorphism in the Irisin-encoding gene (FNDC5) associates with hepatic steatosis by differential miRNA binding to the 3'UTR. <i>Journal of Hepatology</i> , 2019 , 70, 494-500	13.4	46
13	Genetic and Epigenetic Regulation in Nonalcoholic Fatty Liver Disease (NAFLD). <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	64
12	miRNAs in patients with non-alcoholic fatty liver disease: A systematic review and meta-analysis. <i>Journal of Hepatology</i> , 2018 , 69, 1335-1348	13.4	73
11	Simvastatin and metformin inhibit cell growth in hepatitis C virus infected cells via mTOR increasing PTEN and autophagy. <i>PLoS ONE</i> , 2018 , 13, e0191805	3.7	18
10	Metformin modifies glutamine metabolism in an in vitro and in vivo model of hepatic encephalopathy. <i>Revista Espanola De Enfermedades Digestivas</i> , 2018 , 110, 427-433	0.9	3
9	Natural Extracts Abolished Lipid Accumulation in Cells Harboring non-favourable PNPLA3 genotype. <i>Annals of Hepatology</i> , 2018 , 17, 242-249	3.1	9
8	The effects of metabolic status on non-alcoholic fatty liver disease-related outcomes, beyond the presence of obesity. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 48, 1260-1270	6.1	43
7	An extra virgin olive oil rich diet intervention ameliorates the nonalcoholic steatohepatitis induced by a high-fat "Western-type" diet in mice. <i>Molecular Nutrition and Food Research</i> , 2017 , 61, 1600549	5.9	33
6	Lactulose reduces bacterial DNA translocation, which worsens neurocognitive shape in cirrhotic patients with minimal hepatic encephalopathy. <i>Liver International</i> , 2017 , 37, 212-223	7.9	17
5	Imaging biomarkers for steatohepatitis and fibrosis detection in non-alcoholic fatty liver disease. <i>Scientific Reports</i> , 2016 , 6, 31421	4.9	22
4	Oxidized low-density lipoprotein antibodies/high-density lipoprotein cholesterol ratio is linked to advanced non-alcoholic fatty liver disease lean patients. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2016 , 31, 1611-8	4	17
3	Epigenetic mechanisms in non-alcoholic fatty liver disease: An emerging field. <i>World Journal of Hepatology</i> , 2015 , 7, 2497-502	3.4	32
2	Increased soluble CD36 is linked to advanced steatosis in nonalcoholic fatty liver disease. <i>European Journal of Clinical Investigation</i> , 2014 , 44, 65-73	4.6	31
1	Thalidomide with peginterferon alfa-2b and ribavirin in the treatment of non-responders genotype 1 chronic hepatitis C patients: proof of concept. <i>Revista Espanola De Enfermedades Digestivas</i> , 2011 , 103, 619-25	0.9	3